

Practice for Test 3, MAT 266

Determine whether the following series converges absolutely. Justify your answer with the proper series test.

$$1. \sum_{n=0}^{\infty} \frac{(-1)^n (n+5)!}{5!n!4^n} \quad 2. \sum_{n=1}^{\infty} \frac{(-2)^{n+3} n^3}{3^n} \quad 3. \sum_{n=0}^{\infty} \frac{n!}{1,000,000^n} \quad 4. \sum_{n=0}^{\infty} \frac{(2n)!}{3^n (n!)^2}$$

Determine the radius of convergence and the largest open interval of convergence. Justify your answer with the proper series test.

$$5. \sum_{n=0}^{\infty} \frac{(-1)^n 6^n x^{2n}}{n!} \quad 6. \sum_{n=0}^{\infty} \frac{5^n x^n (n+4)}{(n^2 + 16)} \quad 7. \sum_{n=1}^{\infty} \frac{n^2 (x+4)^n}{7^n \sqrt{2n-1}} \quad 8. \sum_{n=1}^{\infty} \frac{(-1)^n n (x-5)^n}{3^{2n} (n+1)}$$

9. For the function $f(x) = \frac{3}{16+x^2}$

(a) Find the power series representation of $f(x)$.

(b) Determine the radius and interval of convergence for the power series in part (a).

10. For the function $f(x) = \frac{1}{6+x}$

(a) Find the power series representation of $f(x)$.

(b) Determine the radius and interval of convergence for the power series in part (a).

(c) Use part (a) to find the power series representation of $g(x) = \ln(6+x)$ along with its radius of convergence.

11. The function $f(x) = \frac{x^2}{1-2x}$ is represented by the power series in x , $f(x) = \sum_{n=0}^{\infty} c_n x^n$. Find the coefficients: $c_0, c_1, c_2, c_3, c_4, c_5$, and c_6 .

12. Write a partial sum for the power series in x which represents the function $f(x) = x \ln(1+3x)$ consisting of the first four nonzero terms. State the radius of convergence.

13. For the function $f(x) = x \cos(x^3)$

a. Find the partial sum of the Maclaurin series for $f(x)$ consisting of the first four nonzero terms.

b. Use the answer in part (a) to find the first four nonzero terms of the partial sum of the Maclaurin series for $F(x) = \int x \cos(x^3) dx$ with $F(0) = 0$.

c. Use the above to estimate the definite integral $\int_0^1 x \cos(x^3) dx$. Round your answer to 8 decimal places.

14. Let $F(x) = \int e^{-x^2} dx$ with $F(0) = 0$.

a. Find the partial sum of the Maclaurin series for $F(x)$ consisting of the first four nonzero terms.

b. Use the answer in part (a) to estimate the definite integral $\int_0^{1/2} e^{-x^2} dx$. Round your answer to 6 decimal places.

15. The function $f(x) = \sin\left(\frac{2x}{3}\right)$ has a Maclaurin series. Find a formula for the Maclaurin series.

16. The function $f(x) = e^{-2x}$ has a Maclaurin series. Find the first four terms of the series.

17. Find the first four terms of the Taylor series of $f(x) = \sqrt{x}$ centered at $a = 16$. Use that result to approximate $\sqrt{15.2}$.

18. Find the first four terms of the Taylor series of $f(x) = \sqrt[3]{x}$ centered at $a = 8$. Use that result to approximate $\sqrt[3]{9}$.
19. Find the first four terms of the Taylor series of $f(x) = \ln(x)$ centered at $a = 5$.
20. Find the first four terms of the Taylor series of $f(x) = \cos(x)$ centered at $a = \frac{\pi}{3}$.
21. Use Maclaurin series to find the following limits:
- a. $\lim_{x \rightarrow 0} \frac{\sin(x^2) - x^2}{x^6}$ b. $\lim_{x \rightarrow 0} \frac{x(\tan^{-1}(x) - x)}{\cos(x) - e^{-x^2/2}}$
22. Eliminate the parameter to find the Cartesian equation for $x = 1 + t$, $y = t^2 + 2$.
23. Eliminate the parameter to find the Cartesian equation for $x = 2\sin(t)$, $y = 3\cos(t)$.
24. Eliminate the parameter to find the Cartesian equation for $x = \sec^2(t)$, $y = 2\tan(t)$.
25. Eliminate the parameter to find the Cartesian equation for $x = 1 + 2\cos(t)$, $y = -2 + 2\sin(t)$.
26. Suppose parametric equations for the line segment between $(1, 2)$ and $(4, 7)$ have the form $x = a + bt$, $y = c + dt$. Find a, b, c and d so that the parametric curve starts at $(1, 2)$ when $t = 0$, and ends at $(4, 7)$ when $t = 1$.
27. Find parametric equations for the path of a particle that moves along the circle $x^2 + y^2 = 25$ twice around clockwise, $0 \leq t \leq 2\pi$, starting at $(5, 0)$.
28. Find the slope of the line tangent to the parametric curve $x = t + \cos(\pi t)$, $y = t + \sin(\pi t)$ at $t = \frac{1}{2}$.
29. Find an equation of the line tangent to the parametric curve $x = \cos^3(t)$, $y = \sin^3(t)$, at $t = \frac{\pi}{3}$.
30. For the parametric curve $x = \frac{t^3}{9} - t$, $y = \frac{t^4}{12} - t^2 + 1$
- a. Find an equation of the line tangent to the curve at $t = 3$.
- b. Find all values of t where the tangent is horizontal or vertical.
31. For the parametric curve $x = e^{2t} - t - 3$, $y = -t^3 + 2t$
- a. Find an equation of the line tangent to the curve at $t = 0$.
- b. Find the exact value of t corresponding to the leftmost point of the curve, and the value(s) of t where the tangent is horizontal.
32. Find the arc length of the parametric curve $x = \cos^3(t)$, $y = \sin^3(t)$, $0 \leq t \leq \frac{\pi}{2}$.
33. Find the exact arc length of the loop of the parametric curve, $x = t - \frac{t^3}{3}$, $y = t^2$, $-\sqrt{3} \leq t \leq \sqrt{3}$.
34. Find the exact arc length of the parametric curve $x = \cos(t) + t\sin(t)$, $y = \sin(t) - t\cos(t)$, $0 \leq t \leq \pi$.
35. Find area under the parametric curve $x = t^3 - 1$, $y = 2 + 3t - t^3$, $-1 \leq t \leq 2$.
36. Find area under the parametric curve $x = t - \frac{1}{t}$, $y = t^2 + 1$, $\frac{1}{2} \leq t \leq 2$.
37. Find area under the parametric curve $x = \sin(t)$, $y = 2\sin(t)\cos(t)$, $0 \leq t \leq \frac{\pi}{2}$.

ANSWER KEY:

1. $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{4}$, thus, by the Ratio Test, the series Converges Absolutely.

2. $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{2}{3}$, thus, by the Ratio Test, the series Converges Absolutely.

3. $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty$, thus, by the Ratio Test, the series diverges.

4. $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{4}{3}$, thus, by the Ratio Test, the series diverges.

5. $I = (-\infty, \infty)$; $R = \infty$.

6. $I = \left(-\frac{1}{5}, \frac{1}{5}\right)$; $R = \frac{1}{5}$.

7. $I = (-11, 3)$; $R = 7$.

8. $I = (-4, 14)$; $R = 9$.

9. (a) $f(x) = \frac{3}{16} - \frac{3x^2}{256} + \frac{3x^4}{4096} - \frac{3x^6}{65536} + \cdots + \frac{(-1)^n 3(x)^{2n}}{16^{n+1}} + \cdots = \sum_{n=0}^{\infty} \frac{(-1)^n 3(x)^{2n}}{16^{n+1}}$

(b) $R = 4$, $I = (-4, 4)$

10. (a) $f(x) = \frac{1}{6} - \frac{x}{36} + \frac{x^2}{216} - \frac{x^3}{1296} + \cdots + \frac{(-1)^n x^n}{6^{n+1}} + \cdots = \sum_{n=0}^{\infty} \frac{(-1)^n x^n}{6^{n+1}}$

Note: for $f(x)$, $c_0 = \frac{1}{6}$, $c_1 = -\frac{1}{36}$, $c_2 = \frac{1}{216}$, and $c_3 = -\frac{1}{1296}$.

(b) $R = 6$, $I = (-6, 6)$

(c) $g(x) = \ln(6) + \frac{x}{6} - \frac{x^2}{72} + \frac{x^3}{648} - \frac{x^4}{5184} + \cdots + \frac{(-1)^n x^{n+1}}{(n+1)6^{n+1}} + \cdots = \ln(6) + \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^n}{n6^n}$; $R = 6$

Note: for $g(x)$, $c_0 = \ln(6)$, $c_1 = \frac{1}{6}$, $c_2 = -\frac{1}{72}$, $c_3 = \frac{1}{648}$, and $c_4 = \frac{1}{5184}$.

11. $c_0 = 0$, $c_1 = 0$, $c_2 = 1$, $c_3 = 2$, $c_4 = 4$, $c_5 = 8$, and $c_6 = 16$.

12. $f(x) \approx 3x^2 - \frac{9x^3}{2} + 9x^4 - \frac{81x^5}{4}$; $R = \frac{1}{3}$

13. (a) $f(x) \approx x - \frac{x^7}{2} + \frac{x^{13}}{24} - \frac{x^{19}}{720}$ (b) $F(x) \approx \frac{x^2}{2} - \frac{x^8}{16} + \frac{x^{14}}{336} - \frac{x^{20}}{14400}$ (c) 0.44040675

14. (a) $F(x) \approx x - \frac{x^3}{3} + \frac{x^5}{10} - \frac{x^7}{42}$ (b) 0.461272

15. $\sum_{n=0}^{\infty} \frac{(-1)^n (2x)^{2n+1}}{3^{2n+1} (2n+1)!}$

16. $f(x) \approx 1 - 2x + \frac{2^2}{2!} x^2 - \frac{2^3}{3!} x^3$

17. $f(x) \approx 4 + \frac{1}{8}(x-16) - \frac{1}{512}(x-16)^2 + \frac{1}{16384}(x-16)^3$; $\sqrt{15.2} = f(15.2) \approx 3.89871875$

$$18. f(x) \approx 2 + \frac{1}{12}(x-8) - \frac{1}{288}(x-8)^2 + \frac{5}{20736}(x-8)^3 ; \quad \sqrt[3]{9} = f(9) \approx 2.080102$$

$$19. f(x) \approx \ln(5) + \frac{1}{5}(x-5) - \frac{1}{50}(x-5)^2 + \frac{1}{375}(x-5)^3$$

$$20. f(x) \approx \frac{1}{2} - \frac{\sqrt{3}}{2}\left(x - \frac{\pi}{3}\right) - \frac{1}{4}\left(x - \frac{\pi}{3}\right)^2 + \frac{\sqrt{3}}{12}\left(x - \frac{\pi}{3}\right)^3$$

$$21. (a) -\frac{1}{6} \quad (b) 4 \quad 22. y = x^2 - 2x + 3 \quad 23. \frac{x^2}{4} + \frac{y^2}{9} = 1$$

$$24. x = \frac{y^2}{4} + 1 \quad 25. (x-1)^2 + (y+2)^2 = 4 \quad 26. x = 1 + 3t, y = 2 + 5t$$

$$27. x = 5\cos(2t), y = -5\sin(2t) \quad 28. \frac{1}{1-\pi} \quad 29. y = -\sqrt{3}x + \frac{\sqrt{3}}{2}$$

$$30. (a) y = \frac{3}{2}x - \frac{5}{4} \quad (b) \text{Horizontal tangent at } t = -\sqrt{6}, 0, \sqrt{6}; \text{Vertical tangent at } t = -\sqrt{3}, \sqrt{3}$$

$$31. (a) y = 2x + 4 \quad (b) \text{leftmost value at } t = \frac{1}{2}\ln\left(\frac{1}{2}\right); \text{Horizontal tangent at } t = -\sqrt{\frac{2}{3}}, \sqrt{\frac{2}{3}}.$$

$$32. \frac{3}{2} \quad 33. 4\sqrt{3} \quad 34. \frac{\pi^2}{2} \quad 35. \frac{81}{4}$$

$$36. \frac{57}{8} \quad 37. \frac{2}{3}$$