

Linear Regression

Dr. Víctor Uc Cetina

Facultad de Matemáticas
Universidad Autónoma de Yucatán

`cetina@informatik.uni-hamburg.de`
`https://sites.google.com/view/victorucetina`

Content


- 1 Problem 1 - Housing Data
- 2 Least Mean Square
- 3 The Normal Equations
- 4 A Probabilistic Interpretation
- 5 Locally Weighted Linear Regression

Housing Data

Suppose we have the following housing data:


Living area (feet square)	Price (USD)
560	37
1012	79
893	76
2196	130
936	72
\vdots	\vdots

Housing Data




One Dimensional Regression Problem

Living area (x_1)	Price (y)
560	37
1012	79
893	76
2196	130
936	72
\vdots	\vdots



One Dimensional Regression Problem

Living area (x_1)	Price (y)
560	37
1012	79
893	76
2196	130
936	72
\vdots	\vdots



We are looking for something like: $h_{\theta}(\mathbf{x}) = \theta_0 + \theta_1 x_1$

Two Dimensional Regression Problem

Living area (x_1)	Bedrooms (x_2)	Price (y)
560	2	37
1012	3	79
893	3	76
2196	4	130
936	3	72
\vdots	\vdots	\vdots

Now, we are looking for something like: $h_{\theta}(\mathbf{x}) = \theta_0 + \theta_1 x_1 + \theta_2 x_2$

Two Dimensional Regression Problem

Living area (x_1)	Bedrooms (x_2)	Price (y)
560	2	37
1012	3	79
893	3	76
2196	4	130
936	3	72
\vdots	\vdots	\vdots

Now, we are looking for something like: $h_{\theta}(\mathbf{x}) = \theta_0 + \theta_1 x_1 + \theta_2 x_2$

Letting $x_0 = 1$ we have: $h(\mathbf{x}) = \sum_{j=0}^n \theta_j x_j$

Two Dimensional Regression Problem

Living area (x_1)	Bedrooms (x_2)	Price (y)
560	2	37
1012	3	79
893	3	76
2196	4	130
936	3	72
\vdots	\vdots	\vdots

Now, we are looking for something like: $h_{\theta}(\mathbf{x}) = \theta_0 + \theta_1 x_1 + \theta_2 x_2$


Letting $x_0 = 1$ we have: $h(\mathbf{x}) = \sum_{j=0}^n \theta_j x_j$

This is the dot product: $\theta^T \mathbf{x}$

Polynomial Functions

$$y = 1$$


$$y = \theta_0$$



Polynomial Functions

$$y = 2x$$


$$y = \theta_1 x$$



Polynomial Functions

$$y = 1 + 2x$$


$$y = \theta_0 + \theta_1 x$$



Polynomial Functions


$$y = 1 + 2x + 2x^2$$

$$y = \theta_0 + \theta_1 x + \theta_2 x^2$$



Polynomial Functions


$$y = 1 + 2x + 2x^2 + 2x^3$$
$$y = \theta_0 + \theta_1x + \theta_2x^2 + \theta_3x^3$$



Polynomial Functions

$$y = 0.1 - 0.2x + 0.2x^2 - 0.156x^3$$

$$y = \theta_0 + \theta_1x + \theta_2x^2 + \theta_3x^3$$



How do we pick θ ?

- One reasonable method is to pick θ such that $h(x)$ is close to y , at least for our m training examples.

How do we pick θ ?

- One reasonable method is to pick θ such that $h(x)$ is close to y , at least for our m training examples.
- We define the cost function $J(\theta) = \frac{1}{2} \sum_{i=1}^m [h_{\theta}(x^{(i)}) - y^{(i)}]^2$.

How do we pick θ ?


- One reasonable method is to pick θ such that $h(x)$ is close to y , at least for our m training examples.
- We define the cost function $J(\theta) = \frac{1}{2} \sum_{i=1}^m [h_{\theta}(x^{(i)}) - y^{(i)}]^2$.
- We can initialize randomly θ and use the gradient descent algorithm to find the θ that minimizes $J(\theta)$.

How do we pick θ ?

- One reasonable method is to pick θ such that $h(x)$ is close to y , at least for our m training examples.
- We define the cost function $J(\theta) = \frac{1}{2} \sum_{i=1}^m [h_{\theta}(x^{(i)}) - y^{(i)}]^2$.
- We can initialize randomly θ and use the gradient descent algorithm to find the θ that minimizes $J(\theta)$.
- $\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta)$.


Estimating parameters

In blue, the initial $h_{\theta}(x)$ function, with randomly generated θ 's. In black, the final $h_{\theta}(x)$ function.




Graph of the error



Plot of the error $J(\theta) = \frac{1}{2} \sum_{i=1}^m [h_{\theta}(x^{(i)}) - y^{(i)}]^2$, after each iteration of stochastic gradient descent.



Gradient Descent



Gradient Descent



Deriving the LMS Learning Rule

$$\frac{\partial}{\partial \theta_j} J(\theta) = \frac{\partial}{\partial \theta_j} \frac{1}{2} (h_{\theta}(x) - y)^2$$

Deriving the LMS Learning Rule

$$\begin{aligned}\frac{\partial}{\partial \theta_j} J(\theta) &= \frac{\partial}{\partial \theta_j} \frac{1}{2} (h_{\theta}(x) - y)^2 \\ &= 2 \cdot \frac{1}{2} (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} (h_{\theta}(x) - y)\end{aligned}$$

Deriving the LMS Learning Rule

$$\begin{aligned}\frac{\partial}{\partial \theta_j} J(\theta) &= \frac{\partial}{\partial \theta_j} \frac{1}{2} (h_{\theta}(x) - y)^2 \\ &= 2 \cdot \frac{1}{2} (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} (h_{\theta}(x) - y) \\ &= (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} \left(\sum_{i=0}^n \theta_i x_i - y \right)\end{aligned}$$

Deriving the LMS Learning Rule

$$\begin{aligned}\frac{\partial}{\partial \theta_j} J(\theta) &= \frac{\partial}{\partial \theta_j} \frac{1}{2} (h_{\theta}(x) - y)^2 \\&= 2 \cdot \frac{1}{2} (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} (h_{\theta}(x) - y) \\&= (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} \left(\sum_{i=0}^n \theta_i x_i - y \right) \\&= (h_{\theta}(x) - y) x_j\end{aligned}$$

Deriving the LMS Learning Rule

$$\begin{aligned}\frac{\partial}{\partial \theta_j} J(\theta) &= \frac{\partial}{\partial \theta_j} \frac{1}{2} (h_{\theta}(x) - y)^2 \\&= 2 \cdot \frac{1}{2} (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} (h_{\theta}(x) - y) \\&= (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} \left(\sum_{i=0}^n \theta_i x_i - y \right) \\&= (h_{\theta}(x) - y) x_j\end{aligned}$$

For a single example, the rule is:

$$\theta_j := \theta_j + \alpha [y^{(i)} - h_{\theta}(x^{(i)})] x_j^{(i)}$$

LMS Algorithms

Batch Gradient Descent

Repeat until convergence {

$$\theta_j := \theta_j + \alpha \sum_{i=1}^m [y^{(i)} - h_{\theta}(x^{(i)})] x_j^{(i)} \quad (\text{for every } j).$$

}

LMS Algorithms

Batch Gradient Descent

Repeat until convergence {

$$\theta_j := \theta_j + \alpha \sum_{i=1}^m [y^{(i)} - h_{\theta}(x^{(i)})] x_j^{(i)} \quad (\text{for every } j).$$

}

Stochastic Gradient Descent

Loop {

for $i = 1$ to m {

$$\theta_j := \theta_j + \alpha [y^{(i)} - h_{\theta}(x^{(i)})] x_j^{(i)} \quad (\text{for every } j).$$

}

}

LMS Algorithms

Mini-Batch Gradient Descent

Repeat until convergence {

$$\theta_j := \theta_j + \alpha \sum_{i=1}^k [y^{(i)} - h_{\theta}(x^{(i)})] x_j^{(i)} \quad (\text{for every } j).$$

}

Here we use mini-batches containing 10 to 1000 examples. This is $k \in [10, 1000]$.

Matrix of Training Examples

Given a training set of m examples, with each example consisting of n variables, then we can construct a $m \times (n + 1)$ matrix:

$$\mathbf{X} = \begin{bmatrix} x_0^{(1)} & x_1^{(1)} & \cdots & x_n^{(1)} \\ x_0^{(2)} & x_1^{(2)} & \cdots & x_n^{(2)} \\ \vdots & \vdots & \ddots & \vdots \\ x_0^{(m)} & x_1^{(m)} & \cdots & x_n^{(m)} \end{bmatrix} = \begin{bmatrix} [\mathbf{x}^{(1)}]^\top \\ [\mathbf{x}^{(2)}]^\top \\ \vdots \\ [\mathbf{x}^{(m)}]^\top \end{bmatrix}$$

Vector of Training Target Values

Let \mathbf{y} be the m -dimensional vector containing the target values from the training set:

$$\mathbf{y} = \begin{bmatrix} y^{(1)} \\ y^{(2)} \\ \vdots \\ y^{(m)} \end{bmatrix}$$

Cost Function $J(\theta)$

We can write the $J(\theta)$ cost function as follows:

$$J(\theta) = \frac{1}{2} \sum_{i=1}^m [h_{\theta}(x^{(i)}) - y^{(i)}]^2$$

Cost Function $J(\theta)$

We can write the $J(\theta)$ cost function as follows:

$$\begin{aligned} J(\theta) &= \frac{1}{2} \sum_{i=1}^m [h_{\theta}(x^{(i)}) - y^{(i)}]^2 \\ &\quad \frac{1}{2} (\mathbf{X}\theta - \mathbf{y})^{\top} (\mathbf{X}\theta - \mathbf{y}) \end{aligned}$$

Cost Function $J(\theta)$

We can write the $J(\theta)$ cost function as follows:

$$J(\theta) = \frac{1}{2} \sum_{i=1}^m [h_{\theta}(x^{(i)}) - y^{(i)}]^2$$
$$\frac{1}{2}(\mathbf{X}\theta - \mathbf{y})^{\top}(\mathbf{X}\theta - \mathbf{y})$$

and the $\nabla_{\theta}J(\theta)$ can be written as:

$$\nabla_{\theta}J(\theta) = \nabla_{\theta}\frac{1}{2}(\mathbf{X}\theta - \mathbf{y})^{\top}(\mathbf{X}\theta - \mathbf{y})$$

Cost Function $J(\theta)$

We can write the $J(\theta)$ cost function as follows:

$$J(\theta) = \frac{1}{2} \sum_{i=1}^m [h_{\theta}(x^{(i)}) - y^{(i)}]^2$$
$$\frac{1}{2}(\mathbf{X}\theta - \mathbf{y})^{\top}(\mathbf{X}\theta - \mathbf{y})$$

and the $\nabla_{\theta}J(\theta)$ can be written as:

$$\nabla_{\theta}J(\theta) = \nabla_{\theta}\frac{1}{2}(\mathbf{X}\theta - \mathbf{y})^{\top}(\mathbf{X}\theta - \mathbf{y})$$
$$\nabla_{\theta}J(\theta) = \mathbf{X}^{\top}\mathbf{X}\theta - \mathbf{X}^{\top}\mathbf{y}$$

Cost Function $J(\theta)$

We can write the $J(\theta)$ cost function as follows:

$$J(\theta) = \frac{1}{2} \sum_{i=1}^m [h_{\theta}(x^{(i)}) - y^{(i)}]^2$$
$$\frac{1}{2}(\mathbf{X}\theta - \mathbf{y})^{\top}(\mathbf{X}\theta - \mathbf{y})$$

and the $\nabla_{\theta}J(\theta)$ can be written as:

$$\nabla_{\theta}J(\theta) = \nabla_{\theta}\frac{1}{2}(\mathbf{X}\theta - \mathbf{y})^{\top}(\mathbf{X}\theta - \mathbf{y})$$

$$\nabla_{\theta}J(\theta) = \mathbf{X}^{\top}\mathbf{X}\theta - \mathbf{X}^{\top}\mathbf{y}$$

$$0 = \mathbf{X}^{\top}\mathbf{X}\theta - \mathbf{X}^{\top}\mathbf{y}$$

Cost Function $J(\theta)$

We can write the $J(\theta)$ cost function as follows:

$$J(\theta) = \frac{1}{2} \sum_{i=1}^m [h_{\theta}(x^{(i)}) - y^{(i)}]^2$$
$$\frac{1}{2}(\mathbf{X}\theta - \mathbf{y})^{\top}(\mathbf{X}\theta - \mathbf{y})$$

and the $\nabla_{\theta}J(\theta)$ can be written as:

$$\nabla_{\theta}J(\theta) = \nabla_{\theta}\frac{1}{2}(\mathbf{X}\theta - \mathbf{y})^{\top}(\mathbf{X}\theta - \mathbf{y})$$

$$\nabla_{\theta}J(\theta) = \mathbf{X}^{\top}\mathbf{X}\theta - \mathbf{X}^{\top}\mathbf{y}$$

$$0 = \mathbf{X}^{\top}\mathbf{X}\theta - \mathbf{X}^{\top}\mathbf{y}$$

$$\mathbf{X}^{\top}\mathbf{X}\theta = \mathbf{X}^{\top}\mathbf{y}$$

Cost Function $J(\theta)$

We can write the $J(\theta)$ cost function as follows:

$$J(\theta) = \frac{1}{2} \sum_{i=1}^m [h_{\theta}(x^{(i)}) - y^{(i)}]^2$$
$$\frac{1}{2}(\mathbf{X}\theta - \mathbf{y})^{\top}(\mathbf{X}\theta - \mathbf{y})$$

and the $\nabla_{\theta}J(\theta)$ can be written as:

$$\nabla_{\theta}J(\theta) = \nabla_{\theta}\frac{1}{2}(\mathbf{X}\theta - \mathbf{y})^{\top}(\mathbf{X}\theta - \mathbf{y})$$

$$\nabla_{\theta}J(\theta) = \mathbf{X}^{\top}\mathbf{X}\theta - \mathbf{X}^{\top}\mathbf{y}$$

$$0 = \mathbf{X}^{\top}\mathbf{X}\theta - \mathbf{X}^{\top}\mathbf{y}$$

$$\mathbf{X}^{\top}\mathbf{X}\theta = \mathbf{X}^{\top}\mathbf{y}$$

$$\theta = (\mathbf{X}^{\top}\mathbf{X})^{-1}\mathbf{X}^{\top}\mathbf{y}.$$

Computing Directly θ

For an n by n square matrix A , the trace of A is defined to be the sum of its diagonal entries

$$\text{tr } A = \sum_{i=1}^n A_{ii}$$

Computing Directly θ

For an n by n square matrix A , the trace of A is defined to be the sum of its diagonal entries

$$\text{tr } A = \sum_{i=1}^n A_{ii}$$

If a is a real number, then

$$\text{tr } a = a$$

Computing Directly θ

For matrices A , B , C and D , we have that

$$\text{tr } AB = \text{tr } BA$$

Computing Directly θ

For matrices A , B , C and D , we have that

$$\text{tr } AB = \text{tr } BA$$

$$\text{tr } ABC = \text{tr } CAB = \text{tr } BCA$$

Computing Directly θ

For matrices A , B , C and D , we have that

$$\text{tr } AB = \text{tr } BA$$

$$\text{tr } ABC = \text{tr } CAB = \text{tr } BCA$$

$$\text{tr } ABCD = \text{tr } DABC = \text{tr } CDAB = \text{tr } BCDA$$

Computing Directly θ

For matrices A and B , and real number a , we have that

$$\text{tr } A = \text{tr } A^T$$

Computing Directly θ

For matrices A and B , and real number a , we have that

$$\text{tr } A = \text{tr } A^{\top}$$

$$\text{tr } A + B = \text{tr } A + \text{tr } B$$

Computing Directly θ

For matrices A and B , and real number a , we have that

$$\text{tr } A = \text{tr } A^T$$

$$\text{tr } A + B = \text{tr } A + \text{tr } B$$

$$\text{tr } aA = a \text{tr } A$$

Computing Directly θ

For matrices A and B , and real number a , we have that

$$\text{tr } A = \text{tr } A^\top$$

$$\text{tr } A + B = \text{tr } A + \text{tr } B$$

$$\text{tr } aA = a \text{tr } A$$

$$\nabla_A \text{tr } AB = B^\top$$

Computing Directly θ

For matrices A and B , and real number a , we have that

$$\text{tr } A = \text{tr } A^\top$$

$$\text{tr } A + B = \text{tr } A + \text{tr } B$$

$$\text{tr } aA = a \text{tr } A$$

$$\nabla_A \text{tr } AB = B^\top$$

$$\nabla_{A^\top} f(A) = (\nabla_A f(A))^\top$$

Computing Directly θ

For matrices A and B , and real number a , we have that

$$\text{tr } A = \text{tr } A^\top$$

$$\text{tr } A + B = \text{tr } A + \text{tr } B$$

$$\text{tr } aA = a \text{tr } A$$

$$\nabla_A \text{tr } AB = B^\top$$

$$\nabla_{A^\top} f(A) = (\nabla_A f(A))^\top$$

$$\nabla_{A^\top} \text{tr } ABA^\top C = B^\top A^\top C^\top + BA^\top C$$

Computing Directly θ

$$\nabla_{\theta} J(\theta) = \nabla_{\theta} \frac{1}{2} (\mathbf{X}\theta - \mathbf{y})^{\top} (\mathbf{X}\theta - \mathbf{y}).$$

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \nabla_{\theta} \frac{1}{2} (\mathbf{X}\theta - \mathbf{y})^{\top} (\mathbf{X}\theta - \mathbf{y}). \\ &= \nabla_{\theta} \frac{1}{2} (\theta^{\top} \mathbf{X}^{\top} - \mathbf{y}^{\top}) (\mathbf{X}\theta - \mathbf{y}).\end{aligned}$$

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \nabla_{\theta} \frac{1}{2} (\mathbf{X}\theta - \mathbf{y})^{\top} (\mathbf{X}\theta - \mathbf{y}). \\ &= \nabla_{\theta} \frac{1}{2} (\theta^{\top} \mathbf{X}^{\top} - \mathbf{y}^{\top}) (\mathbf{X}\theta - \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} (\theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \mathbf{y}^{\top} \mathbf{X} \theta + \mathbf{y}^{\top} \mathbf{y}).\end{aligned}$$

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \nabla_{\theta} \frac{1}{2} (\mathbf{X}\theta - \mathbf{y})^{\top} (\mathbf{X}\theta - \mathbf{y}). \\ &= \nabla_{\theta} \frac{1}{2} (\theta^{\top} \mathbf{X}^{\top} - \mathbf{y}^{\top}) (\mathbf{X}\theta - \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} (\theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \mathbf{y}^{\top} \mathbf{X} \theta + \mathbf{y}^{\top} \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} (\theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \mathbf{y}^{\top} \mathbf{X} \theta + \mathbf{y}^{\top} \mathbf{y}).\end{aligned}$$

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \nabla_{\theta} \frac{1}{2} (\mathbf{X}\theta - \mathbf{y})^{\top} (\mathbf{X}\theta - \mathbf{y}). \\ &= \nabla_{\theta} \frac{1}{2} (\theta^{\top} \mathbf{X}^{\top} - \mathbf{y}^{\top}) (\mathbf{X}\theta - \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} (\theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \mathbf{y}^{\top} \mathbf{X} \theta + \mathbf{y}^{\top} \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} (\theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \mathbf{y}^{\top} \mathbf{X} \theta + \mathbf{y}^{\top} \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta + \operatorname{tr} \mathbf{y}^{\top} \mathbf{y}.\end{aligned}$$

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \nabla_{\theta} \frac{1}{2} (\mathbf{X}\theta - \mathbf{y})^{\top} (\mathbf{X}\theta - \mathbf{y}). \\ &= \nabla_{\theta} \frac{1}{2} (\theta^{\top} \mathbf{X}^{\top} - \mathbf{y}^{\top}) (\mathbf{X}\theta - \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} (\theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \mathbf{y}^{\top} \mathbf{X} \theta + \mathbf{y}^{\top} \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} (\theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \mathbf{y}^{\top} \mathbf{X} \theta + \mathbf{y}^{\top} \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta + \operatorname{tr} \mathbf{y}^{\top} \mathbf{y}.\end{aligned}$$

Using $\operatorname{tr} A = \operatorname{tr} A^{\top}$ and $(ABC)^{\top} = C^{\top} B^{\top} A^{\top}$,
we have $\operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{y} = \operatorname{tr} (\theta^{\top} \mathbf{X}^{\top} \mathbf{y})^{\top} = \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta$.

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \nabla_{\theta} \frac{1}{2} (\mathbf{X}\theta - \mathbf{y})^{\top} (\mathbf{X}\theta - \mathbf{y}). \\ &= \nabla_{\theta} \frac{1}{2} (\theta^{\top} \mathbf{X}^{\top} - \mathbf{y}^{\top}) (\mathbf{X}\theta - \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} (\theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \mathbf{y}^{\top} \mathbf{X} \theta + \mathbf{y}^{\top} \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} (\theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \mathbf{y}^{\top} \mathbf{X} \theta + \mathbf{y}^{\top} \mathbf{y}). \\ &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{y} - \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta + \operatorname{tr} \mathbf{y}^{\top} \mathbf{y}.\end{aligned}$$

Using $\operatorname{tr} A = \operatorname{tr} A^{\top}$ and $(ABC)^{\top} = C^{\top} B^{\top} A^{\top}$,
we have $\operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{y} = \operatorname{tr} (\theta^{\top} \mathbf{X}^{\top} \mathbf{y})^{\top} = \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta$.

$$= \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta.$$

Computing Directly θ

$$\nabla_{\theta} J(\theta) = \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta.$$

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta. \\ &\quad \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \nabla_{\theta} \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta.\end{aligned}$$

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta. \\ &\quad \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \nabla_{\theta} \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta.\end{aligned}$$

Using $\operatorname{tr} AB = \operatorname{tr} BA$, with $A = \mathbf{y}^{\top} \mathbf{X}$, $B = \theta$.

$$\frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \nabla_{\theta} \operatorname{tr} \theta \mathbf{y}^{\top} \mathbf{X}.$$

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta. \\ &\quad \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \nabla_{\theta} \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta.\end{aligned}$$

Using $\operatorname{tr} AB = \operatorname{tr} BA$, with $A = \mathbf{y}^{\top} \mathbf{X}$, $B = \theta$.

$$\frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \nabla_{\theta} \operatorname{tr} \theta \mathbf{y}^{\top} \mathbf{X}.$$

Using $\nabla_{A^{\top}} \operatorname{tr} ABA^{\top} C = B^{\top} A^{\top} C^{\top} + BA^{\top} C$,

with $A^{\top} = \theta$, $B = \mathbf{X}^{\top} \mathbf{X}$, $C = I$,

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta. \\ &\quad \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \nabla_{\theta} \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta.\end{aligned}$$

Using $\operatorname{tr} AB = \operatorname{tr} BA$, with $A = \mathbf{y}^{\top} \mathbf{X}$, $B = \theta$.

$$\frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \nabla_{\theta} \operatorname{tr} \theta \mathbf{y}^{\top} \mathbf{X}.$$

Using $\nabla_{A^{\top}} \operatorname{tr} ABA^{\top} C = B^{\top} A^{\top} C^{\top} + BA^{\top} C$,

with $A^{\top} = \theta$, $B = \mathbf{X}^{\top} \mathbf{X}$, $C = I$,

and using $\nabla_A \operatorname{tr} AB = B^{\top}$, with $A = \theta$, $B = \mathbf{y}^{\top} \mathbf{X}$.

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta. \\ &\quad \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \nabla_{\theta} \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta.\end{aligned}$$

Using $\operatorname{tr} AB = \operatorname{tr} BA$, with $A = \mathbf{y}^{\top} \mathbf{X}$, $B = \theta$.

$$\frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \nabla_{\theta} \operatorname{tr} \theta \mathbf{y}^{\top} \mathbf{X}.$$

Using $\nabla_{A^{\top}} \operatorname{tr} ABA^{\top} C = B^{\top} A^{\top} C^{\top} + BA^{\top} C$,

with $A^{\top} = \theta$, $B = \mathbf{X}^{\top} \mathbf{X}$, $C = I$,

and using $\nabla_A \operatorname{tr} AB = B^{\top}$, with $A = \theta$, $B = \mathbf{y}^{\top} \mathbf{X}$.

$$= \frac{1}{2} (\mathbf{X}^{\top} \mathbf{X} \theta + \mathbf{X}^{\top} \mathbf{X} \theta - 2 \mathbf{X}^{\top} \mathbf{y}).$$

Computing Directly θ

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta. \\ &\quad \frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \nabla_{\theta} \operatorname{tr} \mathbf{y}^{\top} \mathbf{X} \theta.\end{aligned}$$

Using $\operatorname{tr} AB = \operatorname{tr} BA$, with $A = \mathbf{y}^{\top} \mathbf{X}$, $B = \theta$.

$$\frac{1}{2} \nabla_{\theta} \operatorname{tr} \theta^{\top} \mathbf{X}^{\top} \mathbf{X} \theta - 2 \nabla_{\theta} \operatorname{tr} \theta \mathbf{y}^{\top} \mathbf{X}.$$

Using $\nabla_{A^{\top}} \operatorname{tr} ABA^{\top} C = B^{\top} A^{\top} C^{\top} + BA^{\top} C$,

with $A^{\top} = \theta$, $B = \mathbf{X}^{\top} \mathbf{X}$, $C = I$,

and using $\nabla_A \operatorname{tr} AB = B^{\top}$, with $A = \theta$, $B = \mathbf{y}^{\top} \mathbf{X}$.

$$\begin{aligned}&= \frac{1}{2} (\mathbf{X}^{\top} \mathbf{X} \theta + \mathbf{X}^{\top} \mathbf{X} \theta - 2 \mathbf{X}^{\top} \mathbf{y}). \\ &= \mathbf{X}^{\top} \mathbf{X} \theta - \mathbf{X}^{\top} \mathbf{y}.\end{aligned}$$

Why the Cost Function J is Reasonable?

Given a training example i , we may write

$$y^{(i)} = \theta^\top \mathbf{x}^{(i)} + \epsilon^{(i)},$$

with the assumption

$$\epsilon^{(i)} \sim \mathcal{N}(0, \sigma^2).$$

Why the Cost Function J is Reasonable?

Given a training example i , we may write

$$y^{(i)} = \theta^\top \mathbf{x}^{(i)} + \epsilon^{(i)},$$

with the assumption

$$\epsilon^{(i)} \sim \mathcal{N}(0, \sigma^2).$$

Therefore, the density of $\epsilon^{(i)}$ is given by

$$p(\epsilon^{(i)}) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(\epsilon^{(i)})^2}{2\sigma^2}\right).$$

Why the Cost Function J is Reasonable?

Given a training example i , we may write

$$y^{(i)} = \theta^\top \mathbf{x}^{(i)} + \epsilon^{(i)},$$

with the assumption

$$\epsilon^{(i)} \sim \mathcal{N}(0, \sigma^2).$$

Therefore, the density of $\epsilon^{(i)}$ is given by

$$p(\epsilon^{(i)}) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(\epsilon^{(i)})^2}{2\sigma^2}\right).$$

This implies

$$p(y^{(i)} | \mathbf{x}^{(i)}; \theta) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2}{2\sigma^2}\right).$$

Likelihood of θ

The likelihood of θ is:

$$L(\theta) = L(\theta; \mathbf{X}; \mathbf{y}) = p(\mathbf{y}|\mathbf{X}; \theta).$$

Likelihood of θ

The likelihood of θ is:

$$L(\theta) = L(\theta; \mathbf{X}; \mathbf{y}) = p(\mathbf{y}|\mathbf{X}; \theta).$$

Given the independence assumption on the $\epsilon^{(i)}$'s, we can also write:

$$L(\theta) = \prod_{i=1}^m p(y^{(i)}|\mathbf{x}^{(i)}; \theta)$$

Likelihood of θ

The likelihood of θ is:

$$L(\theta) = L(\theta; \mathbf{X}; \mathbf{y}) = p(\mathbf{y}|\mathbf{X}; \theta).$$

Given the independence assumption on the $\epsilon^{(i)}$'s, we can also write:

$$\begin{aligned} L(\theta) &= \prod_{i=1}^m p(y^{(i)}|\mathbf{x}^{(i)}; \theta) \\ &= \prod_{i=1}^m \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2}{2\sigma^2}\right). \end{aligned}$$

Maximum Likelihood of θ

$$\ell = \log L(\theta)$$

Maximum Likelihood of θ

$$\begin{aligned}\ell &= \log L(\theta) \\ &= \log \prod_{i=1}^m \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2}{2\sigma^2}\right)\end{aligned}$$

Maximum Likelihood of θ

$$\begin{aligned}\ell &= \log L(\theta) \\ &= \log \prod_{i=1}^m \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2}{2\sigma^2}\right) \\ &= \sum_{i=1}^m \log \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2}{2\sigma^2}\right)\end{aligned}$$

Maximum Likelihood of θ

$$\begin{aligned}\ell &= \log L(\theta) \\&= \log \prod_{i=1}^m \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2}{2\sigma^2}\right) \\&= \sum_{i=1}^m \log \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2}{2\sigma^2}\right) \\&= m \log \frac{1}{\sqrt{2\pi}\sigma} - \frac{1}{\sigma^2} \cdot \frac{1}{2} \sum_{i=1}^m (y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2\end{aligned}$$

Maximum Likelihood of θ

$$\begin{aligned}\ell &= \log L(\theta) \\&= \log \prod_{i=1}^m \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2}{2\sigma^2}\right) \\&= \sum_{i=1}^m \log \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2}{2\sigma^2}\right) \\&= m \log \frac{1}{\sqrt{2\pi}\sigma} - \frac{1}{\sigma^2} \cdot \frac{1}{2} \sum_{i=1}^m (y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2\end{aligned}$$

Hence, maximizing $\ell(\theta)$ gives the same answer as minimizing

$$\frac{1}{2} \sum_{i=1}^m (y^{(i)} - \theta^\top \mathbf{x}^{(i)})^2.$$

Locally Adjusting the Model

The algorithm works as follows:

- 1 Fit θ to minimize $\sum_i w^{(i)}(y^{(i)} - \theta^\top x^{(i)})^2$.
- 2 Output $\theta^\top x$.

Locally Adjusting the Model

The algorithm works as follows:

- 1 Fit θ to minimize $\sum_i w^{(i)}(y^{(i)} - \theta^\top x^{(i)})^2$.
- 2 Output $\theta^\top x$.

Where $w^{(i)}$'s are non-negative valued weights.

Locally Adjusting the Model

The algorithm works as follows:


- 1 Fit θ to minimize $\sum_i w^{(i)}(y^{(i)} - \theta^\top x^{(i)})^2$.
- 2 Output $\theta^\top x$.

Where $w^{(i)}$'s are non-negative valued weights.

A good choice for the weights is:

$$w^{(i)} = \exp\left(-\frac{(x^{(i)} - x)^2}{2\tau^2}\right)$$

Locally Adjusting the Model



Thank you!

Dr. Víctor Uc Cetina
cetina@informatik.uni-hamburg.de