

# MATHEMATICS

## SECTION A

January 11, 2024

### 1 Matrices

1.  $A$  is a square matrix with  $|A| = 4$ . Then find the value of  $|A \cdot (\text{adj}A)|$ .
2. For the matrix  $A = \begin{pmatrix} 2 & 3 \\ 5 & 7 \end{pmatrix}$ . Find  $(A + A')$  and verify that it is a symmetric matrix.
3. Using elementary row transformations, find the inverse of the matrix  $\begin{pmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{pmatrix}$ .
4. Using matrices, solve the following system of linear equations :

$$x + 2y - 3z = -4$$

$$2x + 3y + 2z = 2$$

$$3x - 3y - 4z = 11$$

5. Using properties of determinants, find the value of  $k$  if  $\begin{vmatrix} x & y & x+y \\ y & x+y & x \\ x+y & x & y \end{vmatrix} = k|x^3 + y^3|$ .

### 2 probability

6. In a multiple choice examination with three possible answers for each of the five questions, what is the probability that a candidate would get four or more correct answers just by guessing ?
7. The probabilities of solving a specific problem independently by  $A$  and  $B$  are  $\frac{1}{3}$  and  $\frac{1}{5}$  respectively. If both try to solve the problem independently, find the probability that the problem is solved.
8. There are three coins. One is a coin having tails on both faces, another is a biased coin that comes up tails 70% of the time and the third is an unbiased coin. One of the coins is chosen at random and tossed, it shows tail. Find the probability that it was a coin with tail on both the faces.

### 3 Algebra

9. Prove that :

$$\cos^{-1}\left(\frac{12}{13}\right) + \sin^{-1}\left(\frac{3}{5}\right) = \sin^{-1}\left(\frac{56}{65}\right)$$

## 4 vector

10. Find the vector equation of the plane which contains the line of intersection of the planes  $\vec{r} \cdot (\hat{i} + 2\hat{j} + 3\hat{k}) - 4 = 0$ ,  $\vec{r} \cdot (2\hat{i} + \hat{j} - \hat{k}) + 5 = 0$  and which is perpendicular to the plane  $\vec{r} \cdot (5\hat{i} + 3\hat{j} - 6\hat{k}) + 8 = 0$ .
11. Find the vector equation of a line passing through the point  $(2, 3, 2)$  and parallel to the line  $\vec{r} = (-2\hat{i} + 3\hat{j})\lambda + (2\hat{i} - 3\hat{j} + 6\hat{k})$ . Also, find the distance between these two lines.
12. Find the value of  $x$  such that the four points with position vectors,  $\mathbf{A}(3\hat{i} + 2\hat{j} + \hat{k})$ ,  $\mathbf{B}(4\hat{i} + x\hat{j} + 5\hat{k})$ ,  $\mathbf{C}(4\hat{i} + 2\hat{j} - 2\hat{k})$ ,  $\mathbf{D}(6\hat{i} + 5\hat{j} - \hat{k})$  are coplanar.
13. Find the vector equation of the plane determined by the points  $\mathbf{A}(3, -1, 2)$ ,  $\mathbf{B}(5, 2, 4)$  and  $\mathbf{C}(-1, -1, 6)$ . Hence, find the distance of the plane, thus obtained, from the origin.
14. Find the coordinates of the foot of the perpendicular  $\mathbf{Q}$  drawn from  $\mathbf{P}(3, 2, 1)$  to the plane  $2x - y + z + 1 = 0$ . Also, find the distance  $\mathbf{PQ}$  and the image of the point  $\mathbf{P}$  treating this plane as a mirror.

## 5 Differentiation

15. A ladder  $13m$  long is leaning against a vertical wall. The bottom of the ladder is dragged away from the wall along the ground at the rate of  $2cm/sec$ . How fast is the height on the wall decreasing when the foot of the ladder is  $5m$  away from the wall?
16. If  $y = (\log x)^x + x^{\log x}$ , find  $\frac{dy}{dx}$ .
17. If  $x = \sin t$ ,  $y = \sin pt$ , prove that  $(1 - x^2)\frac{d^2y}{dx^2} - x\frac{dy}{dx} + p^2y = 0$ .
18. Differentiate

$$\tan^{-1} \left( \frac{\sqrt{1+x^2} - \sqrt{1-x^2}}{\sqrt{1+x^2} + \sqrt{1-x^2}} \right)$$

with respect to  $\cos^{-1} x^2$ .

19. Form the differential equation representing the family of curves  $y = A \sin x$ , by eliminating the arbitrary constant  $A$ .

## 6 Integration

20. Find the particular solution of the differential equation  $\frac{dy}{dx} = \frac{xy}{x^2 + y^2}$ , given that  $y = 1$  when  $x = 0$ .
21. Solve the differential equation  $\frac{dy}{dx} = 1 + x^2 + y^2 + x^2y^2$ , given that  $y = 1$  when  $x = 0$ .
22. Prove that  $\int_0^a f(x) dx = \int_0^a f(a-x) dx$ , and hence evaluate  $\int_0^1 x^2 (1-x)^n dx$ .

## 7 Linear forms

23. Using integration, find the area of the region bounded by the parabola  $y^2 = 4x$  and the circle  $4x^2 + 4y^2 = 9$ .
24. Using the method of integration, find the area of the region bounded by the lines  $3x - 2y + 1 = 0$ ,  $2x + 3y - 21 = 0$  and  $x - 5y + 9 = 0$ .
25. Using integration, find the area of the region bounded by the line  $y = 3x + 2$ , the  $x$ -axis and the ordinates  $x = -2$  and  $x = 1$ .
26. Using integration, find the area of the smaller region bounded by the ellipse  $\frac{x^2}{9} + \frac{y^2}{4} = 1$  and the line  $\frac{x}{3} + \frac{y}{4} = 1$ .

## 8 Functions

27. Let  $A = \mathbb{R} - \{2\}$  and  $B = \mathbb{R} - \{1\}$ . If  $f : A \rightarrow B$  is a function defined by  $f(x) = \frac{x-1}{x-2}$ , show that  $f$  is one-one and onto. Hence, find  $f^{-1}$ .
28. Show that the relation  $S$  in the set  $A = \{x \in \mathbb{Z} : 0 \leq x \leq 12\}$  given by  $S = \{(a, b) : a, b \in \mathbb{Z}, a - b \text{ is divisible by } 3\}$  is an equivalence relation.

## 9 Optimization

29. A dietician wishes to mix two types of food in such a way that the vitamin contents of the mixture contains at least 8 units of vitamin  $A$  and 10 units of vitamin  $C$ . Food  $I$  contains 2 units/kg of vitamin  $A$  and 1 unit/kg of vitamin  $C$ . It costs ₹50 per kg to produce food  $I$ . Food  $II$  contains 1 unit/kg of vitamin  $A$  and 2 units/kg of vitamin  $C$  and it costs ₹70 per kg to produce food  $II$ . Formulate this problem as a LPP to minimise the cost of a mixture that will produce the required diet. Also find the minimum cost.
30. A manufacturer produces nuts and bolts. It takes 1 hour of work on machine  $A$  and 3 hours on machine  $B$  to produce a package of nuts. It takes 3 hours on machine  $A$  and 1 hour on machine  $B$  to produce a package of bolts. He earns a profit of ₹ 35 per package of nuts and ₹ 14 per package of bolts. How many packages of each should be produced each day so as to maximise his profit, if he operates each machine for atmost 12 hours a day ? Convert it into an LPP and solve graphically.