



Clinical Practice Guideline for Providers: Management of Asthma in Children and Adults

Assessment

In the United States, 18.7 million non-institutionalized adults (8%) and 7 million children (9.4%) have been diagnosed with asthma. Rates of asthma are highest among females (10%); race / ethnicity prevalence is highest among Blacks (12%) followed by Whites (8%) and Hispanics (7%). A critical aspect of the diagnosis and management of asthma is the precise and periodic measurement of lung function – both before and after broncho-dilator therapy to determine both the severity and the effectiveness of therapeutic interventions. When establishing the diagnosis of asthma, evaluate:

- **Medical history** including smoking, drug and alcohol use; physical examination; and supportive diagnostic lung function testing.
- **Establish that episodic symptoms of airflow obstruction are present**, and objectively demonstrate that obstruction is at least partially reversible with Spirometry.
- **Exclude the presence of any alternative diagnoses**, particularly COPD or vocal chord obstruction in adults; and aspiration, cardiac failure, inhaled foreign body, structural abnormality or cystic fibrosis in children.
- **Medication requirements.** Short-acting bronchodilators used more than twice per week should prompt daily inhaled corticosteroid administration for persistent asthma, even if mild severity

Measures of assessment and monitoring should include:

- **Spirometry**, to be conducted at least once a year before and after inhaled bronchodilator therapy. Significant reversibility is indicated by an increase of $\geq 12\%$ and 200 ml in FEV1
- **Peak Flow.** Symptomatic patients with normal spirometry should:
 - Have a daily assessment of peak flow monitoring upon rising and before bedtime; and
 - Maintain an accurate log of daily measurements to help detect subtle changes in lung function that may otherwise go unnoticed by the patient or the provider.

Contributing Factors

Assess at the initial evaluation & additional visits based on seasonal variations:

- Smoking and secondhand smoke. If the member smokes, address the value and available resources to aid in smoking cessation.
- Identify possible environmental inhalant allergens, indoor irritants, pet dander, air pollution
- Viral Respiratory Infection component to induction of Reactive Airways Disease
- Identify all the modifiable risk factors: sedentary lifestyle, obesity, stress, smoking, drug use
- Identify other factors: acute/chronic rhino-sinusitis, gastro-esophageal reflux, drugs (ASA/NSAIDS, sulfites, beta-adrenergic blockers in sensitive patients)

Triggers

- Smoking and secondhand smoke
- Air pollution
- Things the member is allergic to: pet dander, dust mites, cockroaches or pollen

- For exercise induced asthma: advise members on the proper use of inhaler use before they exercise
- Dry, cold air
- Infection
- Some medicines, such as aspirin

Pharmacotherapy

Maintain optimal outcomes:

- Control chronic and nocturnal symptoms
- Maintain normal activity levels, including exercise
- Maintain near normal pulmonary function
- Prevent acute episodes of asthma exacerbation
- Avoid adverse effects of asthma medications
- In addition to allergen avoidance, enhance pharmacotherapy for environmental allergy based seasonal asthma, e.g. daily antihistamines and nasal steroid sprays to avoid asthma induction, daily inhaled corticosteroids during season even if not needed outside of season, etc.
- Annual Influenza immunization; Pneumococcal vaccination as appropriate

Pharmacotherapy Based on Individual's Needs

- **Rescue Medication**
 - Short Acting Beta2 Adrenergic Agonist Bronchodilator
 - Primary medication only for infrequent symptoms or pre-exposure prophylaxis
- **First Line Controller Medication**
 - Inhaled Corticosteroids
 - To be added for ALL persistent disease, no matter how mild
- **Second Line Controller Medication**
 - Long Acting Beta2 Adrenergic Agonist Bronchodilators
 - To be added for asthmatics inadequately controlled on steroids
- **Third Line Medications**
 - Other anti-inflammatory inhalers
 - Only added for asthmatics inadequately controlled on 1st & 2nd step therapy
- **Fourth Line Medications**
 - Methylxanthines
 - Available, but rarely required

Leukotriene modifiers: Include Leukotriene Receptor Antagonist (LTRA) and a 5-lipoxygenase inhibitor. Two LTRAs are available—montelukast (for patients >1 year of age) and zafirlukast (for patients ≥7 years of age). The 5-lipoxygenase pathway inhibitor zileuton is available for patients ≥12 years of age; liver function monitoring is essential. LTRAs are alternative, but not preferred, therapy for the treatment of mild persistent asthma. LTRAs can also be used as adjunctive therapy with ICSs, but for youths ≥12 years of age and adults they are not the preferred adjunctive therapy compared to the addition of LABAs. Zileuton can be used as alternative but not preferred adjunctive therapy in adults.

Patient Education

All patients with Asthma should have a written Asthma Action Plan which incorporates all aspects of their Asthma care. This care plan should be re-evaluated at least annually and more often if necessary to help control the patient's changing condition. A team approach, which includes the patient, the PCP, a certified asthma educator, and a pulmonary specialist when appropriate, is the ideal delivery model for the effective and efficient treatment of Asthma. Toward this end, the

patient must understand his/her Asthma Action Plan – which includes:

- Short and long term goals
- Written environmental control recommendations
- Lifestyle changes including sick day interventions
- Self-monitoring of peak flows with use of a recording system (monthly calendar charting seasonal variations in asthma symptoms)
- Basic facts about asthma (provide written material for patient reference)
- List of environmental controls (stress the importance of implementation)
- Appropriate role of Asthma medications:
 - Explain use of controller vs. reliever medications
 - Provide Asthma Action Plan for medication use
 - Provide use instructions for MDI (observe use and critique technique) and the use of Spacer devices
 - Refer to WellCare Asthma Disease Management Program

Monitoring and Reporting

- Establish therapeutic goals: Normal Activity without restriction, rare symptoms.
- Provide instructions for monitoring and reporting.
 - Practice use of peak flow meter as a monitoring tool and instruct patient to record missed school/work days, altered activity, symptom changes.

Follow Up

- Routine office exams seasonally or every 1 to 6 months if stable, with increased frequency in acute cases or if patient's routine "stable" status changes
- Assess attainment of patient goals and concerns
- Adjust treatment plans as often as necessary for optimal control; add inhaled corticosteroids for all persistent (rescue meds > twice per week) asthma, no matter how mild the severity
- Update the Asthma Action Plan and self-management plan at least annually, and more often as indicated for changes in status
- Re-assess patient's peak flow and inhaler technique
- Smoking cessation program referral for smokers

2012 HEDIS Physician Measurement and Assessment of Compliance with Guidelines

Percent of members aged 5-64 years of age during the measurement year who were identified as having persistent asthma during the year prior to the measurement year and who were appropriately prescribed inhaled corticosteroids, leukotriene modifiers, or Nedocromil during the measurement year.

Components of Severity and Therapy Initiation in Children (0-11 years)

Components of Severity		Classifying Asthma Severity and Initiating Therapy in Children							
		Intermittent		Persistent					
				Mild		Moderate		Severe	
		Ages 0-4	Ages 5-11	Ages 0-4	Ages 5-11	Ages 0-4	Ages 5-11	Ages 0-4	Ages 5-11
Impairment	Symptoms	≤2 days/week		≥2 days/week but not daily		Daily		Throughout the day	
	Nighttime awakenings	0	≤2x/month	1-2x/month	3-4x/month	3-4x/month	>1x/week but not nightly	>1x/week	Often 7x/week
	Short-acting beta ₂ -agonist use for symptom control	≤2 days/week		≥2 days/week but not daily		Daily		Several times per day	
	Interference with normal activity	None		Minor limitation		Some limitation		Extremely limited	
	Lung Function			Normal FEV ₁ between exacerbations					
	• FEV ₁ (predicted) or peak flow (personal best)	N/A	>80%	N/A	>80%	N/A	60-80%	N/A	<60%
	• FEV ₁ /FVC		>85%		>80%		75-80%		<75%
Risk	Exacerbations requiring oral systemic corticosteroids (consider severity and interval since last exacerbation)	0-1/year (see notes)		≥2 exacerbations in 6 months requiring oral systemic corticosteroids, or ≥4 wheezing episodes/1 year lasting >1 day AND risk factors for persistent asthma	≥2x/year (see notes) Relative annual risk may be related to FEV ₁				
Recommended Step for Initiating Therapy (See "Stepwise Approach for Managing Asthma" for treatment steps.)		Step 1 (for both age groups)		Step 2 (for both age groups)		Step 3 and consider short course of oral systemic corticosteroids	Step 3: medium-dose ICS option and consider short course of oral systemic corticosteroids	Step 3 and consider short course of oral systemic corticosteroids	Step 3: medium-dose ICS option OR step 4 and consider short course of oral systemic corticosteroids
The stepwise approach is meant to assist, not replace, the clinical decisionmaking required to meet individual patient needs.		In 2-6 weeks, depending on severity, evaluate level of asthma control that is achieved. • Children 0-4 years old: If no clear benefit is observed in 4-6 weeks, stop treatment and consider alternative diagnoses or adjusting therapy. • Children 5-11 years old: Adjust therapy accordingly.							

Assessing Asthma Control and Adjusting Therapy in Children (0-11 years)

Components of Control		Assessing Asthma Control and Adjusting Therapy in Children			
		Well Controlled	Not Well Controlled		Very Poorly Controlled
Impairment	Symptoms	Ages 0-4	Ages 5-11	Ages 0-4	Ages 5-11
		≤2 days/week but not more than once on each day	≤2 days/week or multiple times on ≤2 days/week	>2 days/week or multiple times on ≤2 days/week	Throughout the day
		≤1x/month	≤2x/month	>1x/month	>1x/week
		None	Some limitation	Extremely limited	Extremely limited
		≤2 days/week	>2 days/week	Several times per day	Several times per day
Risk	Lung function	N/A	>80%	N/A	<60%
			>80%		<75%
					≥2x/year
Risk	Exacerbations requiring oral systemic corticosteroids	0-1x/year	2-3x/year	≥2x/year	≥2x/year
Recommended Action for Treatment	Reduction in lung growth	N/A	Requires long-term followup	N/A	N/A
Recommended Action for Treatment	Treatment-related adverse effects	Medication side effects can vary in intensity from none to very troublesome and worrisome. The level of intensity does not correlate to specific levels of control but should be considered in the overall assessment of risk.			
Recommended Action for Treatment	The stepwise approach is meant to assist, not replace, clinical decisionmaking required to meet individual patient needs.				

Assessing Asthma Control and Adjusting Therapy in Youths ≥ 12 years of Age and Adults

Components of Control		Classification of Asthma Control (≥12 years of age)		
		Well Controlled	Not Well Controlled	Very Poorly Controlled
Impairment	Symptoms	≤2 days/week	>2 days/week	Throughout the day
	Nighttime awakenings	≤2x/month	1–3x/week	≥4x/week
	Interference with normal activity	None	Some limitation	Extremely limited
	Short-acting beta ₂ -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week	Several times per day
	FEV ₁ or peak flow	>80% predicted/ personal best	60–80% predicted/ personal best	<60% predicted/ personal best
	Validated questionnaires ATAQ ACQ ACT	0 ≤0.75* ≥20	1–2 ≥1.5 16–19	3–4 N/A ≤15
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year	≥2/year (see note)	
		Consider severity and interval since last exacerbation		
	Progressive loss of lung function	Evaluation requires long-term followup care.		
	Treatment-related adverse effects	Medication side effects can vary in intensity from none to very troublesome and worrisome. The level of intensity does not correlate to specific levels of control but should be considered in the overall assessment of risk.		
Recommended Action for Treatment (See “Stepwise Approach for Managing Asthma” for treatment steps.)		<ul style="list-style-type: none">• Maintain current step.• Regular followup at every 1–6 months to maintain control.• Consider step down if well controlled for at least 3 months.	<ul style="list-style-type: none">• Step up 1 step.• Reevaluate in 2–6 weeks.• For side effects, consider alternative treatment options.	<ul style="list-style-type: none">• Consider short course of oral systemic corticosteroids.• Step up 1–2 steps.• Reevaluate in 2 weeks.• For side effects, consider alternative treatment options.

Classifying Asthma Severity and Initiating Treatment in Youths ≥ 12 Years of Age and Adults

Components of Severity		Classification of Asthma Severity ≥12 years of age			
		Intermittent	Mild	Moderate	Severe
Symptoms	Symptoms	≤2 days/week	>2 days/week but not daily	Daily	Throughout the day
	Nighttime awakenings	≤2x/month	3–4x/month	>1x/week but not nightly	Often 7x/week
	Short-acting beta ₂ -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week but not daily, and not more than 1x on any day	Daily	Several times per day
	Interference with normal activity	None	Minor limitation	Some limitation	Extremely limited
	Lung function	<ul style="list-style-type: none"> • Normal FEV₁ between exacerbations • FEV₁ >80% predicted • FEV₁/FVC normal 	<ul style="list-style-type: none"> • FEV₁ >80% predicted • FEV₁/FVC normal 	<ul style="list-style-type: none"> • FEV₁ >60% but <80% predicted • FEV₁/FVC reduced 5% 	<ul style="list-style-type: none"> • FEV₁ <60% predicted • FEV₁/FVC reduced >5%
Risk	Impairment	Normal FEV ₁ /FVC: 8–19 yr 85% 20–39 yr 80% 40–59 yr 75% 60–80 yr 70%			
	Exacerbations requiring oral systemic corticosteroids	0–1/year (see note)	≥2/year (see note)	Consider severity and interval since last exacerbation. Frequency and severity may fluctuate over time for patients in any severity category. Relative annual risk of exacerbations may be related to FEV ₁ .	
Recommended Step for Initiating Treatment (See "Stepwise Approach for Managing Asthma" for treatment steps.)		Step 1	Step 2	Step 3 and consider short course of oral systemic corticosteroids	Step 4 or 5
		In 2–6 weeks, evaluate level of asthma control that is achieved and adjust therapy accordingly.			

Figure 1. Summary of Recommended Key Clinical Activities for the Diagnosis and Management of Asthma

Clinical Issue	Key Clinical Activities	Action Steps
DIAGNOSIS		
	Establish asthma diagnosis.	<ul style="list-style-type: none">Use medical history and physical examination to determine that symptoms of recurrent episodes of airflow obstruction are present.Use spirometry in all patients ≥5 years of age to determine that airway obstruction is at least partially reversible.Consider alternative causes of airway obstruction.
MANAGING ASTHMA LONG TERM	Goal of asthma therapy is asthma control: <ul style="list-style-type: none">Reduce impairment (prevent chronic symptoms, require infrequent use of short-acting beta2-agonist (SABA), maintain (near) normal lung function and normal activity levels).Reduce risk (prevent exacerbations, minimize need for emergency care or hospitalization, prevent loss of lung function, or for children, prevent reduced lung growth, have minimal or no adverse effects of therapy).	
FOUR COMPONENTS OF CARE		
Assessment and Monitoring	<p>Assess asthma severity to initiate therapy.</p> <p>Assess asthma control to monitor and adjust therapy.</p> <p>Schedule follow-up care.</p>	<ul style="list-style-type: none">Use severity classification chart, assessing both domains of impairment and risk, to determine initial treatment.Use asthma control chart, assessing both domains of impairment and risk, to determine if therapy should be maintained or adjusted (step up if necessary, step down if possible).Use multiple measures of impairment and risk: different measures assess different manifestations of asthma; they may not correlate with each other; and they may respond differently to therapy. Obtain lung function measures by spirometry at least every 1–2 years, more frequently for not-well-controlled asthma.Asthma is highly variable over time; periodic monitoring is essential. Consider scheduling patients at 2- to 6-week intervals while gaining control; at 1–6 month intervals, depending on step of care required or duration of control, to monitor if sufficient control is maintained; at 3-month intervals if a step down in therapy is anticipated.Assess asthma control, medication technique, written asthma action plan, patient adherence and concerns at every visit.
Education	<p>Provide self-management education.</p> <p><i>Tailor education to literacy level of patient. Appreciate the potential role of a patient's cultural beliefs and practices in asthma management.</i></p> <p>Develop a written asthma action plan in partnership w/ patient.</p> <p>Integrate education into all points of care where health professionals interact with patients.</p>	<p>Teach and reinforce:</p> <ul style="list-style-type: none">Self-monitoring to assess level of asthma control and signs of worsening asthma (either symptom or peak flow monitoring shows similar benefits for most patients). Peak flow monitoring may be helpful for patients who have difficulty perceiving symptoms, a history of severe exacerbations, or moderate or severe asthma.Using written asthma action plan (review differences between long-term control and quick-relief medication).Taking medication correctly (inhaler technique and use of devices).Avoiding environmental factors that worsen asthma.Agree on treatment goals and address patient concerns.Provide instructions for (1) daily management (long-term control medication, if appropriate, and environmental control measures) and (2) managing worsening asthma (how to adjust medication, and know when to seek medical care).Involve all members of the health care team in providing / reinforcing education, including physicians, nurses, pharmacists, respiratory therapists, and asthma educators.Encourage education at all points of care: clinics (offering separate self-

		<p>management education programs as well as incorporating education into every patient visit), Emergency Departments and hospitals, pharmacies, schools and other community settings, and patients' homes.</p> <ul style="list-style-type: none"> • Use a variety of educational strategies and methods.
Control Environmental Factors and Comorbid conditions	<p>Recommend measures to control exposures to allergens and pollutants or irritants that make asthma worse.</p> <p>Treat comorbid conditions.</p>	<ul style="list-style-type: none"> • Determine exposures, history of symptoms in presence of exposures, and sensitivities (In patients who have persistent asthma, use skin or in vitro testing to assess sensitivity to perennial indoor allergens.). • Advise patients on ways to reduce exposure to those allergens and pollutants, or irritants to which the patient is sensitive. Multifaceted approaches are beneficial; single steps alone are generally ineffective. Advise all patients and pregnant women to avoid exposure to tobacco smoke. • Consider allergen immunotherapy, by specifically trained personnel, for patients who have persistent asthma and when there is clear evidence of a relationship between symptoms and exposure to an allergen to which the patient is sensitive. • Consider especially: allergic bronchopulmonary aspergillosis; gastroesophageal reflux, obesity, obstructive sleep apnea, rhinitis and sinusitis, and stress or depression. Recognition and treatment of conditions may improve asthma control. • Consider inactivated influenza vaccine for all patients over 6 months of age.
Medications	Select medication and delivery devices to meet patient's needs and circumstances.	<ul style="list-style-type: none"> • Use stepwise approach (See below.) to identify appropriate treatment options. • Inhaled corticosteroids (ICSs) are the most effective long-term control therapy. When choosing among treatment options, consider domain of relevance to the patient (impairment, risk, or both), patient's history of response to the medication, and patient's willingness and ability to use the medication.

Clinical Issue	Key Clinical Activities	Action Steps
STEPWISE APPROACH		
General Principles for All Age Groups	Incorporate four components of care.	<ul style="list-style-type: none"> • Include medications, patient education, environmental control measures, and management of comorbidities at each step. Monitor asthma control regularly (See above, assessment and monitoring.)
	Initiate therapy based on asthma severity.	<ul style="list-style-type: none"> • For patients not taking long-term control therapy, select treatment step based on severity (See figures on stepwise approach for different age groups.). Patients who have persistent asthma require daily long-term control medication.
	Adjust therapy based on asthma control.	<ul style="list-style-type: none"> • Once therapy is initiated, monitor the level of asthma control and adjust therapy accordingly: step up if necessary and step down if possible to identify the minimum amount of medication required to maintain asthma control. • Refer to an asthma specialist for consultation or co-management if there are difficulties achieving or maintaining control; step 4 care or higher is required (step 3 care or higher for children 0–4 years of age); immunotherapy or omalizumab is considered; or additional testing is indicated; or if the patient required 2 bursts of oral systemic corticosteroids in the past year or a hospitalization.

For age specific principles using the Stepwise approach, refer to the National Heart, Lung, and Blood Institute's Expert Panel Report 3 (2007) at <http://www.nhlbi.nih.gov/guidelines/asthma/index.htm> Guidance is also available on treating patients with respect to pregnancy, surgery, and management in home or emergency care settings.

References

National Heart, Lung, and Blood Institute. (2007). Expert panel report 3: guidelines for the diagnosis and management of asthma. Retrieved from <http://www.nhlbi.nih.gov/guidelines/asthma/index.htm>

National Committee for Quality Assurance. (2012). HEDIS 2012 technical specifications for physician measurement: summary table of measure changes. Retrieved from <http://www.ncqa.org/LinkClick.aspx?fileticket=K4wzHTiYHug%3d&tabid=1415>



CLINICAL PRACTICE GUIDELINE FOR PROVIDERS: MANAGEMENT OF ASTHMA IN CHILDREN AND ADULTS HS-1001

Legal Disclaimer: Clinical Practice Guidelines made available by WellCare are informational in nature and are not a substitute for the professional medical judgment of treating physicians or other health care practitioners. These guidelines are based on information available at the time and may not be updated with the most current information available at subsequent times. Individuals should consult with their physician(s) regarding the appropriateness of care or treatment options to meet their specific needs or medical condition. Disclosure of clinical practice guidelines is not a guarantee of coverage. Members of WellCare health plans should consult their individual coverage documents for information regarding covered benefits. WellCare does not offer medical advice or provide medical care, and therefore cannot guarantee any results or outcomes. WellCare does not warrant or guarantee, and shall not be liable for any deficiencies in the information contained herein or for any inaccuracies or recommendations made by independent third parties from whom any of the information contained herein was obtained. *Note: The lines of business (LOB) are subject to change without notice; consult www.wellcare.com/Providers/CPGs for list of current LOBs.*

The WellCare Group of Companies

Easy Choice Health Plan ~ Harmony Health Plan of Illinois, Inc. ~ Missouri Care, Inc.

'Ohana Health Plan, a plan offered by WellCare Health Insurance of Arizona, Inc. ~ WellCare Health Insurance of Illinois, Inc.

WellCare Health Plans of New Jersey, Inc. ~ WellCare Health Insurance of Arizona, Inc. ~ WellCare of Florida, Inc. ~ WellCare of Connecticut, Inc.

WellCare of Georgia, Inc. ~ WellCare of Kentucky, Inc. ~ WellCare of Louisiana, Inc. ~ WellCare of New York, Inc. ~ WellCare of Ohio, Inc.

WellCare of South Carolina, Inc. ~ WellCare of Texas, Inc. ~ WellCare Prescription Insurance, Inc.

Date

6/7/2012

12/1/2011

9/2010

History and Revisions by the Medical Policy Committee

- Approved by MPC. Added *Recommended Key Clinical Activities for the Diagnosis and Management of Asthma* (source: NHLBI, 2007). See pp. 8-9. Updated HEDIS age range, inserted new reference.
- New template design approved by MPC.
- Approved by MPC.