

Search for safety, tips, illness, etc.



Ages & Stages Healthy Living

Safety & Prevention Family Life Health Issues News

Español

SAFETY & PREVENTION LISTEN)

Tips & Tools Our Mission



Safety & Prevention

Immunizations

All Around

At Home

Medication Safety

At Play

On The Go



Ibuprofen Dosage Table for Fever and Pain

Healthy Children > Safety & Prevention > At Home > Medication Safety > Ibuprofen Dosage Table for Fever and Pain

Ibuprofen (i.e., Motrin®, Advil®) is an over-the-counter medicine used to reduce fever and relieve pain. This chart, based on your child's weight, can help determine the right dosage amount, but is no substitute for your pediatrician's advice.

Ibuprofen Dosage Table for Fever and Pain*

Child's Weight (pounds)	12-17	18-23	24-35	36-47	48-59	60-71	72-95	96+	
Infant Drops 50 mg/1.25 mL	1.25	1.875	2.5	3.75	5	-		. "	mL
Liquid 100mg/5 mL	2.5	4	5	7.5	10	12.5	15	20	mL
Chewable 50 mg tablets		-	2	3	4	5	6	8	tabs
Junior strength 100 mg tablets	-	-	-		2	2 1/2	3	4	tabs
Adult 200 mg tablets	-	-		: "4	1	1	1 1/2	2	tabs

^{*}Read table notes below.

Table Notes:

- . Age limit: Don't use under 6 months of age unless directed by your child's pediatrician, because safety has not established and doesn't have FDA approval. Avoid multi-ingredient products in children under 6 years of age.
- Dosage: Determine by finding child's weight in the top row of the dosage table.
- . Measuring the dosage (should be in metric units): Dosing syringes are more accurate than household utensils. Use the syringe or device that comes with the medication. If not available with the medication, medicine syringes are available at pharmacies. Household spoons are not reliable. Note: 1 level teaspoon equals 5 mL and that 1/2 teaspoon equals 2.5 mL.
- . Ibuprofen drops: Ibuprofen infant drops come with a measuring syringe
- · Adult dosage: 400 mg
- Frequency: Repeat every 6-8 hours as needed.

Additional Information & Resources:

- · Fever and Your Baby
- · How to Take a Child's Temperature
- The Healthy Children Show: Fever (Video)
- The Healthy Children Show: Giving Liquid Medicine Safely (Video)
- · Ibuprofen: Info for Consumers (National Library of Medicine)

Last Updated 5/23/2016

Source American Academy of Pediatrics (Copyright © 2016)

The information contained on this Web site should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.









Contact Us About Us Privacy Policy Terms of Use Editorial Policy





© Copyright 2017 American Academy of Pediatrics. All rights reserved.

Edades y Etapas Vida sana Seguridad y Prevención Vida familiar Problemas de salud Noticias Consejos y Herramientas Nuestra misión 👜 ShopAAP



Ingrese su búsqueda



Seguridad y

Prevención Inmunizaciones

En el hogar

Uso seguro de medicamentos

Durante el juego

En marcha

En todas partes

Tabla de dosis del ibuprofeno para la fiebre y el dolor

El ibuprofeno (por ejemplo: Motrin®, Advil®) es un medicamento de venta libre que se usan para tratar la fiebre y aliviar el dolor. La siguiente tabla, basada en el peso corporal del niño, puede ayudarle a determinar la dosis correcta, pero no sustituye las recomendaciones de su médico.

Tabla de dosis del ibuprofeno para aliviar la fiebre y el dolor *

Peso del niño en libras	12-17	18-23	24-35	36-47	48-59	60-71	72-95	96+	
Gotas para bebé 50 mg/1,25 mL	1,25	1,875	2,5	3,75	5				mL
Líquido 100 mg/5 mL	2,5	4	5	7,5	10	12,5	15	20	mL
Masticable 50 mg			2	3	4	5	6	8	Tableta
Para jóvenes 100 mg		-	-	-	2	2 1/3	3	4	Tableta
Para adulto 200 mg					1	1	1 1/2	2	Tableta

^{*}Lea las notas de la tabla a continuación

Notas sobre la tabla:

- Límite de edad: No use en bebés menores de 6 meses de edad a menos que lo
 indique el médico del bebé. (No se ha establecido su seguridad y no cuenta con
 aprobación del FDA). Evite los productos de varios ingredientes en niños menores
 de 6 años de edad.
- Dosis: Se determina al buscar el peso del niño en la fila superior de la tabla de dosis.
- Cómo medir la dosis: (debe medirse en unidades métricas). Las jeringas de dosis son más exactas que los utensilios caseros. Use una jeringa o el dispositivo que viene con el medicamento. Si no viene con el medicamento, puede conseguir una jeringa en cualquier farmacia. Los utensilios caseros no son exactos. Nota: 1 cucharadita nivelada equivale a 5 mL y 1/2 cucharadita equivale a 2,5 mL.
- Ibuprofeno en gotas: Las gotas de ibuprofeno infantil vienen con una jeringa medidora.
- Dosis para adultos: 400 mg

• Frecuencia: Repita cada 6 a 8 horas, según sea necesario

Información adicional:

- La fiebre y su bebé
- · Cómo tomarle la temperatura a un niño

Última actualización 1/12/2017

Fuente American Academy of Pediatrics (Copyright © 2016)

La información contenida en este sitio web no debe usarse como sustituto al consejo y cuidado médico de su pediatra. Puede haber muchas variaciones en el tratamiento que su pediatra podría recomendar basado en hechos y circunstancias individuales.









Contáctenos ¿Quiénes somos? Política de privacidad Términos de uso Política editorial





This site complies with the HONcode standard for trustworthy health information:

© derechos de autor 2017 American Academy of Pediatrics. Todos los derechos reservados.