

Neck Pain, Cervicalgia

Diagnosis/Condition:	Neck Pain, Cervicalgia
Discipline:	DC, ND
ICD-9 Codes:	723.1; 723.3; 739.1; 847.0
ICD-10 Codes:	M54.2; M53.1; M99.01; S13.4XXA; S13.8XXA
Origination Date:	1996
Review/Revised Date:	04/2015
Next Review Date:	04/2017

Neck pain is second only to low back pain as the most common musculoskeletal disorder in population surveys and in primary care. Depending on the case definition of “neck pain,” from 30 to 50% of the adult population experiences neck pain each year.¹ Neck pain is common in children as well. Neck pain has been found in up to 40% of a small sample of pre-adolescents and teenagers.^{2,3} Neck pain is frequently reported to and treated by CAM providers.

Historically segmental dysfunction and cervical sprain/strain are the first and second most common diagnoses reported by CHP chiropractic physicians with neck pain being the second most common diagnosis reported by CHP naturopathic physicians (NDs). There is some evidence that manual therapy techniques have been found to be less costly and more effective than either physiotherapy or general practitioner care in improving neck pain.⁴

While most individuals with acute neck pain do not seek health care, those that do account for a disproportionate amount of health care costs. Most neck pain is not attributable to a specific disease or disorder and is labeled as muscular, mechanical, or postural neck pain. Despite decades of research and posturing to explain chronic neck pain on the basis of a specific disease or injury, and despite increasingly sophisticated diagnostic imaging assessment, little advance has been made in achieving a specific structural diagnosis for this prevalent condition.

Subjective Findings and History

- Risk factors for neck pain include age, gender, and heredity¹
- Modifiable risk factors include tobacco exposure (smoking and second hand), low social support at work, high quantitative work demands, sedentary, precision and repetitive work¹
- Macro trauma: Onset of pain and paraspinal muscle spasm begins either immediately after the injury or gradually over the next 24 hours
- Micro trauma: Repetitive traumatic events not singularly capable of producing injury
- Local pain, sometimes accompanied by referred pain, diffuse (scleratogenous pain distribution)
- Loss of flexibility

- Pain is usually relieved by rest and aggravated by motion

Objective Findings

The validity of most commonly used objective tests is lacking. There is support for subjective self-report assessment in monitoring the patient's course and response to treatment

- Postural evaluation reveals asymmetry, misalignment or decrease of normal spinal curvature
- Decrease/loss of normal spinal ROM
- Palpation: Segmental joint dysfunction/subluxation, tenderness with pressure over involved tissues, muscle spasm or tautness of paravertebral muscles, Myofascial Trigger Points, orthopedic and neurological examination directed at differentiating neurogenic from other sources of pain: absence of nerve compression signs (e.g. absence of muscle weakness); orthopedic tests MAY reproduce the pain (e.g. foraminal compression and other tests that cause spinal motion may increase neck pain)
- Radiographic examination: depending on age and history of prior episodes (see radiographic guidelines at www.chpgroup.com, For Providers tab, Provider Log-in tab, Clinical Tools tab) Note: There is no evidence that common degenerative changes in the cervical spine are a risk factor for neck pain.

Assessment

- Rule out “red flags” of serious pathology in patients seeking care for neck pain without frank trauma. Serious disorders to rule out include: pathologic fractures, neoplasm, systemic inflammatory diseases, infections, cervical myelopathy and/or previous cervical spine or neck surgery, or open injury.⁵
- A complete medical history and description of local and systemic symptoms.
- A thorough physical examination, including musculoskeletal and neurological exams.
- Imaging: refer to CHP Radiographic guidelines. (add link)
- Determining which patients may benefit from treatment for neck pain can aid clinical decision-making. One large study of about 20,000 patients illuminated which ones were likely to experience immediate relief from manipulation. Patients who presented with symptoms of reduced neck, shoulder, arm movement, stiffness, neck pain, upper, mid back pain, headache, shoulder, arm pain, are likely to report immediate improvement in these symptoms after treatment. Patients presenting with any 4 of these symptoms were shown to have the highest probability of immediate improvement.⁶

The Neck Pain Task Force recommends a 4-grade classification system of neck pain severity.⁷

- Grade I neck pain: No signs or symptoms suggestive of major structural pathology and no or minor interference with activities of daily living; will likely respond to minimal intervention such as reassurance and pain control; does not require intensive investigations or ongoing treatment.
- Grade II neck pain: No signs or symptoms of major structural pathology, but major interference with activities of daily living; requires pain relief and early activation/intervention aimed at preventing long-term disability.

- Grade III neck pain: No signs or symptoms of major structural pathology, but presence of neurologic signs such as decreased deep tendon reflexes, weakness, and/or sensory deficits; might require investigation and, occasionally more invasive treatments.
- Grade IV neck pain: Signs or symptoms of major structural pathology, such as fracture, myelopathy, neoplasm, or systemic disease; requires prompt investigation and treatment.

Plan

A number of non-surgical treatments appear to be more beneficial than “usual care” (e.g. medication, advice, exercises), sham, or alternative interventions but none of the active treatments were clearly superior to any other in the short or long term. Educational videos, mobilization, manual therapy, exercises, low-level laser therapy (LLLT), and perhaps acupuncture appeared to have some benefit. Interventions that focus on regaining function and returning to work as soon as possible are relatively more effective than interventions that do not have such a focus.⁷ More recent evidence provides strong recommendations for treatment of chronic neck pain with manipulation, manual therapy and exercise in combination with other modalities.⁸ Moderate to strong recommendations for treatment of chronic neck pain included stretching, strengthening and endurance exercises alone.^{8,9} Moderate recommendations were made for treatment of chronic neck pain with mobilization as well as massage in combination with other therapies and weak recommendations were made for the same condition with manipulation alone. With respect to acute neck pain, moderate recommendations were made for treatment with manipulation and mobilization in combination with other therapies and weak recommendations were made for the same condition with exercise alone.⁸ Treatment of seniors (65 years or older) with neck pain using spinal manipulative therapy proved beneficial when combined with supervised and unsupervised home exercise with the supervised component adding little benefit.¹⁰ There is also evidence of significant and immediate improvement respectively with cervical manipulation in the treatment of symptomatic disc herniations and degenerative cervical radiculopathy.^{11, 12}

Passive Care:

- Cervical and thoracic manipulation, cervical mobilization.^{13,14}
- Physical therapy modalities.
- Prescription/OTC medications: analgesics, NSAIDS, muscle relaxants.
- Topical salicylic acid, other topical analgesic cream.
- Braces/supports.

Active Care:

- Rest from inciting activities.
- Active exercises/stretchers for mobility and strength.
- Ice/heat application at home (hydrotherapy).
- Posture training, ergonomic evaluation, educational interventions.
- Activity/work restrictions, if appropriate.

- Supervised qigong, lyengar yoga, and combined with programs including strengthening, range of motion, and flexibility.¹⁵

Other treatments:

- Nutritional supplementation (Vitamin C, manganese, magnesium, calcium, glucosamine sulfate, chondroitin, methylsulfonylmethane, bromelain and essential fatty acids).
- Botanical supplementation to reduce inflammation (*Curcuma longa* (turmeric), *Capsicum annuum* (cayenne), *Arnica montana* (arnica), *Ruta graveolens* (rue), *Hypericum perforatum* (St. John's wort), and *Gaultheria procumbens* (wintergreen)).
- Topical treatments (e.g. comfrey poultice, hypericum, arnica, wintergreen).
- Homeopathy.
- Acupuncture.
- Massage.¹⁶

Length of Treatment

- Estimated duration of care: 1-6 weeks. Evidence for the dose response to manipulation and physical medicine modalities is sparse. In a small sample (80 patients) the difference between 8 and 16 treatments over an 8 week period was small.¹⁷
- Evaluate progress on an on-going basis.
- Risk factors for chronicity: Significant trauma, co-morbidity (degenerative disc disease, segmental instability, osteoporosis, spine deformity), age, socio-economic factors.

Outcome Assessment Tools

- Visual analog pain scale (VAS), numeric pain rating scale (NRS), quadruple VAS and NRS.
- Neck pain disability index (NDI).

Vertebrobasilar Stroke Risk

There is concern for the association of cervical manipulation, especially upper cervical manipulation, and injury to the vertebrobasilar artery (VBA) resulting in stroke. A population-based case control and case crossover study of 815 VBA stroke cases showed that there was an association between chiropractic services and subsequent vertebrobasilar artery stroke in persons under 45 years of age. But a similar association was also observed among patients receiving general practitioner services. This is likely explained by patients with incipient vertebrobasilar artery dissection-related neck pain or headache seeking care from DCs and MDs before having their stroke and not because of a manipulation.¹⁸ A review of the best evidence indicates that chiropractic care is useful for patients with neck pain for which the risks of serious adverse events should be considered negligible.^{19,20}

Referral Criteria

Referral to an appropriate specialist may be appropriate after 4-6 weeks of care without symptomatic or functional improvement or upon onset of (progressive) neurologic deficit.

Resources for Clinicians

Chiropractic clinical practice guideline: evidence-based treatment of adult neck pain not due to whiplash. Canadian Chiropractic Association Canadian Federation of Chiropractic Regulatory Boards. 2005. 52 pages.

Resources for Patients

Spine-health.com publishes original, award-winning articles written for patients by over 80 physician authors and peer-reviewed by a 23 member Medical Advisory Board. This trusted, independent site is supported by hundreds of physician members and visited by millions of patients and their physicians. <http://www.spine-health.com/pain/neck-pain-0?page=1>

MedlinePlus will direct you to information to help answer health questions. MedlinePlus brings together authoritative information from NLM, the National Institutes of Health (NIH), and other government agencies and health-related organizations.

<http://www.nlm.nih.gov/medlineplus/ency/article/003025.htm>

The Evidence

Raine R, et al. Systematic review of mental health interventions for patients with common somatic symptoms: can research evidence from secondary care be extrapolated to primary care? *BMJ*. 2002; 325(7372):1082.

Gross AR, et al. Manipulation and mobilization for mechanical neck disorders (Cochrane Review) From The Cochrane Library, Issue 1, 2005. Chichester, UK: John Wiley & Sons, Ltd. All rights reserved. Manipulation and mobilization for mechanical neck disorders. <http://www.cochrane.org/cochrane/revabstr/AB004249.htm> (text/html) Mon, 24 Jan 2005

Gordon McMorland, DC, Esther Suter, PhD. Chiropractic management of mechanical neck and low-back pain: A retrospective, outcome-based analysis. *Journal of Manipulative and Physiological Therapeutics* 2000; 23 (5): 307 -311.

Gross AR, et al. Cervical Overview Group. A Cochrane review of manipulation and mobilization for mechanical neck disorders. *Spine*. 2004; 29(14):1541-8.

Walsh MJ. Evaluation of clinical findings for uncomplicated mechanical neck pain. *Chiropractic Journal of Australia* 2000;30(4):122-6.

Wyatt LH. Facet syndrome in the cervical spine. *JACA Journal of the American Chiropractic Association*. 2004; 41(3): 32-7.

Murphy DR. A clinical model for the diagnosis and management of patients with cervical spine syndromes. *Australasian Chiropractic & Osteopathy*. 2004; 12(2): 57-71, 82.

Haas M. et al. Dose Response for Chiropractic Care of Chronic Cervicogenic Headache and Associated Neck Pain: A Randomized Pilot Study. *Journal Of Manipulative And Physiological Therapeutics*. 2004; 27: 547-53.

Chiu TT, Lam TH, Hedley AJ. A Randomized Controlled Trial on the Efficacy of Exercise for Patients With Chronic Neck Pain. *Spine*. 2005;30:E1-7.

Giles LG, Muller R. Chronic Spinal Pain: A Randomized Clinical Trial Comparing Medication, Acupuncture, and Spinal Manipulation. *Spine*. 2003; 28:1490-1501.

Holm L. W. Carroll L. J. David Cassidy J. Ahlbom A. Factors influencing neck pain intensity in whiplash-associated disorders in Sweden. *The Clinical Journal Of Pain*. 2007 ;23:591-7.

Landers MR. Workers' compensation and litigation status influence the functional outcome of patients with neck pain. *The Clinical Journal Of Pain*. 2007;23:676-82.

Hush JM, Maher C.G, Refshauge KM. Risk factors for neck pain in office workers: a prospective study. *BMC Musculoskeletal Disorders*. 2006; 7:81.

Cote P, Cassidy JD, Carroll LJ, Kristman V. The annual incidence and course of neck pain in the general population: a population-based cohort study. *Pain*. 2004; 112:267-73.

Chow RT, Heller GZ, Barnsley L. The effect of 300 mW, 830 nm laser on chronic neck pain: a double-blind, randomized, placebo-controlled study. *Pain*. 2006; 124:201-10.

Hoving JL, et al. Manual therapy, physical therapy, or continued care by the general practitioner for patients with neck pain: long-term results from a pragmatic randomized clinical trial. *The Clinical Journal Of Pain*. 2006;22:370-7.

Tashiro M, Ogura T, Yanai K, et al. Cerebral metabolic changes in men after chiropractic spinal manipulation for neck pain. *Alternative Therapies In Health & Medicine* [serial online]. November 2011;17(6):12-17.

Dunn A, Green B, Formolo L, Chicoine D. Chiropractic Management for Veterans with Neck Pain: A Retrospective Study of Clinical Outcomes. *Journal Of Manipulative & Physiological Therapeutics* [serial online]. October 2011;34(8):533-538.

London W. Justification for chiropractic care to prevent recurrence of neck pain remains elusive. *Focus On Alternative & Complementary Therapies* [serial online]. March 2013;18(1):37-38.

Lawrence D. Spinal manipulation, medication or home exercise for acute and subacute neck pain. *Focus On Alternative & Complementary Therapies* [serial online]. December 2012;17(4):233-234.

Haavik H, Murphy B. Subclinical Neck Pain and the Effects of Cervical Manipulation on Elbow Joint Position Sense. *Journal Of Manipulative & Physiological Therapeutics* [serial online]. February 2011;34(2):88-97.

De camargo VM, Albuquerque-sendín F, Bérzin F, Stefanelli VC, De souza DP, Fernández-de-las-peñas C. Immediate effects on electromyographic activity and pressure pain thresholds after a cervical manipulation in mechanical neck pain: a randomized controlled trial. *J Manipulative Physiol Ther*. 2011;34(4):211-20.

Howell E. The association between neck pain, the Neck Disability Index and cervical ranges of motion: A narrative review. *Journal - Canadian Chiropractic Association*.. September 1, 2011;55(3):211.

Rodine R, Vernon H. Cervical radiculopathy: a systematic review on treatment by spinal manipulation and measurement with the Neck Disability Index. *Journal Of The Canadian Chiropractic Association*. March 2012;56(1):18-28.

Ssavoredra-Hernández M, Castro-Sánchez A, Fernández-de-las-Peñas C, Cleland J, Ortega-Santiago R, Arroyo-Morales M. Predictors for Identifying Patients With Mechanical Neck Pain Who Are Likely to Achieve Short-Term Success With Manipulative Interventions Directed at the Cervical and Thoracic Spine. *Journal Of Manipulative & Physiological Therapeutics*. March 2011;34(3):144-152.

Leaver A, Maher C, Refshauge K, et al. A Randomized Controlled Trial Comparing Manipulation With Mobilization for Recent Onset Neck Pain. *Archives Of Physical Medicine & Rehabilitation* September 2010;91(9):1313-1318.

Ernst E. Chiropractic spinal manipulation: what does the 'best' evidence show?. *Focus On Alternative & Complementary Therapies* [serial online]. December 2012;17(4):202-206.

Klein R, Bareis A, Schneider A, Linde K. Strain-counterstrain to treat restrictions of the mobility of the cervical spine in patients with neck pain-A sham-controlled randomized trial. *Complement Ther Med* 2013 Feb; 21(1): 1 - 7.

Humphreys BKH, Peterson CP, Comparison of outcomes in neck pain patients with and without dizziness undergoing chiropractic treatment: a prospective cohort study with 6-month follow-up. *ChiroMT* 2013 JAN 7; 21.

Kavlak BK, Bakar YB, Sari ZS, Investigation of the Efficacy of Different Physiotherapy Methods for Neck Pain. *J Musculoskel Pain* 2012 Dec; 20(4): 284 - 291.

Kay TM, Gross A, Goldsmith CH, Rutherford S, Voth S, Hoving JL, Brønfort G, Santaguida PL. Exercises for mechanical neck disorders. *Cochrane Database of Systematic Reviews* 2012, Issue 8. Art. No.: CD004250. DOI: 10.1002/14651858.CD004250.pub4

Clinical Pathway Feedback

CHP desires to keep our clinical pathways customarily updated. If you wish to provide additional input, please use the e-mail address listed below and identify which clinical pathway you are referencing. Thank you for taking the time to give us your comments.

Clinical Services Department: providers@chpgroup.com

¹ Hogg-Johnson S, et al. The burden and determinants of neck pain in the general population: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *J Manipulative Physiol Ther.* 2009 Feb;32(2 Suppl):S46-60.

² Weber Hellstenius S. Recurrent neck pain and headaches in preadolescents associated with mechanical, dysfunction of the cervical spine: A cross-sectional observational study with 131 students. *Journal Of Manipulative And Physiological Therapeutics.* October 1, 2009;32(8):625.

³ Ståhl M, et al. Neck pain in adolescence. A 4-year follow-up of pain-free preadolescents. *Pain.* July 2004; 110 (1): 427-431.

⁴ [Tsertsivadze A](#), [Clar C](#), [Court R](#), [Clarke A](#), [Mistry H](#), [Sutcliffe P](#). Cost-effectiveness of manual therapy for the management of musculoskeletal conditions: a systematic review and narrative synthesis of evidence from randomized controlled trials. *J Manipulative Physiol Ther.* 2014 Jul-Aug;37(6):343-62. doi: 10.1016/j.jmpt.2014.05.001. Epub 2014 Jun 27.

⁵ Council on Chiropractic Guidelines and Practice Parameters. Chiropractic management of neck related disorders. http://www.ccgpp.org/neck_related_disorders.pdf

⁶ Thiel, HW, Bolton J. Predictors for immediate and global responses to chiropractic manipulation of the cervical spine. [serial online]. *Journal of Manipulative and Physiological Therapeutics.* 2010; 33(5):A1-A6, 325-406.

⁷ Haldeman S, Carroll L, Cassidy JD, Schubert J, Nygren A. The Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders: executive summary. *J Manipulative Physiol Ther.* 2009;32(2 Suppl):S7-9.

⁸ [Bryans R](#), [Decina P](#), [Descarreaux M](#), [Duranleau M](#), [Marcoux H](#), [Potter B](#), [Ruegg RP](#), [Shaw L](#), [Watkin R](#), [White E](#). Evidence-based guidelines for the chiropractic treatment of adults with neck pain. *J Manipulative Physiol Ther.* 2014 Jan;37(1):42-63. doi: 10.1016/j.jmpt.2013.08.010. Epub 2013 Nov 19.

⁹ 1. Gross A, Kay TM, Paquin JP, Blanchette S, Lalonde P, Christie T, Dupont G, Graham N, Burnie SJ, Gellay G, Goldsmith CH, Forget M, Hoving JL, Brønfort G, Santaguida PL, Cervical Overview Group. Exercises for mechanical neck disorders. *Cochrane Database of Systematic Reviews* 2015, Issue 1. Art. No.: CD004250. DOI: 10.1002/14651858.CD004250.pub5.

¹⁰ [Maiers M](#), [Bronfort G](#), [Evans R](#), [Hartvigsen J](#), [Svendsen K](#), [Bracha Y](#), [Schulz C](#), [Schulz K](#), [Grimm R](#). Spinal manipulative therapy and exercise for seniors with chronic neck pain. *Spine J.* 2014 Sep 1;14(9):1879-89. doi: 10.1016/j.spinee.2013.10.035. Epub 2013 Nov 10.

¹¹ [Peterson CK](#), [Schmid C](#), [Leemann S](#), [Anklin B](#), [Humphreys BK](#). Outcomes from magnetic resonance imaging-confirmed symptomatic cervical disk herniation patients treated with high-velocity, low-amplitude spinal manipulative therapy: a prospective cohort study with 3-month follow-up. *J Manipulative Physiol Ther.* 2013 Oct;36(8):461-7. doi: 10.1016/j.jmpt.2013.07.002. Epub 2013 Aug 12.

¹² [Zhu L](#), [Wei X](#), [Wang S](#). Does cervical spine manipulation reduce pain in people with degenerative cervical radiculopathy? A systematic review of the evidence, and a meta-analysis. *Clin Rehabil.* 2015 Feb 13. pii: 0269215515570382. [Epub ahead of print].

¹³ Gross A, et al. Manipulation or mobilisation for neck pain. *Cochrane Database Syst Rev.* 2010 Jan 20;(1):CD004249.

¹⁴ Leininger BD, Evans R, Bronfort G. Exploring patient satisfaction: a secondary analysis of a randomized clinical trial of spinal manipulation, home exercise, and medication for acute and subacute neck pain. *J Manipulative Physiol Ther.* 2014;37(8):593-601.

-
- ¹⁵ [Southerst D](#), [Nordin MC](#), [Côté P](#), [Shearer HM](#), [Varatharajan S](#), [Yu H](#), [Wong JJ](#), [Sutton DA](#), [Randhawa KA](#), [van der Velde GM](#), [Mior SA](#), [Carroll LJ](#), [Jacobs CL](#), [Taylor-Vaisey AL](#). Is exercise effective for the management of neck pain and associated disorders or whiplash-associated disorders? A systematic review by the Ontario Protocol for Traffic Injury Management (OPTIMA) Collaboration. *Spine J*. 2014 Feb 15. pii: S1529-9430(14)00210-1. doi: 10.1016/j.spinee.2014.02.014. [Epub ahead of print].
- ¹⁶ Sherman KJ, Cook AJ, Wellman RD, et al. Five-week outcomes from a dosing trial of therapeutic massage for chronic neck pain. *Ann Fam Med*. 2014;12(2):112-20.
- ¹⁷ Haas M, Spegman A, Peterson D, Aickin M, Vavrek D. Dose response and efficacy of spinal manipulation for chronic cervicogenic headache: a pilot randomized controlled trial. *Spine J*. 2010 Feb;10(2):117-28.
- ¹⁸ Cassidy J, Boyle E, Cote P, et al. Risk of vertebrobasilar stroke and chiropractic care: Results of a population-based case-control and case-crossover study. *Journal Of Manipulative And Physiological Therapeutics* February 1, 2009;32(2 SUPPL):S201.
- ¹⁹ Rubinstein SM, et al. The benefits outweigh the risks for patients undergoing chiropractic care for neck pain: a prospective, multicenter, cohort study. *Journal Of Manipulative And Physiological Therapeutics*. 2007; 30:408-18.
- ²⁰ Rubinstein S. Adverse events following chiropractic care for subjects with neck or low-back pain: Do the benefits outweigh the risks? *Journal Of Manipulative And Physiological Therapeutics*. July 1, 2008;31(6):461.