

UNIVERSITY OF CALIFORNIA SAN DIEGO

## **ECE 271 Notes**

Instructor: *Prof. Nuno Vasconcelos*

Organized by Ray Tsai

## Bayes Decision Rule

$$\begin{aligned}
 g^*(x) &= \arg \min_{g(x)} \sum_i P_{Y|X}(i|x) L[g(x), i] \\
 &= \arg \max_i P_{Y|X}(i|x) && \text{(for 0-1 loss function)} \\
 &= \arg \max_i P_{X|Y}(x|i) P_Y(i) && \text{(for 0-1 loss function)} \\
 &= \arg \max_i \log P_{X|Y}(x|i) + \log P_Y(i). && \text{(for 0-1 loss function)}
 \end{aligned}$$

For binary classification, the likelihood ratio form is: pick 0 if  $\frac{P_{X|Y}(x|0)}{P_{X|Y}(x|1)} > T^* = \frac{P_Y(1)}{P_Y(0)}$ .

## Associated Risk

$$R^* = \int P_X(x) \sum_{i \neq g^*(x)} P_{Y|X}(i|x) dx = \int P_{Y,X}(y \neq g^*(x), x) dx \quad \text{(For 0-1 loss function)}$$

## Gaussian Classifier

For single variable, we assume  $\sigma_i = \sigma$  and pick 0 if

$$x < \frac{\mu_1 + \mu_0}{2} + \frac{1}{\frac{\mu_1 - \mu_0}{\sigma^2}} \log \frac{P_Y(0)}{P_Y(1)}.$$

Generalizing it to multiple variables, we assume  $\Sigma_i = \Sigma$ , then the BDR becomes

$$i^*(x) = \arg \min_i [d(x, \mu_i) + \alpha_i],$$

where  $d(x, y) = (x - y)^T \Sigma^{-1} (x - y)$  and  $\alpha_i = \log \left[ \frac{(2\pi)^d |\Sigma|}{P_Y(i)} \right] - 2 \log P_Y(i)$ .

Alternatively,

$$i^*(x) = \arg \max_i g_i(x),$$

where  $g_i(x) = w_i^T x + w_{i0}$ ,  $w_i = \Sigma^{-1} \mu_i$ , and  $w_{i0} = -\frac{1}{2} \mu_i^T \Sigma^{-1} \mu_i + \log P_Y(i)$ .

## Geometric Interpretation

Thus, the hyperplane between class 0 and 1 is

$$g_0(x) - g_1(x) = w^T x + b = 0,$$

where  $w = \Sigma^{-1}(\mu_0 - \mu_1)$  and  $b = -\frac{(\mu_0 + \mu_1)^T \Sigma^{-1}(\mu_0 - \mu_1)}{2} + \log \frac{P_Y(0)}{P_Y(1)}$ .

It could also be rewritten as

$$w^T (x - x_0) = 0,$$

where  $w = \Sigma^{-1}(\mu_0 - \mu_1)$  and  $x_0 = \frac{\mu_0 + \mu_1}{2} - \frac{1}{(\mu_0 - \mu_1)^T \Sigma^{-1} (\mu_0 - \mu_1)} \log \frac{P_Y(0)}{P_Y(1)} (\mu_0 - \mu_1)$

## Gaussian Distribution Transformation

Let  $x \sim N(\mu, \Sigma)$ , and let  $y = A^T x$ , for some matrix  $A$ . Then,  $y \sim N(A^T \mu, A^T \Sigma A)$ . A special case of this is the whitening transform  $A_w = \Phi \Lambda^{-1/2}$ , where  $\Phi$  is the matrix of orthonormal eigenvectors of  $\Sigma$ , and  $\Lambda$  is the diagonal matrix of eigenvalues of  $\Sigma$ .

## Sigmoid

Suppose that  $g_1(x) = 1 - g_0(x)$ . Then, we can rewrite

$$g_0(x) = \frac{1}{1 + \frac{P_{X|Y}(x|1)P_Y(1)}{P_{X|Y}(x|0)P_Y(0)}} = \frac{1}{1 + \exp\{d_0(x, \mu_0) - d_1(x, \mu_1) + \alpha_0 - \alpha_1\}},$$

where,  $d(x, y) = (x - y)^T \Sigma^{-1} (x - y)$  and  $\alpha_i = \log[(2\pi)^d |\Sigma_i|] - 2 \log P_Y(i)$ .

## Maximum Likelihood Estimation

Solve for

$$\Theta^* = \arg \max_{\Theta} P_X(D; \Theta) = \arg \max_{\Theta} \log P_X(D; \Theta).$$

Consider the Gaussian example:

Given a sample  $\mathcal{D} = \{x_1, \dots, x_n\}$  of independent points, where  $P_X(x_i) = \frac{1}{\sqrt{(2\pi)^d |\Sigma|}} e^{-\frac{1}{2}(x_i - \mu)^T \Sigma^{-1} (x_i - \mu)}$ .

Then, the likelihood  $L(x_1, \dots, x_n | \mu, \sigma) = \prod_{i=1}^n P_X(x_i)$ . We take the gradient of the natural log of  $L$  with respect to  $\mu$  and get

$$\begin{aligned} \nabla_{\mu}(\log L) &= \nabla_{\mu} \left( -\frac{1}{2} \log[(2\pi)^d |\Sigma|] - \frac{1}{2} \sum_{i=1}^n (x_i - \mu)^T \Sigma^{-1} (x_i - \mu) \right) \\ &= \sum_{i=1}^n \Sigma^{-1} (x_i - \mu) = \sum_{i=1}^n x_i - \sum_{i=1}^n \mu = 0. \end{aligned}$$

Thus, we get  $\hat{\mu} = \frac{1}{n} \sum_{i=1}^n x_i$ . By taking the Hessian, we get  $\nabla_{\mu}^2(\log L) = -\sum_{i=1}^n \Sigma^{-1} = -n\Sigma^{-1}$ . Since the covariance matrix  $\Sigma$  is positive definite,  $-n\Sigma^{-1}$  is negative definite. Thus  $\hat{\mu}$  is the maximum point.

In addition, the MLE of the covariance matrix is

$$\hat{\Sigma} = \frac{1}{n} \sum_{i=1}^n (x_i - \mu)(x_i - \mu)^T.$$

## Bias and Variance

$$\begin{aligned} \text{Bias}(\hat{\theta}) &= E[\hat{\theta} - \theta], \quad \text{Var}(\hat{\theta}) = E\left\{(\hat{\theta} - E[\hat{\theta}])^2\right\}, \\ \text{MSE}(\hat{\theta}) &= E\left[(\hat{\theta} - \theta)^2\right] = \text{Var}(\hat{\theta}) + \text{Bias}^2(\hat{\theta}). \end{aligned}$$

## Least Squares

Consider an overdetermined system  $\Phi\theta = z$ , where we attempt to minimize  $\|z - \Phi\theta\|$ , the least square solution is

$$\theta^* = (\Phi^T \Phi)^{-1} \Phi^T z$$

For an overdetermined system  $W\Phi\theta = Wz$ , where we attempt to minimize  $(z - \Phi\theta)^T W^T W (z - \Phi\theta)$ , the least square solution is

$$\theta^* = (\Phi^T W^T W \Phi)^{-1} \Phi^T W^T W z.$$