

# Double Turán Problem

Ray Tsai

May 12, 2025

# Overview

- 1 Turán Problem
- 2 Double Turán Problem
- 3 Induced Double Turán Problem
- 4 Proof of Theorem B
- 5 Proof of Theorem D
- 6 Concluding Remarks

# Turán Problem

## Turán Problem

Given a graph  $F$ , how many edges can an  $n$ -vertex graph have while containing no copy of  $F$  as a subgraph?

# Turán Problem

## Turán Problem

Given a graph  $F$ , how many edges can an  $n$ -vertex graph have while containing no copy of  $F$  as a subgraph?

## Turán Number

$$\text{ex}(n, F) := \max\{e(G) : |V(G)| = n \text{ and } F \not\subseteq G\}$$

# Turán Problem

## Turán's Theorem [Tur41]

For  $r + 1 \geq 3$ ,

$$\text{ex}(n, K_{r+1}) = e(T_r(n)),$$

with equality for graph  $G$  only if  $G = T_r(n)$ .





Figure:  $T_4(13)$

# Erdős-Stone Theorem

Erdős-Stone Theorem [ES46], Simonovits' Theorem [ES66]


Let  $F$  be any graph of chromatic number  $r + 1 \geq 2$ . Then  $\text{ex}(n, F) = e(T_r(n)) + o(n^2)$  as  $n \rightarrow \infty$ .



# Erdős-Stone Theorem

## Erdős-Stone Theorem [ES46], Simonovits' Theorem [ES66]

Let  $F$  be any graph of chromatic number  $r + 1 \geq 2$ . Then  $\text{ex}(n, F) = e(T_r(n)) + o(n^2)$  as  $n \rightarrow \infty$ .



## Supersaturation Theorem [ES83]

For all  $\epsilon > 0$ , there exists  $\delta > 0$  such that if  $G$  is a  $n$ -vertex graph with  $\text{ex}(n, F) + \epsilon n^2$  edges, then  $G$  contains  $\delta n^{v(F)}$  copies of  $F$ .

# Double Turán Problem: Definition

What is the double Turán problem?



# Double Turán Problem: Definition

What is the double Turán problem?

Let  $G_1, G_2, \dots, G_m$  be graphs on  $[n]$  with  $E(F) \not\subseteq E(G_i) \cap E(G_j)$  for distinct  $i, j \in [m]$ .

# Double Turán Problem: Definition

What is the double Turán problem?

Let  $G_1, G_2, \dots, G_m$  be graphs on  $[n]$  with  $E(F) \not\subseteq E(G_i) \cap E(G_j)$  for distinct  $i, j \in [m]$ . (**Double  $F$ -free**)

# Double Turán Problem: Definition

What is the double Turán problem?

Let  $G_1, G_2, \dots, G_m$  be graphs on  $[n]$  with  $E(F) \not\subseteq E(G_i) \cap E(G_j)$  for distinct  $i, j \in [m]$ . (**Double  $F$ -free**)

## Double Turán Problem

What is the value of  $\phi(m, n, F) = \max \sum_{i=1}^m |G_i|$ ?

# Double Turán Problem: Definition

What is the double Turán problem?

Let  $G_1, G_2, \dots, G_m$  be graphs on  $[n]$  with  $E(F) \not\subseteq E(G_i) \cap E(G_j)$  for distinct  $i, j \in [m]$ . (**Double  $F$ -free**)

## Double Turán Problem

What is the value of  $\phi(m, n, F) = \max \sum_{i=1}^m |G_i|$ ?

$$\binom{m}{2} \geq \phi(m, n, F) \geq \binom{n}{2} + (m-1)\text{ex}(n, F)$$

# Double Turán Problem: Definition

What is the double Turán problem?

Let  $G_1, G_2, \dots, G_m$  be graphs on  $[n]$  with  $E(F) \not\subseteq E(G_i) \cap E(G_j)$  for distinct  $i, j \in [m]$ . (**Double  $F$ -free**)

## Double Turán Problem

What is the value of  $\phi(m, n, F) = \max \sum_{i=1}^m G_i$ ?

$$\binom{m}{2} \geq \phi(m, n, F) \geq \binom{n}{2} + (m-1)\text{ex}(n, F)$$

Is this tight?

# Double Turán Problem: Main Result

## Theorem A

Let  $n \geq 1$  and  $F$  be a non-bipartite graph. Then as  $m \rightarrow \infty$

$$\phi(m, n, F) = (1 + o(1))m \cdot \text{ex}(n, F).$$

# Double Turán Problem: Main Result

## Theorem A

Let  $n \geq 1$  and  $F$  be a non-bipartite graph. Then as  $m \rightarrow \infty$

$$\phi(m, n, F) = (1 + o(1))m \cdot \text{ex}(n, F).$$

There are  $\leq n^{v(F)}$  copies of  $F$  over all  $G_i$ .

# Double Turán Problem: Main Result

## Theorem A

Let  $n \geq 1$  and  $F$  be a non-bipartite graph. Then as  $m \rightarrow \infty$

$$\phi(m, n, F) = (1 + o(1))m \cdot \text{ex}(n, F).$$

There are  $\leq n^{v(F)}$  copies of  $F$  over all  $G_i$ .

## Supersaturation Theorem [ES83]

For all  $\epsilon > 0$ , there exists  $\delta > 0$  such that if  $G$  is a  $n$ -vertex graph with  $\text{ex}(n, F) + \epsilon n^2$  edges, then  $G$  contains  $\delta n^{v(F)}$  copies of  $F$ .



# Double Turán Problem: Main Results

The case for  $F = K_r$  can be reduced to a finite computation.

# Double Turán Problem: Main Results

The case for  $F = K_r$  can be reduced to a finite computation.

$$\phi(3, n, K_3) = \binom{n}{2} + \lfloor n/2 \rfloor$$

# Double Turán Problem: Main Results

The case for  $F = K_r$  can be reduced to a finite computation.

$$\phi(3, n, K_3) = \binom{n}{2} + \lfloor n/2 \rfloor$$

But

$$\lim_{n \rightarrow \infty} \frac{\phi(4, n, K_3)}{\binom{n}{2} + 3\lfloor n/4 \rfloor} > 1$$

# Main Results




Figure: Construction for four graphs not containing a double- $K_3$

# Double Turán Problem: Main Result

## Theorem B (Wilson)

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

# Double Turán Problem: Main Result

## Theorem B (Wilson)

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

By the Erdős-Stone theorem, this only applies to certain bipartite graphs.

# Double Turán Problem: Main Result


## Theorem B (Wilson)

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

By the Erdős-Stone theorem, this only applies to certain bipartite graphs.

e.g. The extremal graph for  $P_2$  is a matching, which has maximum degree 1 and  $\text{ex}(n, P_2) = \lfloor n/2 \rfloor$ .



# Double Turán Problem: Motivation

Double Turán problems are closely related to Turán problems for 3-uniform hypergraphs  $H$  through **link graphs**.



# Double Turán Problem: Motivation

Double Turán problems are closely related to Turán problems for 3-uniform hypergraphs  $H$  through **link graphs**.

## Link Graph

For  $i \in V(H)$ , define graph  $H_i$  with

$$V(H_i) = V(H) \setminus \{i\} \quad \text{and} \quad E(H_i) = \{\{j, k\} : \{i, j, k\} \in E(H)\}.$$

# Double Turán Problem: Motivation

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $K_{2,2,2}^{(3)}$

# Double Turán Problem: Motivation

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $K_{2,2,2}^{(3)}$

Conjecture [Erd64]

$$\text{ex}(n, K_{2,2,2}^{(3)}) = \Theta(n^{11/4})$$

# Double Turán Problem: Motivation

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $K_{2,2,2}^{(3)}$

Conjecture [Erd64]

$$\text{ex}(n, K_{2,2,2}^{(3)}) = \Theta(n^{11/4})$$

# Double Turán Problem: Motivation

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $K_{2,2,2}^{(3)}$

Conjecture [Erd64]

$$\text{ex}(n, K_{2,2,2}^{(3)}) = \Theta(n^{11/4})$$

# Double Turán Problem: Motivation

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $K_{2,2,2}^{(3)}$

Conjecture [Erd64]

$$\text{ex}(n, K_{2,2,2}^{(3)}) = \Theta(n^{11/4})$$

# Double Turán Problem: Motivation

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $K_{2,2,2}^{(3)}$

Conjecture [Erd64]

$$\text{ex}(n, K_{2,2,2}^{(3)}) = \Theta(n^{11/4})$$

$H$  is octahedron-free  $\implies H_1, H_2, \dots, H_n$  are double  $K_{2,2}$ -free.

$$\text{ex}(n, K_{2,2,2}^{(3)}) \leq \phi(n, n, K_{2,2})$$

# Double Turán Problem: Motivation

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $K_{2,2,2}^{(3)}$

Conjecture [Erd64]

$$\text{ex}(n, K_{2,2,2}^{(3)}) = \Theta(n^{11/4})$$

$H$  is octahedron-free  $\implies H_1, H_2, \dots, H_n$  are double  $K_{2,2}$ -free.

$$\text{ex}(n, K_{2,2,2}^{(3)}) = \Theta(\phi(n, n, K_{2,2}))$$



# Induced Double Turán Problem: Definition

What is the **induced** double Turán problem?

# Induced Double Turán Problem: Definition

What is the **induced** double Turán problem?

## Induced Graphs

We call graphs  $G_1, G_2, \dots, G_m$  **induced** if each  $G_i$  is an induced subgraph of  $\bigcup_{i=1}^m G_i$ .

# Induced Double Turán Problem: Definition

What is the **induced** double Turán problem?

## Induced Graphs

We call graphs  $G_1, G_2, \dots, G_m$  **induced** if each  $G_i$  is an induced subgraph of  $\bigcup_{i=1}^m G_i$ .

In other words, if  $\{u, v\} \in E(G_i)$  and  $u, v \in V(G_j)$ , then  $\{u, v\} \in E(G_j)$ .

# Induced Double Turán Problem: Definition

What is the **induced** double Turán problem?

## Induced Graphs

We call graphs  $G_1, G_2, \dots, G_m$  **induced** if each  $G_i$  is an induced subgraph of  $\bigcup_{i=1}^m G_i$ .

In other words, if  $\{u, v\} \in E(G_i)$  and  $u, v \in V(G_j)$ , then  $\{u, v\} \in E(G_j)$ .

Let graphs  $G_1, G_2, \dots, G_m$  be induced and double  $F$ -free.

## Induced Double Turán Problem

What is the value of  $\phi^*(m, n, F) = \max \sum_{i=1}^m e(G_i)$ ?

# Induced Double Turán Problem: Definition

What is the **induced** double Turán problem?

## Induced Graphs

We call graphs  $G_1, G_2, \dots, G_m$  **induced** if each  $G_i$  is an induced subgraph of  $\bigcup_{i=1}^m G_i$ .

In other words, if  $\{u, v\} \in E(G_i)$  and  $u, v \in V(G_j)$ , then  $\{u, v\} \in E(G_j)$ .

Let graphs  $G_1, G_2, \dots, G_m$  be induced and double  $F$ -free.

## Induced Double Turán Problem

What is the value of  $\phi^*(m, n, F) = \max \sum_{i=1}^m e(G_i)$ ?

$$\phi(m, n, F) \geq \phi^*(m, n, F)$$

# Induced Double Turán Problem: Conjecture

## Induced Double Turán Problem

What is the value of  $\phi^*(m, n, F) = \max \sum_{i=1}^m e(G_i)$ ?

$$\phi^*(m, n, F) \geq \max \left\{ \binom{n}{2} + \binom{v(F) - 1}{2} (m - 1), m \cdot \text{ex}(n, F) \right\}.$$

# Induced Double Turán Problem: Conjecture

## Induced Double Turán Problem

What is the value of  $\phi^*(m, n, F) = \max \sum_{i=1}^m e(G_i)$ ?

$$\phi^*(m, n, F) \geq \max \left\{ \binom{n}{2} + \binom{v(F)-1}{2} (m-1), m \cdot \text{ex}(n, F) \right\}.$$

Is it tight?

# Induced Double Turán Problem: Conjecture

## Induced Double Turán Problem

What is the value of  $\phi^*(m, n, F) = \max \sum_{i=1}^m e(G_i)$ ?

$$\phi^*(m, n, F) \geq \max \left\{ \binom{n}{2} + \binom{v(F)-1}{2} (m-1), m \cdot \text{ex}(n, F) \right\}.$$

Is it tight?

## Conjecture

Let  $F$  be any non-empty graph and  $m, n \geq 1$ . Then

$$\phi^*(m, n, F) = \Theta(m \cdot \text{ex}(n, F) + n^2).$$



## Theorem C

For  $m \geq 3$  and non-bipartite  $F$ , if  $n$  is large enough, then

$$\phi^*(m, n, F) = m \cdot \text{ex}(n, F),$$

with equality only for identical extremal  $n$ -vertex  $F$ -free graphs.

## Theorem C

For  $m \geq 3$  and non-bipartite  $F$ , if  $n$  is large enough, then

$$\phi^*(m, n, F) = m \cdot \text{ex}(n, F),$$

with equality only for identical extremal  $n$ -vertex  $F$ -free graphs.

$\implies$  Conjecture is true for non-bipartite  $F$ .

## Theorem C

For  $m \geq 3$  and non-bipartite  $F$ , if  $n$  is large enough, then

$$\phi^*(m, n, F) = m \cdot \text{ex}(n, F),$$

with equality only for identical extremal  $n$ -vertex  $F$ -free graphs.

$\implies$  Conjecture is true for non-bipartite  $F$ .

For  $F = K_r$ , this theorem applies for all  $n$ .

# Induced Double Turán Problem: Motivation

## Generalized Turán problem

What is the maximum number  $\text{ex}(n, F, K_3)$  of triangles in an  $n$ -vertex  $F$ -free graph  $G$ ?

Studied by Alon and Shikhelman [AS16] and Kostochka, Mubayi and Verstraete [KMV15; MM23; MV16].

# Induced Double Turán Problem: Motivation

## Generalized Turán problem

What is the maximum number  $\text{ex}(n, F, K_3)$  of triangles in an  $n$ -vertex  $F$ -free graph  $G$ ?

Studied by Alon and Shikhelman [AS16] and Kostochka, Mubayi and Verstraete [KMV15; MM23; MV16].

For  $i \in V(G)$ , define  $G_i$  with

$$V(G_i) = V(G) \quad \text{and} \quad E(G_i) = \{\{j, k\} : \{i, j\}, \{j, k\}, \{i, k\} \in E(G)\}$$

# Induced Double Turán Problem: Motivation

Ex. Octahedron-free graph  $G$ .




Figure: Octahedron




Figure: Octahedron Graph  $K_{2,2,2}$

# Induced Double Turán Problem: Motivation

Ex. Octahedron-free graph  $G$ .




Figure: Octahedron




Figure: Octahedron Graph  $K_{2,2,2}$

# Induced Double Turán Problem: Motivation

Ex. Octahedron-free graph  $G$ .




Figure: Octahedron




Figure: Octahedron Graph  $K_{2,2,2}$



# Induced Double Turán Problem: Motivation

Ex. Octahedron-free graph  $G$ .




Figure: Octahedron




Figure: Octahedron Graph  $K_{2,2,2}$

# Induced Double Turán Problem: Motivation

Ex. Octahedron-free graph  $G$ .




Figure: Octahedron




Figure: Octahedron Graph  $K_{2,2,2}$

$G$  is  $K_{2,2,2}$ -free  $\implies G_1, G_2, \dots, G_n$  are induced and double  $K_{2,2}$ -free.

$$\text{ex}(n, K_{2,2,2}, K_3) \leq \phi^*(n, n, K_{2,2})$$

# Induced Double Turán Problem: Motivation

Ex. Octahedron-free graph  $G$ .




Figure: Octahedron




Figure: Octahedron Graph  $K_{2,2,2}$

$G$  is  $K_{2,2,2}$ -free  $\implies G_1, G_2, \dots, G_n$  are induced and double  $K_{2,2}$ -free.

$$\text{ex}(n, K_{2,2,2}, K_3) = \Theta(\phi^*(n, n, K_{2,2}))$$

# Induced Double Turán Problem: Motivation

## Conjecture

Let  $F$  be any non-empty graph and  $m, n \geq 1$ . Then

$$\phi^*(m, n, F) = \Theta(m \cdot \text{ex}(n, F) + n^2).$$

Conjecture implies

$$\text{ex}(n, K_{2,2,2}, K_3) = \Theta(\phi^*(n, n, K_{2,2})) = \Theta(n^{5/2}).$$

## Theorem D

Let  $P$  be a path with two edges. Then  $\phi(n, n, P) = \Omega(n^{5/2})$ , whereas  $\phi^*(n, n, P) = o(n^{5/2})$ , as  $n \rightarrow \infty$ . In particular,

$$\lim_{n \rightarrow \infty} \frac{\phi^*(n, n, P)}{\phi(n, n, P)} = 0.$$

## Theorem D

Let  $P$  be a path with two edges. Then  $\phi(n, n, P) = \Omega(n^{5/2})$ , whereas  $\phi^*(n, n, P) = o(n^{5/2})$ , as  $n \rightarrow \infty$ . In particular,

$$\lim_{n \rightarrow \infty} \frac{\phi^*(n, n, P)}{\phi(n, n, P)} = 0.$$

This shows that  $\phi(m, n, F)$  and  $\phi^*(m, n, F)$  can be very different problems.

# Proof of Theorem B: Upper Bound

## Theorem B (Wilson)

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

# Proof of Theorem B: Upper Bound

## Theorem B (Wilson)

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

We need to show

$$\phi(m, n, F) \leq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$



# Proof of Theorem B: Upper Bound

## Theorem B (Wilson)

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

We need to show

$$\phi(m, n, F) \leq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

Let  $E_S$  be the set of edges in exactly  $\{G_i\}_{i \in S}$ .

# Proof of Theorem B: Upper Bound

## Theorem B (Wilson)

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

We need to show

$$\phi(m, n, F) \leq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

Let  $E_S$  be the set of edges in exactly  $\{G_i\}_{i \in S}$ .

$$\implies \sum_{i=1}^m e(G_i) = \sum_{S \subseteq [m]} |S| |E_S| \leq \binom{n}{2} + \sum_{S \subseteq [m], |S| \geq 2} (|S| - 1) |E_S|.$$

# Proof of Theorem B: Upper Bound

$$\sum_{\substack{S \subseteq [m] \\ |S| \geq 2}} (|S| - 1) |E_S| = \sum_{\substack{S \subseteq [m] \\ |S|=2}} \sum_{T \supseteq S} \frac{(|T| - 1) |E_T|}{\binom{|T|}{2}} \leq \sum_{\substack{S \subseteq [m] \\ |S|=2}} \sum_{T \supseteq S} |E_T|,$$

as each  $T \in [m]$  with  $|T| \geq 2$  is counted  $\binom{|T|}{2}$  times.

# Proof of Theorem B: Upper Bound

Observation: If  $|S| \geq 2$ , the edge set  $\bigcup_{T \supseteq S} E_T$  is  $F$ -free.

# Proof of Theorem B: Upper Bound

Observation: If  $|S| \geq 2$ , the edge set  $\bigcup_{T \supseteq S} E_T$  is  $F$ -free.

$$\left| \bigcup_{T \supseteq S} E_T \right| = \sum_{T \supseteq S} |E_T| \leq \text{ex}(n, F)$$

# Proof of Theorem B: Upper Bound

Observation: If  $|S| \geq 2$ , the edge set  $\bigcup_{T \supseteq S} E_T$  is  $F$ -free.

$$\left| \bigcup_{T \supseteq S} E_T \right| = \sum_{T \supseteq S} |E_T| \leq \text{ex}(n, F)$$

$$\Rightarrow \sum_{\substack{S \subseteq [m] \\ |S| \geq 2}} (|S| - 1) |E_S| \leq \sum_{\substack{S \subseteq [m], \\ |S|=2}} \sum_{T \supseteq S} |E_T| \leq \binom{m}{2} \text{ex}(n, F)$$

This proves the upper bound.

# Proof of Theorem B: Lower Bound

We need to show

$$\phi(m, n, F) \geq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

# Proof of Theorem B: Lower Bound

We need to show

$$\phi(m, n, F) \geq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

For  $1 \leq i < j \leq m$ , let  $H_{ij}$  be an extremal  $F$ -free graph on  $[n]$  with maximum degree  $\Delta(H_{ij}) \leq \sqrt{n}/m^2$



# Proof of Theorem B: Lower Bound

We need to show

$$\phi(m, n, F) \geq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

For  $1 \leq i < j \leq m$ , let  $H_{ij}$  be an extremal  $F$ -free graph on  $[n]$  with maximum degree  $\Delta(H_{ij}) \leq \sqrt{n}/m^2$

$\implies$  We can pack these  $\binom{m}{2}$   $H_{ij}$ 's into  $[n]$  s.t.  $E(H_{i_1j_1}) \cap E(H_{i_2j_2}) = \emptyset$

# Proof of Theorem B: Lower Bound

For  $i \in [m]$ , put

$$G_i = H_{1i} \cup H_{2i} \cup \cdots \cup H_{im}$$

# Proof of Theorem B: Lower Bound

For  $i \in [m]$ , put

$$G_i = H_{1i} \cup H_{2i} \cup \cdots \cup H_{im}$$

Then  $G_i \cap G_j = H_{ij} \implies F$ -free

# Proof of Theorem B: Lower Bound

For  $i \in [m]$ , put

$$G_i = H_{1i} \cup H_{2i} \cup \cdots \cup H_{im}$$

Then  $G_i \cap G_j = H_{ij} \implies F$ -free

Add edges not in any of  $H_{ij}$  to  $G_1$ .

$$\sum_{i=1}^m e(G_i) = 2 \underbrace{\sum e(H_{ij})}_{\binom{m}{2} \text{ex}(n, F)} + \underbrace{\left| E(K_n) \setminus \bigcup E(H_{ij}) \right|}_{\binom{n}{2} - \binom{m}{2} \text{ex}(n, F)} = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F)$$

## Theorem D

Let  $m, n, r \geq 3$ . Then

$$\phi^*(m, n, K_r) = m \cdot e(T_{r-1}(n)),$$

with equality for induced  $K_r$ -free graphs  $G_1, G_2, \dots, G_m$  only if  $G_1 = G_2 = \dots = G_m = T_{r-1}(n)$ .

## Theorem D


Let  $m, n, r \geq 3$ . Then

$$\phi^*(m, n, K_r) = m \cdot e(T_{r-1}(n)),$$


with equality for induced  $K_r$ -free graphs  $G_1, G_2, \dots, G_m$  only if  $G_1 = G_2 = \dots = G_m = T_{r-1}(n)$ .

IDEA: Reduce to  $m = 2$ .

# Proof of Theorem D




# Proof of Theorem D



$G_1 \cap G_2$  (gray area) is double  $K_r$ -free.



# Proof of Theorem D




$G_1 \cap G_2$  (gray area) is double  $K_r$ -free.

If  $G_1, G_2$  intersects in  $t$  vertices, then

$$e(G_1) + e(G_2) \leq \binom{n-t}{2} + (n-t)t + 2T_{r-1}(t)$$

# Proof of Theorem D




$G_1 \cap G_2$  (gray area) is double  $K_r$ -free.

If  $G_1, G_2$  intersects in  $t$  vertices, then


$$e(G_1) + e(G_2) \leq \binom{n-t}{2} + (n-t)t + 2T_{r-1}(t)$$

$\implies$  we are done if the unique maximum occurs at  $t = n$ .

# Proof of Theorem D



# Proof of Theorem D




Put  $f(t, r) := \binom{n-t}{2} + (n-t)t + 2T_{r-1}(t)$

For example, if  $r = 4$ ,

$$f(t+1, 4) - f(t, 4) = -t + 1 + 2 \underbrace{[T_3(t+1) - T_3(t)]}_{t - \left\lceil \frac{t}{2} \right\rceil} = t + 1 - 2 \left\lceil \frac{t}{2} \right\rceil > 0$$

# Proof of Theorem D




Put  $f(t, r) := \binom{n-t}{2} + (n-t)t + 2T_{r-1}(t)$

For example, if  $r = 4$ ,

$$f(t+1, 4) - f(t, 4) = -t + 1 + 2 \underbrace{[T_3(t+1) - T_3(t)]}_{t - \lceil \frac{t}{2} \rceil} = t + 1 - 2 \left\lceil \frac{t}{2} \right\rceil > 0$$

$\implies$  The larger the intersection, the larger the sum of edges.

# Proof of Theorem D



$$\text{Put } f(t, r) := \binom{n-t}{2} + (n-t)t + 2T_{r-1}(t)$$

For example, if  $r = 4$ ,

$$f(t+1, 4) - f(t, 4) = -t + 1 + 2 \underbrace{[T_3(t+1) - T_3(t)]}_{t - \lceil \frac{t}{2} \rceil} = t + 1 - 2 \left\lceil \frac{t}{2} \right\rceil > 0$$

$\implies$  The larger the intersection, the larger the sum of edges.

$\implies$  The best we can do is  $G_1 = G_2 = T_3(n)$ .

# Concluding Remarks: Induced Double Turán Problem

## Conjecture

Let  $F$  be any non-empty graph and  $m, n \geq 1$ . Then

$$\phi^*(m, n, F) = \Theta(m \cdot \text{ex}(n, F) + n^2).$$


# Concluding Remarks: Induced Double Turán Problem

## Conjecture

Let  $F$  be any non-empty graph and  $m, n \geq 1$ . Then

$$\phi^*(m, n, F) = \Theta(m \cdot \text{ex}(n, F) + n^2).$$

This conjecture is broadly open for bipartite graphs.






# Concluding Remarks: Induced Double Turán Problem

## Conjecture

Let  $F$  be any non-empty graph and  $m, n \geq 1$ . Then

$$\phi^*(m, n, F) = \Theta(m \cdot \text{ex}(n, F) + n^2).$$

This conjecture is broadly open for bipartite graphs.



Solving the conjecture would give a solution to  $\text{ex}(n, K_{2,2,2}, K_3)$ .

# Concluding Remarks: Double Turán Problem

The non-bipartite case is likely intractable.

# Concluding Remarks: Double Turán Problem

The non-bipartite case is likely intractable.

We only know  $\phi(3, n, K_3) = \binom{n}{2} + \lfloor n/2 \rfloor$ . The rest is open.

# Concluding Remarks: Double Turán Problem

The non-bipartite case is likely intractable.

We only know  $\phi(3, n, K_3) = \binom{n}{2} + \lfloor n/2 \rfloor$ . The rest is open.

$$\lim_{n \rightarrow \infty} \frac{\phi(4, n, K_3)}{m\binom{n}{2}} = ??$$

- [AS16] Noga Alon and Clara Shikhelman. “Many  $T$  copies in  $H$ -free graphs”. In: *J. Combin. Theory Ser. B* 121 (2016), pp. 146–172. ISSN: 0095-8956,1096-0902. DOI: 10.1016/j.jctb.2016.03.004. URL: <https://doi.org/10.1016/j.jctb.2016.03.004>.
- [Erd64] P. Erdős. “On extremal problems of graphs and generalized graphs”. In: *Israel Journal of Mathematics* 2.3 (1964), pp. 183–190. DOI: 10.1007/BF02759942. URL: <https://doi.org/10.1007/BF02759942>.
- [ES46] P. Erdős and A. H. Stone. “On the structure of linear graphs”. In: *Bull. Amer. Math. Soc.* 52 (1946), pp. 1087–1091. ISSN: 0002-9904. DOI: 10.1090/S0002-9904-1946-08715-7. URL: <https://doi.org/10.1090/S0002-9904-1946-08715-7>.

- [ES66] P. Erdős and M. Simonovits. “A limit theorem in graph theory”. In: *Studia Sci. Math. Hungar* 1 (1966), pp. 51–57. ISSN: 0081-6906.
- [ES83] Paul Erdős and Miklós Simonovits. “Supersaturated graphs and hypergraphs”. In: *Combinatorica* 3.2 (1983), pp. 181–192. ISSN: 0209-9683. DOI: 10.1007/BF02579292. URL: <https://doi.org/10.1007/BF02579292>.
- [KMV15] Alexandr Kostochka et al. “Turán problems and shadows III: expansions of graphs”. In: *SIAM J. Discrete Math.* 29.2 (2015), pp. 868–876. ISSN: 0895-4801, 1095-7146. DOI: 10.1137/140977138. URL: <https://doi.org/10.1137/140977138>.

- [MM23] Dhruv Mubayi and Sayan Mukherjee. “Triangles in graphs without bipartite suspensions”. In: *Discrete Math.* 346.6 (2023), Paper No. 113355, 19. ISSN: 0012-365X, 1872-681X. DOI: 10.1016/j.disc.2023.113355. URL: <https://doi.org/10.1016/j.disc.2023.113355>.
- [MV16] Dhruv Mubayi and Jacques Verstraëte. “A survey of Turán problems for expansions”. In: *Recent trends in combinatorics*. Vol. 159. IMA Vol. Math. Appl. Springer, [Cham], 2016, pp. 117–143. ISBN: 978-3-319-24296-5; 978-3-319-24298-9. DOI: 10.1007/978-3-319-24298-9\\_5. URL: [https://doi.org/10.1007/978-3-319-24298-9\\_5](https://doi.org/10.1007/978-3-319-24298-9_5).
- [Tur41] Paul Turán. “Eine Extremalaufgabe aus der Graphentheorie”. In: *Mat. Fiz. Lapok* 48 (1941), pp. 436–452.