

# Double Turán Problem

Ray Tsai

May 9, 2025

# Overview

- 1 Preliminaries
- 2 Double Turán Problem
- 3 Induced Double Turán Problem
- 4 Main Results
- 5 Proof of Theorem B
- 6 Proof of Theorem D

What is the Turán problem?

What is the Turán problem?

## Question

Given a graph  $F$ , how many edges can an  $n$ -vertex graph have while containing no copy of  $F$  as a subgraph?

What is the Turán problem?

## Question

Given a graph  $F$ , how many edges can an  $n$ -vertex graph have while containing no copy of  $F$  as a subgraph?

We call the following quantity the **Turán number** or **extremal number** of  $F$ :

## Definition

$$\text{ex}(n, F) := \max\{e(G) : |V(G)| = n \text{ and } F \not\subseteq G\}$$

## Turán's theorem

The maximum number of edges in an  $n$ -vertex graph containing no clique of order  $r + 1$  is  $e(T_r(n))$ , with equality only for  $T_r(n)$ .

## Turán's theorem

The maximum number of edges in an  $n$ -vertex graph containing no clique of order  $r + 1$  is  $e(T_r(n))$ , with equality only for  $T_r(n)$ .




Figure:  $T_4(13)$

## Erdős-Stone Theorem, Simonovits' Theorem

Let  $F$  be any graph of chromatic number  $r + 1 \geq 3$ . Then  
 $\text{ex}(n, F) = (1 + o(1)) T_r(n)$  as  $n \rightarrow \infty$ .



# Double Turán Problem

What is the double Turán problem?

# Double Turán Problem

What is the double Turán problem?

Let  $G_1, G_2, \dots, G_m$  be graphs on  $[n]$  with  $E(F) \not\subseteq E(G_i) \cap E(G_j)$  for distinct  $i, j \in [m]$ .

# Double Turán Problem

What is the double Turán problem?

Let  $G_1, G_2, \dots, G_m$  be graphs on  $[n]$  with  $E(F) \not\subseteq E(G_i) \cap E(G_j)$  for distinct  $i, j \in [m]$ . (**Double  $F$ -free**)

# Double Turán Problem

What is the double Turán problem?

Let  $G_1, G_2, \dots, G_m$  be graphs on  $[n]$  with  $E(F) \not\subseteq E(G_i) \cap E(G_j)$  for distinct  $i, j \in [m]$ . (**Double  $F$ -free**)

## Question

What is the value of  $\phi(m, n, F) = \max \sum_{i=1}^m |G_i|$ ?

# Double Turán Problem

Double Turán problems are closely related to Turán problems for 3-uniform hypergraphs  $H$  through **link graphs**.

# Double Turán Problem

Double Turán problems are closely related to Turán problems for 3-uniform hypergraphs  $H$  through **link graphs**.

## Definition

For  $i \in V(H)$ , define graph  $H_i$  with

$$V(H_i) = V(H) \setminus \{i\} \quad \text{and} \quad E(H_i) = \{\{j, k\} : \{i, j, k\} \in E(H)\}.$$

# Double Turán Problem

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $O$

# Double Turán Problem

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $O$



# Double Turán Problem

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $O$

# Double Turán Problem

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $O$

# Double Turán Problem

Example: Octahedron-free 3-uniform hypergraph  $H$




Figure: Octahedron  $O$

$H$  is octahedron-free  $\implies H_1, H_2, \dots, H_n$  are double  $C_4$ -free.

$$\boxed{\text{ex}(n, O) \leq \phi(n, n, C_4)}$$

# Induced Double Turán Problem

What is the **induced** double Turán problem?

# Induced Double Turán Problem

What is the **induced** double Turán problem?

## Definition

We call graphs  $G_1, G_2, \dots, G_m$  **induced** if each  $G_i$  is an induced subgraph of  $\bigcup_{i=1}^m G_i$ .

# Induced Double Turán Problem

What is the **induced** double Turán problem?

## Definition

We call graphs  $G_1, G_2, \dots, G_m$  **induced** if each  $G_i$  is an induced subgraph of  $\bigcup_{i=1}^m G_i$ .

In other words,  $e \in E(G_i)$  iff  $e \in E(G_j)$  for all  $j = 1, \dots, m$ .

# Induced Double Turán Problem

What is the **induced** double Turán problem?

## Definition

We call graphs  $G_1, G_2, \dots, G_m$  **induced** if each  $G_i$  is an induced subgraph of  $\bigcup_{i=1}^m G_i$ .

In other words,  $e \in E(G_i)$  iff  $e \in E(G_j)$  for all  $j = 1, \dots, m$ .

Let graphs  $G_1, G_2, \dots, G_m$  be induced and double  $F$ -free.

## Question

What is the value of  $\phi^*(m, n, F) = \max \sum_{i=1}^m e(G_i)$ ?

# Induced Double Turán Problem

## Generalized Turán problem

What is the maximum number  $ex(n, F, K_3)$  of triangles in a graph  $H$  on  $[n]$  with no copy of  $F$  as a subgraph?



# Induced Double Turán Problem

## Generalized Turán problem

What is the maximum number  $\text{ex}(n, F, K_3)$  of triangles in a graph  $H$  on  $[n]$  with no copy of  $F$  as a subgraph?

For  $i \in V(G)$ , define  $G_i$  with

$$V(G_i) = V(G) \quad \text{and} \quad E(G_i) = \{\{j, k\} : \{i, j\}, \{j, k\}, \{i, k\} \in E(G)\}$$

# Induced Double Turán Problem

Ex. Octahedron-free graph  $G$ .




Figure: Octahedron




Figure: Octahedron Graph  $K_{2,2,2}$

# Induced Double Turán Problem

Ex. Octahedron-free graph  $G$ .




Figure: Octahedron




Figure: Octahedron Graph  $K_{2,2,2}$

# Induced Double Turán Problem

Ex. Octahedron-free graph  $G$ .




Figure: Octahedron




Figure: Octahedron Graph  $K_{2,2,2}$

# Induced Double Turán Problem

Ex. Octahedron-free graph  $G$ .




Figure: Octahedron




Figure: Octahedron Graph  $K_{2,2,2}$

$G$  is  $K_{2,2,2}$ -free  $\implies G_1, G_2, \dots, G_n$  are induced and double  $K_{2,2}$ -free.

$$\text{ex}(n, K_{2,2,2}, K_3) \leq \phi^*(n, n, K_{2,2})$$

## Theorem A

For  $m \geq 3$  and non-bipartite  $F$ , if  $n$  is large enough, then

$$\phi^*(m, n, F) = m \cdot \text{ex}(n, F),$$

with equality only for identical extremal  $n$ -vertex  $F$ -free graphs.

# Main Results

## Theorem A

For  $m \geq 3$  and non-bipartite  $F$ , if  $n$  is large enough, then

$$\phi^*(m, n, F) = m \cdot \text{ex}(n, F),$$

with equality only for identical extremal  $n$ -vertex  $F$ -free graphs.

## Theorem B

For  $m, n, r \geq 3$ ,

$$\phi^*(m, n, K_r) = m \cdot e(T_{r-1}(n)),$$

with equality for induced  $K_r$ -free graphs  $G_1, G_2, \dots, G_m$  only if  $G_1 = G_2 = \dots = G_m = T_{r-1}(n)$ .

# Main Results

The non-bipartite case for  $\phi(m, n, K_r)$  is not as simple.

Intuitively, we might guess

$$\phi(m, n, K_r) = \binom{n}{2} + (m-1)\text{ex}(n, K_r)$$



# Main Results

The non-bipartite case for  $\phi(m, n, K_r)$  is not as simple.

Intuitively, we might guess

$$\phi(m, n, K_r) = \binom{n}{2} + (m-1)\text{ex}(n, K_r)$$

# Main Results

We need the idea of  $(m, n, k)$ -blowup to state the result for  $\phi(m, n, K_r)$ .

# Main Results

We need the idea of  $(m, n, k)$ -blowup to state the result for  $\phi(m, n, K_r)$ .




Figure: Example of an  $(4, n, 5)$ -blowup not containing a double  $K_3$ .

# Main Results

Let  $f(m, n, r)$  denote the maximum possible sum of edges in an double  $K_r$ -free  $(m, n, k)$ -blowup with  $k < R_{\binom{m}{2}}(r)$  ( $\binom{m}{2}$ -color Ramsey number for  $K_r$ ).

# Main Results

Let  $f(m, n, r)$  denote the maximum possible sum of edges in an double  $K_r$ -free  $(m, n, k)$ -blowup with  $k < R_{\binom{m}{2}}(r)$  ( $\binom{m}{2}$ -color Ramsey number for  $K_r$ ).

## Theorem C

For  $n \geq 1$ ,

- ① if  $r \geq 2$  and  $m \geq 1$ ,

$$\phi(m, n, K_r) = f(m, n, r).$$

②

$$\phi(3, n, K_3) = \binom{n}{2} + \left\lfloor \frac{n^2}{2} \right\rfloor.$$

In particular,

$$\lim_{n \rightarrow \infty} \frac{\phi(4, n, K_3)}{\binom{n}{2} + 3 \left\lfloor \frac{n^2}{4} \right\rfloor} > 1$$

## Conjecture

Let  $F$  be any non-empty graph and  $m, n \geq 1$ . Then

$$\phi^*(m, n, F) = \Theta(m \cdot \text{ex}(n, F) + n^2).$$

## Conjecture

Let  $F$  be any non-empty graph and  $m, n \geq 1$ . Then

$$\phi^*(m, n, F) = \Theta(m \cdot \text{ex}(n, F) + n^2).$$

$$\phi^*(m, n, F) \geq \max\left\{\binom{n}{2}, m \cdot \text{ex}(n, F)\right\}.$$

## Conjecture

Let  $F$  be any non-empty graph and  $m, n \geq 1$ . Then

$$\phi^*(m, n, F) = \Theta(m \cdot \text{ex}(n, F) + n^2).$$

By Theorem 1, the conjecture is true when  $F$  is non-bipartite.



## Conjecture

Let  $F$  be any non-empty graph and  $m, n \geq 1$ . Then

$$\phi^*(m, n, F) = \Theta(m \cdot \text{ex}(n, F) + n^2).$$

Since

$$\text{ex}(n, K_{2,2,2}, K_3) \leq \phi^*(n, n, K_{2,2})$$

the conjecture implies

$$\text{ex}(n, K_{2,2,2}, K_3) \leq O(n^2)$$

which solves a conjecture of Mubayi and Verstraete.

## Theorem D

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

## Theorem D

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

If  $P$  is a path of length 2 and  $m = o(n^{1/4})$ ,

$$\binom{n}{2} + m - 1 \leq \phi^*(m, n, P) \leq \phi(m, n, P) = \binom{n}{2} + \binom{m}{2} \left\lfloor \frac{n}{2} \right\rfloor.$$

## Theorem E

Let  $P$  be a path with two edges. Then  $\phi(n, n, P) = \Omega(n^{5/2})$ , whereas  $\phi^*(n, n, P) = o(n^{5/2})$ , as  $n \rightarrow \infty$ . In particular,

$$\lim_{n \rightarrow \infty} \frac{\phi^*(n, n, P)}{\phi(n, n, P)} = 0.$$

## Theorem E

Let  $P$  be a path with two edges. Then  $\phi(n, n, P) = \Omega(n^{5/2})$ , whereas  $\phi^*(n, n, P) = o(n^{5/2})$ , as  $n \rightarrow \infty$ . In particular,

$$\lim_{n \rightarrow \infty} \frac{\phi^*(n, n, P)}{\phi(n, n, P)} = 0.$$

This shows that  $\phi(n, n, P)$  and  $\phi^*(n, n, P)$  are very different problems.

# Proof of Theorem B

## Theorem B

Let  $m, n, r \geq 3$ . Then

$$\phi^*(m, n, K_r) = m \cdot e(T_{r-1}(n)),$$

with equality for induced  $K_r$ -free graphs  $G_1, G_2, \dots, G_m$  only if  $G_1 = G_2 = \dots = G_m = T_{r-1}(n)$ .

## Proof Roadmap

- Step 1: Reduce to the case of smaller  $m$
- Step 2: Further reduce to an optimization problem
- Step 3: Solve the optimization problem

# Step 1: Reduce to the case of smaller $m$

## Proof Roadmap

- **Step 1:** Reduce to the case of smaller  $m$
- Step 2: Further reduce to an optimization problem
- Step 3: Solve the optimization problem


## Step 2: Further reduce to an optimization problem

### Proof Roadmap


- Step 1: Reduce to the case of smaller  $m$
- **Step 2:** Further reduce to an optimization problem
- Step 3: Solve the optimization problem



## Step 2: Further reduce to an optimization problem




## Step 2: Further reduce to an optimization problem



The intersection of  $G_1, G_2$  (gray area) is double  $K_r$ -free.

## Step 2: Further reduce to an optimization problem




The intersection of  $G_1, G_2$  (gray area) is double  $K_r$ -free.

### Observation

If  $G_1, G_2$  intersects in  $t$  vertices, Then

$$e(G_1) + e(G_2) \leq \binom{n-t}{2} + (n-t)t + 2\text{ex}(t, F)$$

## Step 2: Further reduce to an optimization problem



The intersection of  $G_1, G_2$  (gray area) is double  $K_r$ -free.

### Observation

If  $G_1, G_2$  intersects in  $t$  vertices, Then

$$e(G_1) + e(G_2) \leq \binom{n-t}{2} + (n-t)t + 2\text{ex}(t, F)$$

$\implies$  we are done if the unique maximum on  $t$  is at  $t = n$ .

# Step 3: Solve the optimization problem

## Proof Roadmap

- Step 1: Reduce to the case of smaller  $m$
- Step 2: Further reduce to an optimization problem
- **Step 3:** Solve the optimization problem

# Proof of Theorem D: Upper Bound

## Theorem D

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

# Proof of Theorem D: Upper Bound

## Theorem D

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

We need to show

$$\phi(m, n, F) \leq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

# Proof of Theorem D: Upper Bound

## Theorem D

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

We need to show

$$\phi(m, n, F) \leq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

Let  $E_S$  be the set of edges in exactly  $\{G_i\}_{i \in S}$ .



# Proof of Theorem D: Upper Bound

## Theorem D

Let  $F$  be a graph. If there exists an extremal  $F$ -free  $n$ -vertex graph with maximum degree at most  $\sqrt{n}/m^2$ , then

$$\phi(m, n, F) = \binom{n}{2} + \binom{m}{2} \text{ex}(n, F).$$

We need to show

$$\phi(m, n, F) \leq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

Let  $E_S$  be the set of edges in exactly  $\{G_i\}_{i \in S}$ .

$$\Rightarrow \sum_{i=1}^m e(G_i) = \sum_{S \subseteq [m]} |S| |E_S| \leq \binom{n}{2} + \sum_{S \subseteq [m], |S| \geq 2} (|S| - 1) |E_S|.$$

# Proof of Theorem D: Upper Bound

$$\sum_{\substack{S \subseteq [m] \\ |S| \geq 2}} (|S| - 1) |E_S| = \sum_{\substack{S \subseteq [m] \\ |S| = 2}} \sum_{T \supseteq S} \frac{(|T| - 1) |E_T|}{\binom{|T|}{2}} \leq \sum_{\substack{S \subseteq [m] \\ |S| = 2}} \sum_{T \supseteq S} |E_T|,$$

as each  $T \in [m]$  with  $|T| \geq 2$  is counted  $\binom{|T|}{2}$  times.

# Proof of Theorem D: Upper Bound

Observation: If  $|S| \geq 2$ , the edge set  $\bigcup_{T \supseteq S} E_T$  is  $F$ -free.

# Proof of Theorem D: Upper Bound

Observation: If  $|S| \geq 2$ , the edge set  $\bigcup_{T \supseteq S} E_T$  is  $F$ -free.

$$\left| \bigcup_{T \supseteq S} E_T \right| = \sum_{T \supseteq S} |E_T| \leq \text{ex}(n, F)$$

# Proof of Theorem D: Upper Bound

Observation: If  $|S| \geq 2$ , the edge set  $\bigcup_{T \supseteq S} E_T$  is  $F$ -free.

$$\left| \bigcup_{T \supseteq S} E_T \right| = \sum_{T \supseteq S} |E_T| \leq \text{ex}(n, F)$$

$$\Rightarrow \sum_{\substack{S \subseteq [m] \\ |S| \geq 2}} (|S| - 1) |E_S| \leq \sum_{\substack{S \subseteq [m], \\ |S|=2}} \sum_{T \supseteq S} |E_T| \leq \binom{m}{2} \text{ex}(n, F)$$

This proves the upper bound.

# Proof of Theorem D: Lower Bound

We need to show

$$\phi(m, n, F) \geq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

# Proof of Theorem D: Lower Bound

We need to show

$$\phi(m, n, F) \geq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

Let  $H_1, \dots, H_{\binom{m}{2}}$  be extremal  $F$ -free graphs on  $[n]$  with  $\Delta(H_i) \leq \sqrt{n}/m^2$

# Proof of Theorem D: Lower Bound

We need to show

$$\phi(m, n, F) \geq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

Let  $H_1, \dots, H_{\binom{m}{2}}$  be extremal  $F$ -free graphs on  $[n]$  with  $\Delta(H_i) \leq \sqrt{n}/m^2$

We are done if  $H_1, \dots, H_{\binom{m}{2}}$  can be embedded onto  $[n]$  s.t. edges are pairwise disjoint.



# Proof of Theorem D: Lower Bound

We need to show

$$\phi(m, n, F) \geq \binom{n}{2} + \text{ex}(n, F) \binom{m}{2}.$$

Let  $H_1, \dots, H_{\binom{m}{2}}$  be extremal  $F$ -free graphs on  $[n]$  with  $\Delta(H_i) \leq \sqrt{n}/m^2$

We are done if  $H_1, \dots, H_{\binom{m}{2}}$  can be embedded onto  $[n]$  s.t. edges are pairwise disjoint.

IDEA: start with any embedding and iteratively decrease overlapping edges

# Proof of Theorem D: Lower Bound

Define a  $(u, v, i)$ -swap by swapping the embedding of vertex  $u$  and  $v$  of  $H_i$

# Proof of Theorem D: Lower Bound

Define a  $(u, v, i)$ -swap by swapping the embedding of vertex  $u$  and  $v$  of  $H_i$

$\implies (u, v, i)$ -swap preserves graph isomorphism.

# Proof of Theorem D: Lower Bound

For vertex  $v$ , let  $N(v) = N_{H_1}(v) \cup \cdots \cup N_{H_{\binom{m}{2}}}(v)$

# Proof of Theorem D: Lower Bound

For vertex  $v$ , let  $N(v) = N_{H_1}(v) \cup \cdots \cup N_{H_{\binom{m}{2}}}(v)$

Suppose there exists  $\{u, w\} \in E(H_i) \cap E(H_j)$

# Proof of Theorem D: Lower Bound

For vertex  $v$ , let  $N(v) = N_{H_1}(v) \cup \dots \cup N_{H_{\binom{m}{2}}}(v)$

Suppose there exists  $\{u, w\} \in E(H_i) \cap E(H_j)$

$$|N(u)| \leq M \cdot \Delta \leq n^{1/2}/2$$

# Proof of Theorem D: Lower Bound

For vertex  $v$ , let  $N(v) = N_{H_1}(v) \cup \dots \cup N_{H_{\binom{m}{2}}}(v)$

Suppose there exists  $\{u, w\} \in E(H_i) \cap E(H_j)$

$$|N(u)| \leq M \cdot \Delta \leq n^{1/2}/2$$

$$\implies |N(u) \cup N(N(u))| \leq \Delta + \Delta(\Delta - 1) \leq n/4$$

# Proof of Theorem D: Lower Bound

For vertex  $v$ , let  $N(v) = N_{H_1}(v) \cup \cdots \cup N_{H_{\binom{m}{2}}}(v)$

Suppose there exists  $\{u, w\} \in E(H_i) \cap E(H_j)$

$$|N(u)| \leq M \cdot \Delta \leq n^{1/2}/2$$

$$\implies |N(u) \cup N(N(u))| \leq \Delta + \Delta(\Delta - 1) \leq n/4$$

$$\implies \exists v \notin N(u) \cup N(N(u)), \text{ so } N(u) \cap N(v) = \emptyset.$$



# Proof of Theorem D: Lower Bound

For vertex  $v$ , let  $N(v) = N_{H_1}(v) \cup \dots \cup N_{H_{\binom{m}{2}}}(v)$

Suppose there exists  $\{u, w\} \in E(H_i) \cap E(H_j)$

$$|N(u)| \leq M \cdot \Delta \leq n^{1/2}/2$$

$$\implies |N(u) \cup N(N(u))| \leq \Delta + \Delta(\Delta - 1) \leq n/4$$

$$\implies \exists v \notin N(u) \cup N(N(u)), \text{ so } N(u) \cap N(v) = \emptyset.$$

$$\implies (u, v, i)\text{-swap makes } \{u, w\} \text{ not overlapping!}$$

# Proof of Theorem D: Lower Bound

For vertex  $v$ , let  $N(v) = N_{H_1}(v) \cup \dots \cup N_{H_{\binom{m}{2}}}(v)$

Suppose there exists  $\{u, w\} \in E(H_i) \cap E(H_j)$

$$|N(u)| \leq M \cdot \Delta \leq n^{1/2}/2$$

$$\implies |N(u) \cup N(N(u))| \leq \Delta + \Delta(\Delta - 1) \leq n/4$$

$$\implies \exists v \notin N(u) \cup N(N(u)), \text{ so } N(u) \cap N(v) = \emptyset.$$

$$\implies (u, v, i)\text{-swap makes } \{u, w\} \text{ not overlapping!}$$

Keep swapping until no overlapping edges left