

Arithmetic Progression

General term : $a_n = a + (n - 1)d$

n^{th} term from end : $a + (m - n)d$

where n^{th} term from end = $(m - n + 1)$ from start

n^{th} term from end: $l - (n - 1)d$

if A,B,C are in AP: $2B = A + C$

$$a_1 + a_n = a_2 + a_{n-1}$$

$$2a_n = a_{n+k} + a_{n-k}$$

selecting middle term of AP:

$$\text{ODD} \Rightarrow -a + \quad \quad \text{CD} \rightarrow d$$

$$\text{EVEN} \Rightarrow (a - d)(a + d) \quad \text{CD} \rightarrow 2d$$

Sum of AP :

$$\text{with common difference } S_n = \frac{n}{2}\{2a + (n - 1)d\}$$

$$\text{with last term } S_n = \frac{n}{2}\{a + l\}$$

results of sum of AP :

• Seq is AP \rightarrow if sum of n terms form $An^2 + Bn$

• if ratio of sum is given, then ratio of n^{th} term is
 \rightarrow replace n by $2n - 1$

• if ratio of n^{th} term is given, then ratio of sum is
 \rightarrow replace n by $\frac{n+1}{2}$

Geometric Progression

General terms : $a_n = ar^{n-1}$

n^{th} term from end : $a_n = ar^{m-n}$

using last term : $a_n = l \left(\frac{1}{r}\right)^{n-1}$

Properties of GP :

• GP divided/multiplied by constant, stays GP

• reciprocal of GP, is GP

$$\bullet a_1 a_n = a_2 a_{n-1} = \dots = a_k a_{n-k+1}$$

$$\bullet b^2 = ac$$

$$\bullet \text{for a GP } a_1, a_2 \dots a_n \\ \Rightarrow GM = (a_1 a_2 \dots a_n)^{\frac{1}{n}}$$

• if $a_1, a_2 \dots a_n \Rightarrow$ GP
 $\log a_1, \log a_2 \dots \log a_n \Rightarrow$ AP and vice-versa

results of sum of GP :

$$S_n = \begin{cases} a \left(\frac{1 - r^n}{1 - r} \right), & r \neq 1 \\ na, & r \end{cases}$$

$$S_n = \frac{a - lr}{1 - r} = \frac{lr - a}{r - 1}$$

$$S_\infty = \begin{cases} \frac{a}{1 - r} & -1 < r < 1 \end{cases}$$

Harmonic Progression

if $a_1, a_2 \dots a_n$ are in HP

then $\frac{1}{a_1}, \frac{1}{a_2} \dots \frac{1}{a_n}$ are in AP

$$\text{common difference : } d = \frac{1}{a_2} - \frac{1}{a_1}$$

$$\text{general term : } a_n = \frac{1}{\frac{1}{a_1} + (n - 1)d}$$

Harmonic Mean(H)

$$\bullet H = \frac{2ab}{a + b}$$

• $a, H_1, H_2 \dots, H_n, b$ are in HP

then $\frac{1}{a}, \frac{1}{H_1}, \frac{1}{H_2} \dots, \frac{1}{H_n}, \frac{1}{b}$ are in AP

$$\rightarrow \frac{1}{b} = (n + 2)^{\text{th}}$$

$$\rightarrow \frac{1}{b} = \frac{1}{a} + (n + 1)D \quad D = \frac{a - b}{(n + 1)ab}$$

Arithmetico-geometric Sequence

$a, (a + d)r, (a + 2d)r^2, \dots$ is a A.G. sequence

$$n^{\text{th}} \text{ term : } a_n = \{a + (n - 1)d\} \cdot r^{n-1}$$

$$\text{sum of } \infty \text{ term : } S_\infty = \frac{a}{1 - r} + \frac{d \cdot r}{(1 - r)^2}$$

Sum of Some Sequence

$$\text{first } n \text{ natural no : } \frac{n(n + 1)}{2}$$

$$\text{first } n \text{ odd no : } n^2$$

$$\text{first } n \text{ even no : } n(n + 1)$$

$$\text{square of first } n \text{ no : } \frac{n(n + 1)(2n + 1)}{6}$$

$$\text{cube of first } n \text{ no : } \left\{ \frac{n(n + 1)}{2} \right\}^2$$

$$4^{\text{th}} \text{ power of } n \text{ no : } \frac{n(n + 1)(2n + 1)(3n^2 + 3n - 1)}{30}$$

Relation between AM, GM and HM **Tips and Tricks**

(black space for tips,tricks and important question)

- **A,G and H between 2 numbers(a and b):**

$$A = \frac{a+b}{2} \quad G = \sqrt{AB} \quad H = \frac{2ab}{a+b}$$

- $A > G > H$
- quadratic eq having a and b as its roots

$$x^2 - 2Ax + G^2 = 0$$

- the two numbers(a,b) are $A \pm \sqrt{A^2 - G^2}$
- if A and G are in the ratio $m : n$, then the number(a,b) are in ratio

$$m + \sqrt{m^2 - n^2} : m - \sqrt{m^2 - n^2}$$

- **A,G and H between 3 numbers(a, b, c):**

$$A = \frac{a+b+c}{3} \quad G = (abc)^{\frac{1}{3}} \quad \frac{1}{H} = \frac{1}{3} \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right)$$

- cubic equation where a,b,c are the roots

$$x^3 - 3Ax^2 + \frac{3G^3}{H}x - G^3 = 0$$