PRACTICE SHEET

1. In a manufacture of readymade garments, which average is used to find the most frequent size?
(a) Arithmetic mean (b) Geometric mean
(c) Mode (d) Harmonic mean

2. What is the arithmetic mean of the series ${}^nC_0, {}^nC_1, \ldots {}^nC_n$, ?

(a) $\frac{2^n}{n}$ (b) $\frac{2^n}{(n+1)}$

(c) $\frac{2^{(n+1)}}{n}$ (d) $\frac{2^{(n+1)}}{(n+1)}$

3. If from the point of intersection of two gives, a perpendicular is drawn on the x – axis, what does the x – coordinate give?
(a) Arithmetic Mean (b) Mode
(c) Median (d) Geometric Mean

4. The definition of Mode fails if:
(a)The maximum frequency is repeated
(b)The maximum frequency is not repeated
(c)The maximum frequency occurs in the middle
(d)The curve drawn with the help of given data is symmetrical

5. Which one of the following statement is not correct?
(a)Median divides distributions into two equal subgrops
(b)The third quartile is the same as the 75th percentile
(c)The 5th decile is the same as the 50th percentile
(d)The 50th decile is the same as the 5th percentile

Town	Population
A	6863
B	519
C	12185
D	1785

What is the most appropriate diagram to present the above data?
(a) Pie diagram (b) Bar chart
(c) Cubic chart (d) Histogram

11. Consider the two series observations A and B as follows:

Series A	1019	1008	1015	1006	1002
Series B	1.9	0.8	1.5	0.6	0.2

If the standard deviation of the Series A is $\sqrt{38}$, then what is the standard deviation of the Series B?
(a) 3.8 (b) $\sqrt{0.38}$
(c) 0.38 (d) $\sqrt{38}$

12. Let $\bar{X}$ be the mean of n observations $x_1, x_2, \ldots x_n$. If (a – b) is added to each observation, then what is the mean of new set of observations?
(a) 0 (b) $\bar{X}$
(c) $\bar{X} - (a - b)$ (d) $\bar{X} + (a - b)$

13. What is the values of *n* for which the numbers1, 2, 3, ..., n have variance 2?