

# MATH 8510 GALOIS THEORY

LU JUNYU

## 1. WEEK 1

Let's agree on some facts and conventions from elementary abstract algebra, in particular those with polynomial rings before we dig into Galois theory.

A ring is always commutative with multiplicative identity 1 unless otherwise stated.  $R^*$  is the multiplicative group of units in  $R$  and  $R^\times = R \setminus \{0\}$ . We can use these two notations interchangeably when  $R$  is a field.

Let  $F$  be a field. A polynomial ring  $F[X]$  with an indeterminate  $X$  is an  $F$ -vector space with basis  $1, X, X^2, \dots, X^n, \dots$  with the multiplication

$$\left(\sum_i a_i X^i\right)\left(\sum_j b_j X^j\right) = \sum_k \left(\sum_{i+j=k} a_i b_j\right) X_k,$$

where  $X^0$  is defined to be 1. Alternatively, we can identify  $R[X]$  with

$$R^{(\mathbb{N})} = \{(a_i) : a_i \in R, a_i = 0 \text{ for all but finitely many } i \in \mathbb{N}\}$$

in an obvious way. But usually, we want to say  $R$  embeds into  $R[X]$  although the most formal way is to identify  $R$  with a subring of  $R[X]$ . The degree function has the following properties:

- (1)  $\deg(f + g) \leq \max(\deg f, \deg g)$ ,
- (2)  $\deg(fg) = \deg f + \deg g$ .

There are plenty results by arguing over the degree of a polynomial. We have  $(R[X])^* = R^*$  if  $R$  is an integral domain. We have the division algorithm on  $R[X]$ .

**Theorem 1.1.** *Let  $F$  be a commutative ring. Then  $F[X]$  is a PID if and only if  $F$  is a field.*

Hence or otherwise,  $\mathbb{Z}[X]$  is not a PID. Indeed,  $\langle 2, X \rangle$  is an example of an ideal that cannot be generated by a single polynomial.  $K[X, Y]$  is not a PID as  $\langle X, Y \rangle$  is not principal.

**Theorem 1.2.** *An ideal in a PID is prime if and only if it is maximal.*

**Definition 1.3.** If  $f(X) \in F[X]$  where  $F$  is a field, then a **root** of  $f$  in  $F$  is an element  $\alpha \in F$  such that  $f(\alpha) = 0$ .

Given a polynomial  $f[X] \in F[X]$  and any  $u \in F$ , the division algorithm give us:

$$f(X) = q(X)(X - u) + f(u).$$

And lying in the center of proving that every finite subgroup of  $F^\times$  is cyclic is the following theorem.

**Theorem 1.4.** *Let  $F$  be a field and  $f[X] \in F[X]$  a polynomial of degree  $n$ . Then  $f$  has at most  $n$  roots.*

**Definition 1.5.** Let  $F$  be a field. A nonzero polynomial  $p(X) \in F[X]$  is said to be **irreducible** over  $F$  (or **irreducible** in  $F[X]$ ) if  $\deg p \geq 1$  and there is no factorization  $p = fg$  in  $F[X]$  with  $\deg f < \deg p$  and  $\deg g < \deg p$ .

A quadratic or cubic polynomial is irreducible in  $F[X]$  if and only if it has no root in  $F$ .

**Theorem 1.6** (Gauss's Lemma). *A polynomial  $f(X) \in \mathbb{Z}[X]$  is irreducible if and only if it is irreducible over  $\mathbb{Q}[X]$ .*

**Theorem 1.7** (Eisenstein Criterion). *Let  $f(X) = a_0 + a_1X + \dots + a_nX^n \in \mathbb{Z}[X]$  be a polynomial over integers with  $a_n \neq 0$ . Suppose that there exists a prime  $p$  such that*

- (1)  $p \nmid a_n$ ,
- (2)  $p \mid a_i$  for  $i = 0, 1, \dots, n-1$ ,
- (3)  $p^2 \nmid a_0$ .

*Then  $f(X)$  is irreducible over  $\mathbb{Z}[X]$ .*

**Theorem 1.8.** *Let  $F$  be a field. Then  $(f(X))$  is a prime ideal in  $F[X]$  if and only if  $f(X)$  is irreducible. Equivalently,  $f$  is irreducible if and only if  $K[X]/(f)$  is a field.*

Making use of above results, we finally reach the very last theorem which functions as a cornerstone in many arguments.

**Theorem 1.9.** *Let  $k$  be a field and  $f[X]$  a monic irreducible polynomial in  $k[X]$  of degree  $d$ . Let  $K = k[X]/I$ , where  $I = (f)$ , and  $\beta = X + I \in K$ . Then:*

- (1)  $K$  is a field and  $k' = \{a + I : a \in k\}$  is a subfield of  $K$  isomorphic to  $k$ ,
- (2)  $\beta$  is a root of  $g$  in  $K$ ,
- (3) if  $g(X) \in k[X]$  and  $\beta$  is a root of  $g$  in  $K$ , then  $f \mid g$  in  $k[X]$ ,
- (4)  $f$  is the unique monic irreducible polynomial in  $k[X]$  having  $\beta$  as a root,
- (5)  $1, \beta, \beta^2, \dots, \beta^{d-1}$  forms a basis of  $K$  as a vector space over  $k$  and so  $\dim_k(K) = d$ .

.....