

Assignment 1

Wendy Nieuwkamer

16-9-2016

1 Question 1

Suppose that we have historical data of result of soccer matches of teams playing against Ajax. We want to use this information to learn to predict at a certain moment whether a team will win, lose or draw against Ajax. Our approach will be based on Machine Learning.

1.1 Define the given and the goal of the prediction task. Classify the learning task.

Given are the results of soccer matches, the goal is to decide whether a team will win, lose or draw when playing. The learning task is supervised, as we are given inputs with their wanted result. It is a classification problem as every input should be classified as one of three possibilities.

1.2 What would be the form of training data for the learning task? Give a small training set.

An example of a training set could be the following table:

team	results
1	win
2	draw
3	lose
1	lose
3	draw
4	win

Table 1: training examples

2 Question 2

Use the training data in the table to answer all the subquestions.

x	y
3	6
5	7
6	10

2.1 Manually calculate two iterations of the gradient descent algorithm for univariate linear regression function.

Initialize the parameters such that the regression function passes through the origin $(0, 0)$ and has an angle of 45 degrees. Use a learning rate of 0.1. Give the intermediate results of your calculations and also compute the mean-squared error of the function after 2 iterations.

Gradient Descent Algorithm

Repeat until convergence {

$$\begin{aligned}\theta_0 &:= \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \\ \theta_1 &:= \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x^{(i)}\end{aligned}$$

}

The hypothesis h_{θ} is:

$$h_{\theta}(x^{(i)}) = \theta_0 + \theta_1 x^{(i)} \quad (1)$$

For this exercise the learning rate $\alpha = 0.1$ and the amount of learning examples $m = 3$ are given. The line that passes through the origin and has an angle of 45 degrees relative to the x-axis is $f(x) = x$, thus our starting values are $\theta_0 = 0$ and $\theta_1 = 1$.

First Iteration First, calculate the new values for θ_0 and θ_1 .

$$\begin{aligned}\theta_0 &= 0 - 0.1 \cdot \frac{1}{3} \sum_{i=1}^3 (x^{(i)} - y^{(i)}) \\ &= -\frac{1}{30}(-3 + -2 + -4) \\ &= 0.3\end{aligned}$$

$$\begin{aligned}\theta_1 &= 1 - 0.1 \cdot \frac{1}{3} \sum_{i=1}^3 (x^{(i)} - y^{(i)})x^{(i)} \\ &= 1 - \frac{1}{30}(-9 + -10 + -24) \\ &= 1 + \frac{43}{30} = \frac{73}{30} = 2.34\end{aligned}$$

Then, update θ_0 and θ_1 .

$$\begin{aligned}\theta_0 &:= 0.3 \\ \theta_1 &:= 2.34\end{aligned}$$

Second Iteration The values for m and α are still the same, but now we use the new values for θ_0 and θ_1 . Again, first the calculations.

$$\begin{aligned}\theta_0 &= 0.30 - 0.1 \cdot \frac{1}{3} \sum_{i=1}^3 (0.30 + 2.34x^{(i)} - y^{(i)}) \\ &= 0.30 - \frac{1}{30}(1.6 + 5.47 + 4.9) \\ &= 0.30 - 0.40 = -0.10\end{aligned}$$

$$\begin{aligned}\theta_1 &= 2.34 - 0.1 \cdot \frac{1}{3} \sum_{i=1}^3 (0.30 + 2.34x^{(i)} - y^{(i)})x^{(i)} \\ &= 2.34 - \frac{1}{30}(8.15 + 27.33 + 29.40) \\ &= 2.34 - 2.16 = 0.18\end{aligned}$$

Then, update θ_0 and θ_1 .

$$\begin{aligned}\theta_0 &:= -0.10 \\ \theta_1 &:= 0.18\end{aligned}$$

Mean-Squared Error Now we can calculate the mean-squared error to see how good our hypothesis $h_\theta(x^{(i)}) = -0.10 + 0.18x^{(i)}$ is. The formula we use is:

$$J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)})^2$$

This function is also called the cost function. We can now insert our values $m = 3$, $\theta_0 = -0.10$ and $\theta_1 = 0.18$.

$$\begin{aligned}J(\theta_0, \theta_1) &= \frac{1}{6} \sum_{i=1}^3 (0.18x^{(i)} - 0.10 - y^{(i)})^2 \\ &= \frac{1}{6} (30.91 + 38.44 + 81.36) \\ &= 25.12\end{aligned}$$

Thus, the mean-squared error for $m = 3$, $\theta_0 = -0.10$, $\theta_1 = 0.18$ and the given training examples is 25.12.

2.2 Convert the data to z-scores, repeat the calculations above and compare the results.

To calculate the z-scores use mean $\mu = 0$ and standard deviation $\sigma = 1$. The formula for calculating a z-score is:

$$Z = \frac{X - \mu}{\sigma}$$

Where X is the input value of the training example. Inserting our μ and σ into this formula gives:

$$Z = \frac{X - 0}{1} = X$$

Redoing the calculations would not be very useful as with this μ and σ the Z-score would be the same as the original input. It would be more useful to choose $\mu = 4.5$ and $\sigma = 1.125$ for x as this actually changes the values.

3 Question 3