

# Deformations of space curves lying on a del Pezzo surface

Hirokazu Nasu

Tokai University

June 12, 2025  
2025 Algebraic Geometry Conference in Jeonju Hanok Village



Today's slide

## Photo 1 (at Prof. Miyazaki's 65, 2025)



## Photo 2 (at Kumamoto, 2025)



## Photo 3 (at a workshop on IBS, 2024)



# Plan of Talk

- ➊ Hilbert schemes and deformation of flags
- ➋ Kleppe-Ellia conjecture and its generalization (Main result)  
(cf. [arXiv:2501.15788](https://arxiv.org/abs/2501.15788))
- ➌ Applications and Examples

## Plan of Talk

- ➊ Hilbert schemes and deformation of flags
- ➋ Kleppe-Ellia conjecture and its generalization (Main result)  
(cf. [arXiv:2501.15788](https://arxiv.org/abs/2501.15788))
- ➌ Applications and Examples

## Plan of Talk

- ➊ Hilbert schemes and deformation of flags
- ➋ Kleppe-Ellia conjecture and its generalization (Main result)  
(cf. [arXiv:2501.15788](https://arxiv.org/abs/2501.15788))
- ➌ Applications and Examples

# §1 Hilbert schemes and deformation of flags

# the Hilbert scheme

Given a projective scheme  $X$ , and given Hilbert polynomial  $P$ ,

$$\mathbf{Hilb}_P X = \{C \subset X \mid \text{closed subscheme of } P(C) = P\}$$

is called the **Hilbert scheme** of  $X$ . The Hilbert scheme has the following nice properties:

- fine moduli scheme, i.e. it has a **universal family**  $\mathcal{C} \subset X \times \mathbf{Hilb}_P X$  such that every deformation of  $C$  in  $X$  derived from  $\mathcal{C}$
- **projective** ( $\mathbf{Hilb}_P X \hookrightarrow \mathbf{Gr}$ )
- existence of nice **deformation theories**, e.g., if  $C$  is a loc. c.i., then  $H^0(N_{C/X})$  and  $H^1(N_{C/X})$  resp. represent the tangent and obstruction spaces at  $[C]$ .

while it also has “not so nice” properties: e.g.

- it may have **bad singularities** (e.g. non-reduced components),
- may be **highly reducible** (for some  $P$ ).

# the Hilbert scheme

Given a projective scheme  $X$ , and given Hilbert polynomial  $P$ ,

$$\mathbf{Hilb}_P X = \{C \subset X \mid \text{closed subscheme of } P(C) = P\}$$

is called the **Hilbert scheme** of  $X$ . The Hilbert scheme has the following nice properties:

- fine moduli scheme, i.e. it has a **universal family**  $\mathcal{C} \subset X \times \mathbf{Hilb}_P X$  such that every deformation of  $C$  in  $X$  derived from  $\mathcal{C}$
- **projective** ( $\mathbf{Hilb}_P X \hookrightarrow \mathbf{Gr}$ )
- existence of nice **deformation theories**, e.g., if  $C$  is a loc. c.i., then  $H^0(N_{C/X})$  and  $H^1(N_{C/X})$  resp. represent the tangent and obstruction spaces at  $[C]$ .

while it also has “not so nice” properties: e.g.

- it may have **bad singularities** (e.g. non-reduced components),
- may be **highly reducible** (for some  $P$ ).

# the Hilbert scheme

Given a projective scheme  $X$ , and given Hilbert polynomial  $P$ ,

$$\mathbf{Hilb}_P X = \{C \subset X \mid \text{closed subscheme of } P(C) = P\}$$

is called the **Hilbert scheme** of  $X$ . The Hilbert scheme has the following nice properties:

- fine moduli scheme, i.e. it has a **universal family**  $\mathcal{C} \subset X \times \mathbf{Hilb}_P X$  such that every deformation of  $C$  in  $X$  derived from  $\mathcal{C}$
- **projective** ( $\mathbf{Hilb}_P X \hookrightarrow \mathbf{Gr}$ )
- existence of nice **deformation theories**, e.g., if  $C$  is a loc. c.i., then  $H^0(N_{C/X})$  and  $H^1(N_{C/X})$  resp. represent the tangent and obstruction spaces at  $[C]$ .

while it also has “not so nice” properties: e.g.

- it may have **bad singularities** (e.g. non-reduced components),
- may be **highly reducible** (for some  $P$ ).

# the Hilbert scheme

Given a projective scheme  $X$ , and given Hilbert polynomial  $P$ ,

$$\mathbf{Hilb}_P X = \{C \subset X \mid \text{closed subscheme of } P(C) = P\}$$

is called the **Hilbert scheme** of  $X$ . The Hilbert scheme has the following nice properties:

- fine moduli scheme, i.e. it has a **universal family**  $\mathcal{C} \subset X \times \mathbf{Hilb}_P X$  such that every deformation of  $C$  in  $X$  derived from  $\mathcal{C}$
- **projective** ( $\mathbf{Hilb}_P X \hookrightarrow \mathbf{Gr}$ )
- existence of nice **deformation theories**, e.g., if  $C$  is a loc. c.i., then  $H^0(N_{C/X})$  and  $H^1(N_{C/X})$  resp. represent the tangent and obstruction spaces at  $[C]$ .

while it also has “not so nice” properties: e.g.

- it may have **bad singularities** (e.g. non-reduced components),
- may be **highly reducible** (for some  $P$ ).

# the Hilbert scheme

Given a projective scheme  $X$ , and given Hilbert polynomial  $P$ ,

$$\mathbf{Hilb}_P X = \{C \subset X \mid \text{closed subscheme of } P(C) = P\}$$

is called the **Hilbert scheme** of  $X$ . The Hilbert scheme has the following nice properties:

- fine moduli scheme, i.e. it has a **universal family**  $\mathcal{C} \subset X \times \mathbf{Hilb}_P X$  such that every deformation of  $C$  in  $X$  derived from  $\mathcal{C}$
- **projective** ( $\mathbf{Hilb}_P X \hookrightarrow \mathbf{Gr}$ )
- existence of nice **deformation theories**, e.g., if  $C$  is a loc. c.i., then  $H^0(N_{C/X})$  and  $H^1(N_{C/X})$  resp. represent the tangent and obstruction spaces at  $[C]$ .

while it also has “not so nice” properties: e.g.

- it may have **bad singularities** (e.g. non-reduced components),
- may be **highly reducible** (for some  $P$ ).

# the Hilbert scheme

Given a projective scheme  $X$ , and given Hilbert polynomial  $P$ ,

$$\mathbf{Hilb}_P X = \{C \subset X \mid \text{closed subscheme of } P(C) = P\}$$

is called the **Hilbert scheme** of  $X$ . The Hilbert scheme has the following nice properties:

- fine moduli scheme, i.e. it has a **universal family**  $\mathcal{C} \subset X \times \mathbf{Hilb}_P X$  such that every deformation of  $C$  in  $X$  derived from  $\mathcal{C}$
- **projective** ( $\mathbf{Hilb}_P X \hookrightarrow \mathbf{Gr}$ )
- existence of nice **deformation theories**, e.g., if  $C$  is a loc. c.i., then  $H^0(N_{C/X})$  and  $H^1(N_{C/X})$  resp. represent the tangent and obstruction spaces at  $[C]$ .

while it also has “not so nice” properties: e.g.

- it may have **bad singularities** (e.g. non-reduced components),
- may be **highly reducible** (for some  $P$ ).

# the Hilbert scheme

Given a projective scheme  $X$ , and given Hilbert polynomial  $P$ ,

$$\mathbf{Hilb}_P X = \{C \subset X \mid \text{closed subscheme of } P(C) = P\}$$

is called the **Hilbert scheme** of  $X$ . The Hilbert scheme has the following nice properties:

- fine moduli scheme, i.e. it has a **universal family**  $\mathcal{C} \subset X \times \mathbf{Hilb}_P X$  such that every deformation of  $C$  in  $X$  derived from  $\mathcal{C}$
- **projective** ( $\mathbf{Hilb}_P X \hookrightarrow \mathbf{Gr}$ )
- existence of nice **deformation theories**, e.g., if  $C$  is a loc. c.i., then  $H^0(N_{C/X})$  and  $H^1(N_{C/X})$  resp. represent the tangent and obstruction spaces at  $[C]$ .

while it also has “not so nice” properties: e.g.

- it may have **bad singularities** (e.g. non-reduced components),
- may be **highly reducible** (for some  $P$ ).

# the Hilbert scheme

Given a projective scheme  $X$ , and given Hilbert polynomial  $P$ ,

$$\mathbf{Hilb}_P X = \{C \subset X \mid \text{closed subscheme of } P(C) = P\}$$

is called the **Hilbert scheme** of  $X$ . The Hilbert scheme has the following nice properties:

- fine moduli scheme, i.e. it has a **universal family**  $\mathcal{C} \subset X \times \mathbf{Hilb}_P X$  such that every deformation of  $C$  in  $X$  derived from  $\mathcal{C}$
- **projective** ( $\mathbf{Hilb}_P X \hookrightarrow \mathbf{Gr}$ )
- existence of nice **deformation theories**, e.g., if  $C$  is a loc. c.i., then  $H^0(N_{C/X})$  and  $H^1(N_{C/X})$  resp. represent the tangent and obstruction spaces at  $[C]$ .

while it also has “not so nice” properties: e.g.

- it may have **bad singularities** (e.g. non-reduced components),
- may be **highly reducible** (for some  $P$ ).

## Dimension of $\text{Hilb } X$

For simplicity, we assume  $C$  is a c.i. in  $X$ . Then

$$\underbrace{h^0(C, N_{C/X}) - h^1(C, N_{C/X})}_{\text{exp.dim.} (= \chi(N_{C/X}) \text{ if } C \text{ is a curve})} \leq \dim_{[C]} \text{Hilb } X \leq \underbrace{h^0(C, N_{C/X})}_{\text{tangential dimension}}.$$

However, when  $H^1(N_{C/X}) \neq 0$ , it is hard to determine the dimension of  $\text{Hilb } X$  at  $[C]$  (depending on whether  $C$  is obstructed or not).

To resolve this problem, we take an intermediate variety  $C \subset S \subset X$  and use [Hilbert-flag scheme](#)

$$\text{HF } X = \{(C, S) \mid C \subset S \subset X\} \subset \text{Hilb } X \times \text{Hilb } X.$$

Let

$$N_{(C,S)} := N_{C/X} \times_{N_{S/X}|_C} N_{S/X}$$

be the normal sheaf of  $(C, S)$  in  $X$ .

## Dimension of $\text{Hilb } X$

For simplicity, we assume  $C$  is a c.i. in  $X$ . Then

$$\underbrace{h^0(C, N_{C/X}) - h^1(C, N_{C/X})}_{\text{exp.dim.} (= \chi(N_{C/X}) \text{ if } C \text{ is a curve})} \leq \dim_{[C]} \text{Hilb } X \leq \underbrace{h^0(C, N_{C/X})}_{\text{tangential dimension}} .$$

However, when  $H^1(N_{C/X}) \neq 0$ , it is hard to determine the dimension of  $\text{Hilb } X$  at  $[C]$  (depending on whether  $C$  is obstructed or not).

To resolve this problem, we take an intermediate variety  $C \subset S \subset X$  and use [Hilbert-flag scheme](#)

$$\text{HF } X = \{(C, S) \mid C \subset S \subset X\} \subset \text{Hilb } X \times \text{Hilb } X.$$

Let

$$N_{(C,S)} := N_{C/X} \times_{N_{S/X}|_C} N_{S/X}$$

be the normal sheaf of  $(C, S)$  in  $X$ .

## Dimension of $\text{Hilb } X$

For simplicity, we assume  $C$  is a c.i. in  $X$ . Then

$$\underbrace{h^0(C, N_{C/X}) - h^1(C, N_{C/X})}_{\text{exp.dim.} (= \chi(N_{C/X}) \text{ if } C \text{ is a curve})} \leq \dim_{[C]} \text{Hilb } X \leq \underbrace{h^0(C, N_{C/X})}_{\text{tangential dimension}} .$$

However, when  $H^1(N_{C/X}) \neq 0$ , it is hard to determine the dimension of  $\text{Hilb } X$  at  $[C]$  (depending on whether  $C$  is obstructed or not).

To resolve this problem, we take an intermediate variety  $C \subset S \subset X$  and use [Hilbert-flag scheme](#)

$$\text{HF } X = \{(C, S) \mid C \subset S \subset X\} \subset \text{Hilb } X \times \text{Hilb } X.$$

Let

$$N_{(C,S)} := N_{C/X} \times_{N_{S/X}|_C} N_{S/X}$$

be the normal sheaf of  $(C, S)$  in  $X$ .

## Dimension of $\text{Hilb } X$

For simplicity, we assume  $C$  is a c.i. in  $X$ . Then

$$\underbrace{h^0(C, N_{C/X}) - h^1(C, N_{C/X})}_{\text{exp.dim.} (= \chi(N_{C/X}) \text{ if } C \text{ is a curve})} \leq \dim_{[C]} \text{Hilb } X \leq \underbrace{h^0(C, N_{C/X})}_{\text{tangential dimension}} .$$

However, when  $H^1(N_{C/X}) \neq 0$ , it is hard to determine the dimension of  $\text{Hilb } X$  at  $[C]$  (depending on whether  $C$  is obstructed or not).

To resolve this problem, we take an intermediate variety  $C \subset S \subset X$  and use [Hilbert-flag scheme](#)

$$\text{HF } X = \{(C, S) \mid C \subset S \subset X\} \subset \text{Hilb } X \times \text{Hilb } X.$$

Let

$$N_{(C,S)} := N_{C/X} \times_{N_{S/X}|_C} N_{S/X}$$

be the normal sheaf of  $(C, S)$  in  $X$ .

## Dimension of $\text{Hilb } X$

For simplicity, we assume  $C$  is a c.i. in  $X$ . Then

$$\underbrace{h^0(C, N_{C/X}) - h^1(C, N_{C/X})}_{\text{exp.dim.} (= \chi(N_{C/X}) \text{ if } C \text{ is a curve})} \leq \dim_{[C]} \text{Hilb } X \leq \underbrace{h^0(C, N_{C/X})}_{\text{tangential dimension}} .$$

However, when  $H^1(N_{C/X}) \neq 0$ , it is hard to determine the dimension of  $\text{Hilb } X$  at  $[C]$  (depending on whether  $C$  is obstructed or not).

To resolve this problem, we take an intermediate variety  $C \subset S \subset X$  and use [Hilbert-flag scheme](#)

$$\text{HF } X = \{(C, S) \mid C \subset S \subset X\} \subset \text{Hilb } X \times \text{Hilb } X.$$

Let

$$N_{(C,S)} := N_{C/X} \times_{N_{S/X}|_C} N_{S/X}$$

be the normal sheaf of  $(C, S)$  in  $X$ .

## Dimension of $\text{Hilb } X$

For simplicity, we assume  $C$  is a c.i. in  $X$ . Then

$$\underbrace{h^0(C, N_{C/X}) - h^1(C, N_{C/X})}_{\text{exp.dim.} (= \chi(N_{C/X}) \text{ if } C \text{ is a curve})} \leq \dim_{[C]} \text{Hilb } X \leq \underbrace{h^0(C, N_{C/X})}_{\text{tangential dimension}} .$$

However, when  $H^1(N_{C/X}) \neq 0$ , it is hard to determine the dimension of  $\text{Hilb } X$  at  $[C]$  (depending on whether  $C$  is obstructed or not).

To resolve this problem, we take an intermediate variety  $C \subset S \subset X$  and use [Hilbert-flag scheme](#)

$$\text{HF } X = \{(C, S) \mid C \subset S \subset X\} \subset \text{Hilb } X \times \text{Hilb } X.$$

Let

$$N_{(C,S)} := N_{C/X} \times_{N_{S/X}|_C} N_{S/X}$$

be the normal sheaf of  $(C, S)$  in  $X$ .

## Naive question

Then

$$H^1(N_{(C,S)/X}) = 0 \implies \text{HF } X \text{ is nonsingular at } (C, S)$$

and if moreover  $H^i(N_{(C,S)/X}) = 0$  for all  $i > 0$ , then  $\text{HF } X$  is of expected dimension

$$\chi(N_{(C,S)/X}) = \chi(N_{C/S}) + \chi(N_{S/X}).$$

Let  $\mathcal{W}_{C,S}$  be an irreducible component passing through  $(C, S)$ , and  $pr_1 : \text{HF } X \rightarrow \text{Hilb } X$ ,  $(C', S') \mapsto [C']$ , the 1st projection.

### Question 1

When is the image  $pr_1(\mathcal{W}_{C,S})$  of  $\mathcal{W}_{C,S}$  an irreducible component of  $\text{Hilb } X$ ?

## Naive question

Then

$$H^1(N_{(C,S)/X}) = \mathbf{0} \implies \text{HF } X \text{ is nonsingular at } (C, S)$$

and if moreover  $H^i(N_{(C,S)/X}) = \mathbf{0}$  for all  $i > 0$ , then  $\text{HF } X$  is of expected dimension

$$\chi(N_{(C,S)/X}) = \chi(N_{C/S}) + \chi(N_{S/X}).$$

Let  $\mathcal{W}_{C,S}$  be an irreducible component passing through  $(C, S)$ , and  $pr_1 : \text{HF } X \rightarrow \text{Hilb } X$ ,  $(C', S') \mapsto [C']$ , the 1st projection.

### Question 1

When is the image  $pr_1(\mathcal{W}_{C,S})$  of  $\mathcal{W}_{C,S}$  an irreducible component of  $\text{Hilb } X$ ?

## Naive question

Then

$$H^1(N_{(C,S)/X}) = 0 \implies \text{HF } X \text{ is nonsingular at } (C, S)$$

and if moreover  $H^i(N_{(C,S)/X}) = 0$  for all  $i > 0$ , then  $\text{HF } X$  is of expected dimension

$$\chi(N_{(C,S)/X}) = \chi(N_{C/S}) + \chi(N_{S/X}).$$

Let  $\mathcal{W}_{C,S}$  be an irreducible component passing through  $(C, S)$ , and  $pr_1 : \text{HF } X \rightarrow \text{Hilb } X$ ,  $(C', S') \mapsto [C']$ , the 1st projection.

### Question 1

When is the image  $pr_1(\mathcal{W}_{C,S})$  of  $\mathcal{W}_{C,S}$  an irreducible component of  $\text{Hilb } X$ ?

## Naive question

Then

$$H^1(N_{(C,S)/X}) = \mathbf{0} \implies \mathbf{HF} X \text{ is nonsingular at } (C, S)$$

and if moreover  $H^i(N_{(C,S)/X}) = \mathbf{0}$  for all  $i > 0$ , then  $\mathbf{HF} X$  is of expected dimension

$$\chi(N_{(C,S)/X}) = \chi(N_{C/S}) + \chi(N_{S/X}).$$

Let  $\mathcal{W}_{C,S}$  be an irreducible component passing through  $(C, S)$ , and  $pr_1 : \mathbf{HF} X \rightarrow \mathbf{Hilb} X$ ,  $(C', S') \mapsto [C']$ , the 1st projection.

### Question 1

When is the image  $pr_1(\mathcal{W}_{C,S})$  of  $\mathcal{W}_{C,S}$  an irreducible component of  $\mathbf{Hilb} X$ ?

## Stable degeneration

Let  $C \subset S \subset X$  a sequence of closed subvarieties with  $H^i(N_{(C,S)/X}) = 0$  for all  $i > 0$ .

### Definition 2

We say  $C$  is **stably degenerate** or **stably contained in  $S$** , if for every small global deformation  $C'$  of  $C$  in  $X$ , there exists a global deformation  $S'$  of  $S$  in  $X$  such that  $S' \subset C'$ .

We have the following implications:

$$(1) \Rightarrow (2) \Rightarrow (3)$$

where

- ①  $H^1(\mathcal{I}_{C/S} \otimes_S N_{S/X}) = 0$ .
- ②  $pr_1 : \text{HF } X \rightarrow \text{Hilb } X$ ,  $(C', S') \mapsto [C']$  is smooth at  $(C, S)$ .
- ③  $C$  is stably degenerate and  $pr_1(\mathcal{W}_{C,S})$  is a component of  $\text{Hilb } X$ .

The implication “ $(2) \Rightarrow (3)$ ” follows from the fact that  $pr'_1 = pr_1|_{\mathcal{W}_{C,S}}$  is locally surjective at  $[C]$ .

## Stable degeneration

Let  $C \subset S \subset X$  a sequence of closed subvarieties with  $H^i(N_{(C,S)/X}) = 0$  for all  $i > 0$ .

### Definition 2

We say  $C$  is **stably degenerate** or **stably contained in  $S$** , if for every small global deformation  $C'$  of  $C$  in  $X$ , there exists a global deformation  $S'$  of  $S$  in  $X$  such that  $S' \subset C'$ .

We have the following implications:

$$(1) \Rightarrow (2) \Rightarrow (3)$$

where

- ①  $H^1(\mathcal{I}_{C/S} \otimes_S N_{S/X}) = 0$ .
- ②  $pr_1 : \text{HF } X \rightarrow \text{Hilb } X$ ,  $(C', S') \mapsto [C']$  is smooth at  $(C, S)$ .
- ③  $C$  is stably degenerate and  $pr_1(\mathcal{W}_{C,S})$  is a component of  $\text{Hilb } X$ .

The implication “ $(2) \Rightarrow (3)$ ” follows from the fact that  $pr'_1 = pr_1|_{\mathcal{W}_{C,S}}$  is locally surjective at  $[C]$ .

## Stable degeneration

Let  $C \subset S \subset X$  a sequence of closed subvarieties with  $H^i(N_{(C,S)/X}) = 0$  for all  $i > 0$ .

### Definition 2

We say  $C$  is **stably degenerate** or **stably contained in  $S$** , if for every small global deformation  $C'$  of  $C$  in  $X$ , there exists a global deformation  $S'$  of  $S$  in  $X$  such that  $S' \subset C'$ .

We have the following implications:

$$(1) \Rightarrow (2) \Rightarrow (3)$$

where

- ①  $H^1(\mathcal{I}_{C/S} \otimes_S N_{S/X}) = 0$ .
- ②  $pr_1 : \text{HF } X \rightarrow \text{Hilb } X$ ,  $(C', S') \mapsto [C']$  is smooth at  $(C, S)$ .
- ③  $C$  is stably degenerate and  $pr_1(\mathcal{W}_{C,S})$  is a component of  $\text{Hilb } X$ .

The implication “ $(2) \Rightarrow (3)$ ” follows from the fact that  $pr'_1 = pr_1|_{\mathcal{W}_{C,S}}$  is locally surjective at  $[C]$ .

## Stable degeneration

Let  $C \subset S \subset X$  a sequence of closed subvarieties with  $H^i(N_{(C,S)/X}) = 0$  for all  $i > 0$ .

### Definition 2

We say  $C$  is **stably degenerate** or **stably contained in  $S$** , if for every small global deformation  $C'$  of  $C$  in  $X$ , there exists a global deformation  $S'$  of  $S$  in  $X$  such that  $S' \subset C'$ .

We have the following implications:

$$(1) \Rightarrow (2) \Rightarrow (3)$$

where

- ①  $H^1(\mathcal{I}_{C/S} \otimes_S N_{S/X}) = 0$ .
- ②  $pr_1 : \text{HF } X \rightarrow \text{Hilb } X$ ,  $(C', S') \mapsto [C']$  is smooth at  $(C, S)$ .
- ③  $C$  is stably degenerate and  $pr_1(\mathcal{W}_{C,S})$  is a component of  $\text{Hilb } X$ .

The implication “ $(2) \Rightarrow (3)$ ” follows from the fact that  $pr'_1 = pr_1|_{\mathcal{W}_{C,S}}$  is locally surjective at  $[C]$ .

## Stable degeneration

Let  $C \subset S \subset X$  a sequence of closed subvarieties with  $H^i(N_{(C,S)/X}) = 0$  for all  $i > 0$ .

### Definition 2

We say  $C$  is **stably degenerate** or **stably contained in  $S$** , if for every small global deformation  $C'$  of  $C$  in  $X$ , there exists a global deformation  $S'$  of  $S$  in  $X$  such that  $S' \subset C'$ .

We have the following implications:

$$(1) \Rightarrow (2) \Rightarrow (3)$$

where

- ➊  $H^1(\mathcal{I}_{C/S} \otimes_S N_{S/X}) = 0$ .
- ➋  $pr_1 : \text{HF } X \rightarrow \text{Hilb } X$ ,  $(C', S') \mapsto [C']$  is smooth at  $(C, S)$ .
- ➌  $C$  is stably degenerate and  $pr_1(W_{C,S})$  is a component of  $\text{Hilb } X$ .

The implication “ $(2) \Rightarrow (3)$ ” follows from the fact that  $pr'_1 = pr_1|_{W_{C,S}}$  is locally surjective at  $[C]$ .

## Stable degeneration

Let  $C \subset S \subset X$  a sequence of closed subvarieties with  $H^i(N_{(C,S)/X}) = 0$  for all  $i > 0$ .

### Definition 2

We say  $C$  is **stably degenerate** or **stably contained in  $S$** , if for every small global deformation  $C'$  of  $C$  in  $X$ , there exists a global deformation  $S'$  of  $S$  in  $X$  such that  $S' \subset C'$ .

We have the following implications:

$$(1) \Rightarrow (2) \Rightarrow (3)$$

where

- ➊  $H^1(I_{C/S} \otimes_S N_{S/X}) = 0$ .
- ➋  $pr_1 : \mathbf{HF} X \rightarrow \mathbf{Hilb} X, (C', S') \mapsto [C']$  is smooth at  $(C, S)$ .
- ➌  $C$  is stably degenerate and  $pr_1(W_{C,S})$  is a component of  $\mathbf{Hilb} X$ .

The implication “(2)  $\Rightarrow$  (3)” follows from the fact that  $pr'_1 = pr_1|_{W_{C,S}}$  is locally surjective at  $[C]$ .

## Stable degeneration

Let  $C \subset S \subset X$  a sequence of closed subvarieties with  $H^i(N_{(C,S)/X}) = 0$  for all  $i > 0$ .

### Definition 2

We say  $C$  is **stably degenerate** or **stably contained in  $S$** , if for every small global deformation  $C'$  of  $C$  in  $X$ , there exists a global deformation  $S'$  of  $S$  in  $X$  such that  $S' \subset C'$ .

We have the following implications:

$$(1) \Rightarrow (2) \Rightarrow (3)$$

where

- ①  $H^1(\mathcal{I}_{C/S} \otimes_S N_{S/X}) = 0$ .
- ②  $pr_1 : \mathbf{HF} X \rightarrow \mathbf{Hilb} X$ ,  $(C', S') \mapsto [C']$  is smooth at  $(C, S)$ .
- ③  $C$  is stably degenerate and  $pr_1(\mathcal{W}_{C,S})$  is a component of  $\mathbf{Hilb} X$ .

The implication “ $(2) \Rightarrow (3)$ ” follows from the fact that  $pr'_1 = pr_1|_{\mathcal{W}_{C,S}}$  is locally surjective at  $[C]$ .

## §2 Kleppe-Ellia conjecture and its generalization

## Mumford's example and Kleppe's generalization

Given a projective scheme, we denote by  $\mathbf{Hilb}^{sc} X$  the Hilbert scheme of smooth connected curves in  $X$ , i.e.,

$$\mathbf{Hilb}^{sc} X = \{C \subset X \mid C: \text{smooth connected curve}\}.$$

Theorem 3 (Mumford'62, a pathology)

$\mathbf{Hilb}^{sc} \mathbb{P}^3$  contains a generically non-reduced component.

Every its general member, i.e., a smooth curve  $C \subset \mathbb{P}^3$ , was contained in a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ .

Later, Kleppe['87] generalized this example systematically by using the coordinate of  $(a; b_1, \dots, b_6)$  of  $[C]$  in  $\mathbf{Pic} S_3 \simeq \mathbb{Z}^7$ :

$$C \sim al - \sum_{i=1}^6 b_i e_i \quad \longleftrightarrow \quad (a; b_1, \dots, b_6)$$

where  $l = [\mathcal{O}_{\mathbb{P}^2}(1)]$  and  $e_i$  ( $i = 1, \dots, 6$ ) are 6 exceptional curves.

## Mumford's example and Kleppe's generalization

Given a projective scheme, we denote by  $\mathbf{Hilb}^{sc} X$  the Hilbert scheme of smooth connected curves in  $X$ , i.e.,

$$\mathbf{Hilb}^{sc} X = \{C \subset X \mid C: \text{smooth connected curve}\}.$$

Theorem 3 (Mumford'62, a pathology)

$\mathbf{Hilb}^{sc} \mathbb{P}^3$  contains a generically non-reduced component.

Every its general member, i.e., a smooth curve  $C \subset \mathbb{P}^3$ , was contained in a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ .

Later, Kleppe['87] generalized this example systematically by using the coordinate of  $(a; b_1, \dots, b_6)$  of  $[C]$  in  $\mathbf{Pic} S_3 \simeq \mathbb{Z}^7$ :

$$C \sim al - \sum_{i=1}^6 b_i e_i \quad \longleftrightarrow \quad (a; b_1, \dots, b_6)$$

where  $l = [\mathcal{O}_{\mathbb{P}^2}(1)]$  and  $e_i$  ( $i = 1, \dots, 6$ ) are 6 exceptional curves.

## Mumford's example and Kleppe's generalization

Given a projective scheme, we denote by  $\mathbf{Hilb}^{sc} X$  the Hilbert scheme of smooth connected curves in  $X$ , i.e.,

$$\mathbf{Hilb}^{sc} X = \{C \subset X \mid C: \text{smooth connected curve}\}.$$

Theorem 3 (Mumford'62, a pathology)

$\mathbf{Hilb}^{sc} \mathbb{P}^3$  contains a **generically non-reduced** component.

Every its general member, i.e., a smooth curve  $C \subset \mathbb{P}^3$ , was contained in a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ .

Later, Kleppe['87] generalized this example systematically by using the coordinate of  $(a; b_1, \dots, b_6)$  of  $[C]$  in  $\mathbf{Pic} S_3 \simeq \mathbb{Z}^7$ :

$$C \sim al - \sum_{i=1}^6 b_i e_i \quad \longleftrightarrow \quad (a; b_1, \dots, b_6)$$

where  $l = [\mathcal{O}_{\mathbb{P}^2}(1)]$  and  $e_i$  ( $i = 1, \dots, 6$ ) are **6** exceptional curves.

## Mumford's example and Kleppe's generalization

Given a projective scheme, we denote by  $\mathbf{Hilb}^{sc} X$  the Hilbert scheme of smooth connected curves in  $X$ , i.e.,

$$\mathbf{Hilb}^{sc} X = \{C \subset X \mid C: \text{smooth connected curve}\}.$$

Theorem 3 (Mumford'62, a pathology)

$\mathbf{Hilb}^{sc} \mathbb{P}^3$  contains a **generically non-reduced** component.

Every its general member, i.e., a smooth curve  $C \subset \mathbb{P}^3$ , was contained in a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ .

Later, Kleppe['87] generalized this example systematically by using the coordinate of  $(a; b_1, \dots, b_6)$  of  $[C]$  in  $\mathbf{Pic} S_3 \simeq \mathbb{Z}^7$ :

$$C \sim al - \sum_{i=1}^6 b_i e_i \quad \longleftrightarrow \quad (a; b_1, \dots, b_6)$$

where  $l = [\mathcal{O}_{\mathbb{P}^2}(1)]$  and  $e_i$  ( $i = 1, \dots, 6$ ) are **6** exceptional curves.

## Mumford's example and Kleppe's generalization

Given a projective scheme, we denote by  $\mathbf{Hilb}^{sc} X$  the Hilbert scheme of smooth connected curves in  $X$ , i.e.,

$$\mathbf{Hilb}^{sc} X = \{C \subset X \mid C: \text{smooth connected curve}\}.$$

Theorem 3 (Mumford'62, a pathology)

$\mathbf{Hilb}^{sc} \mathbb{P}^3$  contains a **generically non-reduced** component.

Every its general member, i.e., a smooth curve  $C \subset \mathbb{P}^3$ , was contained in a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ .

Later, Kleppe['87] generalized this example systematically by using the coordinate of  $(a; b_1, \dots, b_6)$  of  $[C]$  in  $\mathbf{Pic} S_3 \simeq \mathbb{Z}^7$ :

$$C \sim al - \sum_{i=1}^6 b_i e_i \quad \longleftrightarrow \quad (a; b_1, \dots, b_6)$$

where  $l = [\mathcal{O}_{\mathbb{P}^2}(1)]$  and  $e_i$  ( $i = 1, \dots, 6$ ) are 6 exceptional curves.

## Mumford's example and Kleppe's generalization

Given a projective scheme, we denote by  $\mathbf{Hilb}^{sc} X$  the Hilbert scheme of smooth connected curves in  $X$ , i.e.,

$$\mathbf{Hilb}^{sc} X = \{C \subset X \mid C: \text{smooth connected curve}\}.$$

Theorem 3 (Mumford'62, a pathology)

$\mathbf{Hilb}^{sc} \mathbb{P}^3$  contains a **generically non-reduced** component.

Every its general member, i.e., a smooth curve  $C \subset \mathbb{P}^3$ , was contained in a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ .

Later, Kleppe['87] generalized this example systematically by using the coordinate of  $(a; b_1, \dots, b_6)$  of  $[C]$  in  $\mathbf{Pic} S_3 \simeq \mathbb{Z}^7$ :

$$C \sim a\mathbf{l} - \sum_{i=1}^6 b_i \mathbf{e}_i \quad \longleftrightarrow \quad (a; b_1, \dots, b_6)$$

where  $\mathbf{l} = [\mathcal{O}_{\mathbb{P}^2}(1)]$  and  $\mathbf{e}_i$  ( $i = 1, \dots, 6$ ) are 6 exceptional curves.

## Mumford's example and Kleppe's generalization

Given a projective scheme, we denote by  $\mathbf{Hilb}^{sc} X$  the Hilbert scheme of smooth connected curves in  $X$ , i.e.,

$$\mathbf{Hilb}^{sc} X = \{C \subset X \mid C: \text{smooth connected curve}\}.$$

Theorem 3 (Mumford'62, a pathology)

$\mathbf{Hilb}^{sc} \mathbb{P}^3$  contains a **generically non-reduced** component.

Every its general member, i.e., a smooth curve  $C \subset \mathbb{P}^3$ , was contained in a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ .

Later, Kleppe['87] generalized this example systematically by using the coordinate of  $(a; b_1, \dots, b_6)$  of  $[C]$  in  $\mathbf{Pic} S_3 \simeq \mathbb{Z}^7$ :

$$C \sim a\mathbf{l} - \sum_{i=1}^6 b_i \mathbf{e}_i \quad \longleftrightarrow \quad (a; b_1, \dots, b_6)$$

where  $\mathbf{l} = [\mathcal{O}_{\mathbb{P}^2}(1)]$  and  $\mathbf{e}_i$  ( $i = 1, \dots, 6$ ) are 6 exceptional curves.

## Mumford's example and Kleppe's generalization

Given a projective scheme, we denote by  $\mathbf{Hilb}^{sc} X$  the Hilbert scheme of smooth connected curves in  $X$ , i.e.,

$$\mathbf{Hilb}^{sc} X = \{C \subset X \mid C: \text{smooth connected curve}\}.$$

Theorem 3 (Mumford'62, a pathology)

$\mathbf{Hilb}^{sc} \mathbb{P}^3$  contains a **generically non-reduced** component.

Every its general member, i.e., a smooth curve  $C \subset \mathbb{P}^3$ , was contained in a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ .

Later, Kleppe['87] generalized this example systematically by using the coordinate of  $(a; b_1, \dots, b_6)$  of  $[C]$  in  $\mathbf{Pic} S_3 \simeq \mathbb{Z}^7$ :

$$C \sim al - \sum_{i=1}^6 b_i e_i \quad \longleftrightarrow \quad (a; b_1, \dots, b_6)$$

where  $l = [O_{\mathbb{P}^2}(1)]$  and  $e_i$  ( $i = 1, \dots, 6$ ) are **6** exceptional curves.

## Kleppe-Ellia conjecture

He proposed a conjecture, which can be reformulated as follows:

Conjecture (Kleppe'87, modified by Ellia'87)

Let  $C \subset S_3 \subset \mathbb{P}^3$  be a smooth curve of degree  $d \geq 14$  and genus  $g$  lying on a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ . Then  $C$  is stably degenerate if

- ①  $\chi(\mathcal{I}_C(3)) \geq 1$  ( $\Leftrightarrow g \geq 3d - 18$ ),
- ②  $C$  is linearly normal, and
- ③  $C$  is general in  $[C] \in \text{Pic } S_3$ .

K-E conj. is non-trivial only if  $C$  is not 3-normal ( $\Leftrightarrow H^1(\mathcal{I}_C(3)) \neq 0$ ). In fact, otherwise,  $pr_1 : \text{HF } \mathbb{P}^3 \rightarrow \text{Hilb } \mathbb{P}^3$  is smooth at  $(C, S)$  by

$$H^0(N_{(C,S)/\mathbb{P}^3}) \xrightarrow{p_1} H^0(N_{C/\mathbb{P}^3}) \longrightarrow \underbrace{H^1(N_{S/\mathbb{P}^3}(-C))}_{\simeq H^1(\mathcal{I}_C(3))} \longrightarrow 0.$$

Moreover, the first two assumptions are necessary for the conclusion.

## Kleppe-Ellia conjecture

He proposed a conjecture, which can be reformulated as follows:

Conjecture (Kleppe'87, modified by Ellia'87)

Let  $C \subset S_3 \subset \mathbb{P}^3$  be a smooth curve of degree  $d \geq 14$  and genus  $g$  lying on a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ . Then  $C$  is stably degenerate if

- ①  $\chi(\mathcal{I}_C(3)) \geq 1$  ( $\Leftrightarrow g \geq 3d - 18$ ),
- ②  $C$  is linearly normal, and
- ③  $C$  is general in  $[C] \in \text{Pic } S_3$ .

K-E conj. is non-trivial only if  $C$  is not 3-normal ( $\Leftrightarrow H^1(\mathcal{I}_C(3)) \neq 0$ ). In fact, otherwise,  $pr_1 : \text{HF } \mathbb{P}^3 \rightarrow \text{Hilb } \mathbb{P}^3$  is smooth at  $(C, S)$  by

$$H^0(N_{(C,S)/\mathbb{P}^3}) \xrightarrow{p_1} H^0(N_{C/\mathbb{P}^3}) \longrightarrow \underbrace{H^1(N_{S/\mathbb{P}^3}(-C))}_{\simeq H^1(\mathcal{I}_C(3))} \longrightarrow 0.$$

Moreover, the first two assumptions are necessary for the conclusion.

## Kleppe-Ellia conjecture

He proposed a conjecture, which can be reformulated as follows:

### Conjecture (Kleppe'87, modified by Ellia'87)

Let  $C \subset S_3 \subset \mathbb{P}^3$  be a smooth curve of degree  $d \geq 14$  and genus  $g$  lying on a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ . Then  $C$  is stably degenerate if

- ①  $\chi(\mathcal{I}_C(3)) \geq 1$  ( $\Leftrightarrow g \geq 3d - 18$ ),
- ②  $C$  is linearly normal, and
- ③  $C$  is general in  $[C] \in \text{Pic } S_3$ .

K-E conj. is non-trivial only if  $C$  is not 3-normal ( $\Leftrightarrow H^1(\mathcal{I}_C(3)) \neq 0$ ). In fact, otherwise,  $pr_1 : \text{HF } \mathbb{P}^3 \rightarrow \text{Hilb } \mathbb{P}^3$  is smooth at  $(C, S)$  by

$$H^0(N_{(C,S)/\mathbb{P}^3}) \xrightarrow{p_1} H^0(N_{C/\mathbb{P}^3}) \longrightarrow \underbrace{H^1(N_{S/\mathbb{P}^3}(-C))}_{\simeq H^1(\mathcal{I}_C(3))} \longrightarrow 0.$$

Moreover, the first two assumptions are necessary for the conclusion.

## Kleppe-Ellia conjecture

He proposed a conjecture, which can be reformulated as follows:

### Conjecture (Kleppe'87, modified by Ellia'87)

Let  $C \subset S_3 \subset \mathbb{P}^3$  be a smooth curve of degree  $d \geq 14$  and genus  $g$  lying on a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ . Then  $C$  is stably degenerate if

- ①  $\chi(\mathcal{I}_C(3)) \geq 1$  ( $\Leftrightarrow g \geq 3d - 18$ ),
- ②  $C$  is linearly normal, and
- ③  $C$  is general in  $[C] \in \text{Pic } S_3$ .

K-E conj. is non-trivial only if  $C$  is not 3-normal ( $\Leftrightarrow H^1(\mathcal{I}_C(3)) \neq 0$ ). In fact, otherwise,  $pr_1 : \text{HF } \mathbb{P}^3 \rightarrow \text{Hilb } \mathbb{P}^3$  is smooth at  $(C, S)$  by

$$H^0(N_{(C,S)/\mathbb{P}^3}) \xrightarrow{p_1} H^0(N_{C/\mathbb{P}^3}) \longrightarrow \underbrace{H^1(N_{S/\mathbb{P}^3}(-C))}_{\simeq H^1(\mathcal{I}_C(3))} \longrightarrow 0.$$

Moreover, the first two assumptions are necessary for the conclusion.

## Kleppe-Ellia conjecture

He proposed a conjecture, which can be reformulated as follows:

### Conjecture (Kleppe'87, modified by Ellia'87)

Let  $C \subset S_3 \subset \mathbb{P}^3$  be a smooth curve of degree  $d \geq 14$  and genus  $g$  lying on a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ . Then  $C$  is stably degenerate if

- ①  $\chi(\mathcal{I}_C(3)) \geq 1$  ( $\Leftrightarrow g \geq 3d - 18$ ),
- ②  $C$  is linearly normal, and
- ③  $C$  is general in  $[C] \in \text{Pic } S_3$ .

K-E conj. is non-trivial only if  $C$  is not 3-normal ( $\Leftrightarrow H^1(\mathcal{I}_C(3)) \neq 0$ ). In fact, otherwise,  $pr_1 : \text{HF } \mathbb{P}^3 \rightarrow \text{Hilb } \mathbb{P}^3$  is smooth at  $(C, S)$  by

$$H^0(N_{(C,S)/\mathbb{P}^3}) \xrightarrow{p_1} H^0(N_{C/\mathbb{P}^3}) \longrightarrow \underbrace{H^1(N_{S/\mathbb{P}^3}(-C))}_{\simeq H^1(\mathcal{I}_C(3))} \longrightarrow 0.$$

Moreover, the first two assumptions are necessary for the conclusion.

## Kleppe-Ellia conjecture

He proposed a conjecture, which can be reformulated as follows:

### Conjecture (Kleppe'87, modified by Ellia'87)

Let  $C \subset S_3 \subset \mathbb{P}^3$  be a smooth curve of degree  $d \geq 14$  and genus  $g$  lying on a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ . Then  $C$  is stably degenerate if

- ①  $\chi(\mathcal{I}_C(3)) \geq 1$  ( $\Leftrightarrow g \geq 3d - 18$ ),
- ②  $C$  is linearly normal, and
- ③  $C$  is general in  $[C] \in \text{Pic } S_3$ .

K-E conj. is non-trivial only if  $C$  is not 3-normal ( $\Leftrightarrow H^1(\mathcal{I}_C(3)) \neq 0$ ). In fact, otherwise,  $pr_1 : \text{HF } \mathbb{P}^3 \rightarrow \text{Hilb } \mathbb{P}^3$  is smooth at  $(C, S)$  by

$$H^0(N_{(C,S)/\mathbb{P}^3}) \xrightarrow{p_1} H^0(N_{C/\mathbb{P}^3}) \longrightarrow \underbrace{H^1(N_{S/\mathbb{P}^3}(-C))}_{\simeq H^1(\mathcal{I}_C(3))} \longrightarrow 0.$$

Moreover, the first two assumptions are necessary for the conclusion.

## Kleppe-Ellia conjecture

He proposed a conjecture, which can be reformulated as follows:

### Conjecture (Kleppe'87, modified by Ellia'87)

Let  $C \subset S_3 \subset \mathbb{P}^3$  be a smooth curve of degree  $d \geq 14$  and genus  $g$  lying on a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ . Then  $C$  is stably degenerate if

- ①  $\chi(\mathcal{I}_C(3)) \geq 1$  ( $\Leftrightarrow g \geq 3d - 18$ ),
- ②  $C$  is linearly normal, and
- ③  $C$  is general in  $[C] \in \text{Pic } S_3$ .

K-E conj. is non-trivial only if  $C$  is not 3-normal ( $\Leftrightarrow H^1(\mathcal{I}_C(3)) \neq 0$ ). In fact, otherwise,  $pr_1 : \text{HF } \mathbb{P}^3 \rightarrow \text{Hilb } \mathbb{P}^3$  is smooth at  $(C, S)$  by

$$H^0(N_{(C,S)/\mathbb{P}^3}) \xrightarrow{p_1} H^0(N_{C/\mathbb{P}^3}) \longrightarrow \underbrace{H^1(N_{S/\mathbb{P}^3}(-C))}_{\simeq H^1(\mathcal{I}_C(3))} \longrightarrow 0.$$

Moreover, the first two assumptions are necessary for the conclusion.

## Kleppe-Ellia conjecture

He proposed a conjecture, which can be reformulated as follows:

### Conjecture (Kleppe'87, modified by Ellia'87)

Let  $C \subset S_3 \subset \mathbb{P}^3$  be a smooth curve of degree  $d \geq 14$  and genus  $g$  lying on a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ . Then  $C$  is stably degenerate if

- ①  $\chi(\mathcal{I}_C(3)) \geq 1$  ( $\Leftrightarrow g \geq 3d - 18$ ),
- ②  $C$  is linearly normal, and
- ③  $C$  is general in  $[C] \in \text{Pic } S_3$ .

K-E conj. is non-trivial only if  $C$  is not 3-normal ( $\Leftrightarrow H^1(\mathcal{I}_C(3)) \neq 0$ ). In fact, otherwise,  $pr_1 : \text{HF } \mathbb{P}^3 \rightarrow \text{Hilb } \mathbb{P}^3$  is smooth at  $(C, S)$  by

$$H^0(N_{(C,S)/\mathbb{P}^3}) \xrightarrow{p_1} H^0(N_{C/\mathbb{P}^3}) \longrightarrow \underbrace{H^1(N_{S/\mathbb{P}^3}(-C))}_{\simeq H^1(\mathcal{I}_C(3))} \longrightarrow 0.$$

Moreover, the first two assumptions are necessary for the conclusion.

## Kleppe-Ellia conjecture

He proposed a conjecture, which can be reformulated as follows:

### Conjecture (Kleppe'87, modified by Ellia'87)

Let  $C \subset S_3 \subset \mathbb{P}^3$  be a smooth curve of degree  $d \geq 14$  and genus  $g$  lying on a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ . Then  $C$  is stably degenerate if

- ①  $\chi(\mathcal{I}_C(3)) \geq 1$  ( $\Leftrightarrow g \geq 3d - 18$ ),
- ②  $C$  is linearly normal, and
- ③  $C$  is general in  $[C] \in \text{Pic } S_3$ .

K-E conj. is non-trivial only if  $C$  is not 3-normal ( $\Leftrightarrow H^1(\mathcal{I}_C(3)) \neq 0$ ). In fact, otherwise,  $pr_1 : \mathbf{HF} \mathbb{P}^3 \rightarrow \mathbf{Hilb} \mathbb{P}^3$  is smooth at  $(C, S)$  by

$$H^0(N_{(C,S)/\mathbb{P}^3}) \xrightarrow{p_1} H^0(N_{C/\mathbb{P}^3}) \longrightarrow \underbrace{H^1(N_{S/\mathbb{P}^3}(-C))}_{\simeq H^1(\mathcal{I}_C(3))} \longrightarrow 0.$$

Moreover, the first two assumptions are necessary for the conclusion.

## Kleppe-Ellia conjecture

He proposed a conjecture, which can be reformulated as follows:

**Conjecture (Kleppe'87, modified by Ellia'87)**

Let  $C \subset S_3 \subset \mathbb{P}^3$  be a smooth curve of degree  $d \geq 14$  and genus  $g$  lying on a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ . Then  $C$  is stably degenerate if

- ①  $\chi(\mathcal{I}_C(3)) \geq 1$  ( $\Leftrightarrow g \geq 3d - 18$ ),
- ②  $C$  is linearly normal, and
- ③  $C$  is general in  $[C] \in \text{Pic } S_3$ .

K-E conj. is non-trivial only if  $C$  is not 3-normal ( $\Leftrightarrow H^1(\mathcal{I}_C(3)) \neq 0$ ). In fact, otherwise,  $pr_1 : \text{HF } \mathbb{P}^3 \rightarrow \text{Hilb } \mathbb{P}^3$  is smooth at  $(C, S)$  by

$$H^0(N_{(C,S)/\mathbb{P}^3}) \xrightarrow{p_1} H^0(N_{C/\mathbb{P}^3}) \longrightarrow \underbrace{H^1(N_{S/\mathbb{P}^3}(-C))}_{\simeq H^1(\mathcal{I}_C(3))} \longrightarrow 0.$$

Moreover, the first two assumptions are necessary for the conclusion.

## Kleppe-Ellia conjecture

He proposed a conjecture, which can be reformulated as follows:

### Conjecture (Kleppe'87, modified by Ellia'87)

Let  $C \subset S_3 \subset \mathbb{P}^3$  be a smooth curve of degree  $d \geq 14$  and genus  $g$  lying on a smooth cubic surface  $S_3 \subset \mathbb{P}^3$ . Then  $C$  is stably degenerate if

- ①  $\chi(\mathcal{I}_C(3)) \geq 1$  ( $\Leftrightarrow g \geq 3d - 18$ ),
- ②  $C$  is linearly normal, and
- ③  $C$  is general in  $[C] \in \text{Pic } S_3$ .

K-E conj. is non-trivial only if  $C$  is not 3-normal ( $\Leftrightarrow H^1(\mathcal{I}_C(3)) \neq 0$ ). In fact, otherwise,  $pr_1 : \text{HF } \mathbb{P}^3 \rightarrow \text{Hilb } \mathbb{P}^3$  is smooth at  $(C, S)$  by

$$H^0(N_{(C,S)/\mathbb{P}^3}) \xrightarrow{p_1} H^0(N_{C/\mathbb{P}^3}) \longrightarrow \underbrace{H^1(N_{S/\mathbb{P}^3}(-C))}_{\simeq H^1(\mathcal{I}_C(3))} \longrightarrow 0.$$

Moreover, the first two assumptions are necessary for the conclusion.

## Some remarks

K-E conjecture is Known to be true if

- $C$  is not 3-normal and  $g >> d$  (Kleppe'87 and Ellia'87), or
- $C$  is 2-normal, i.e.  $H^1(\mathcal{I}_C(2)) = 0$  (N'23)

## Some remarks

K-E conjecture is Known to be true if

- $C$  is not 3-normal and  $g >> d$  (Kleppe'87 and Ellia'87), or
- $C$  is 2-normal, i.e.  $H^1(\mathcal{I}_C(2)) = 0$  (N'23)

## Some remarks

K-E conjecture is Known to be true if

- $C$  is not 3-normal and  $g >> d$  (Kleppe'87 and Ellia'87), or
- $C$  is 2-normal, i.e.  $H^1(\mathcal{I}_C(2)) = \mathbf{0}$  (N'23)

# Del Pezzo surfaces

## Definition 4

A smooth projective surface  $S$  is called **del Pezzo** if  $-K_S$  is ample.

Every del Pezzo surface  $S$  is isomorphic to a blow-up of  $\mathbb{P}^2$  (at  $9 - n$  points) or  $\mathbb{P}^1 \times \mathbb{P}^1$ . The number  $n = (-K_S)^2$  is called the *degree* of  $S$ , and  $1 \leq n \leq 9$ .

## Example 1 (del Pezzo surfaces)

degree $n$	a description of $S_n$	$-K_S$
$\vdots$	$\vdots$	
3	cubic surface $S_3 \subset \mathbb{P}^3$	
4	quartic c.i. $S_{2,2} \subset \mathbb{P}^4$	
5	lin. section $[\mathrm{Gr}(2,5) \hookrightarrow \mathbb{P}^9] \cap \mathbb{L}^{(5)}$	v.a.
$\vdots$	$\vdots$	

# Del Pezzo surfaces

## Definition 4

A smooth projective surface  $S$  is called **del Pezzo** if  $-K_S$  is ample.

Every del Pezzo surface  $S$  is isomorphic to a blow-up of  $\mathbb{P}^2$  (at  $9 - n$  points) or  $\mathbb{P}^1 \times \mathbb{P}^1$ . The number  $n = (-K_S)^2$  is called the *degree* of  $S$ , and  $1 \leq n \leq 9$ .

## Example 1 (del Pezzo surfaces)

degree $n$	a description of $S_n$	$-K_S$
$\vdots$	$\vdots$	
3	cubic surface $S_3 \subset \mathbb{P}^3$	
4	quartic c.i. $S_{2,2} \subset \mathbb{P}^4$	
5	lin. section $[\mathrm{Gr}(2,5) \hookrightarrow \mathbb{P}^9] \cap \mathbb{L}^{(5)}$	v.a.
$\vdots$	$\vdots$	

# Del Pezzo surfaces

## Definition 4

A smooth projective surface  $S$  is called **del Pezzo** if  $-K_S$  is ample.

Every del Pezzo surface  $S$  is isomorphic to a blow-up of  $\mathbb{P}^2$  (at  $9 - n$  points) or  $\mathbb{P}^1 \times \mathbb{P}^1$ . The number  $n = (-K_S)^2$  is called the *degree* of  $S$ , and  $1 \leq n \leq 9$ .

## Example 1 (del Pezzo surfaces)

degree $n$	a description of $S_n$	$-K_S$
$\vdots$	$\vdots$	
3	cubic surface $S_3 \subset \mathbb{P}^3$	
4	quartic c.i. $S_{2,2} \subset \mathbb{P}^4$	
5	lin. section $[\mathrm{Gr}(2,5) \hookrightarrow \mathbb{P}^9] \cap \mathbb{L}^{(5)}$	v.a.
$\vdots$	$\vdots$	

# Del Pezzo surfaces

## Definition 4

A smooth projective surface  $S$  is called **del Pezzo** if  $-K_S$  is ample.

Every del Pezzo surface  $S$  is isomorphic to a blow-up of  $\mathbb{P}^2$  (at  $9 - n$  points) or  $\mathbb{P}^1 \times \mathbb{P}^1$ . The number  $n = (-K_S)^2$  is called the *degree* of  $S$ , and  $1 \leq n \leq 9$ .

## Example 1 (del Pezzo surfaces)

degree $n$	a description of $S_n$	$-K_S$
$\vdots$	$\vdots$	
3	cubic surface $S_3 \subset \mathbb{P}^3$	
4	quartic c.i. $S_{2,2} \subset \mathbb{P}^4$	
5	lin. section $[\mathrm{Gr}(2, 5) \hookrightarrow \mathbb{P}^9] \cap \mathbb{L}^{(5)}$	v.a.
$\vdots$	$\vdots$	

## Why on del Pezzo?

### Proposition 5 (smoothness of flag-scheme)

Let  $C \subset S = S_n \subset \mathbb{P}^n$  be a smooth curve of degree  $d$  and genus  $g$  lying on a del Pezzo surface  $S_n$  ( $n \geq 3$ ). Then the Hilbert-flag scheme  $\text{HF } \mathbb{P}^n$  is nonsingular at  $(C, S)$  of expected dimension

$$\chi(N_{(C,S)/\mathbb{P}^n}) = d + g + n^2 + 9,$$

and  $H^i(N_{(C,S)/\mathbb{P}^n}) = 0$  for all  $i > 0$ .

In fact,  $H^i(N_{C/S}) = H^i(N_{S/\mathbb{P}^n}) = 0$  for  $i > 0$ , which implies  $C \subset S$  and  $S \subset \mathbb{P}^n$  have nice (unobstructed) deformations and hence so does  $(C, S)$  in  $\mathbb{P}^n$ .

## Why on del Pezzo?

### Proposition 5 (smoothness of flag-scheme)

Let  $C \subset S = S_n \subset \mathbb{P}^n$  be a smooth curve of degree  $d$  and genus  $g$  lying on a del Pezzo surface  $S_n$  ( $n \geq 3$ ). Then the Hilbert-flag scheme  $\mathbf{HF} \mathbb{P}^n$  is nonsingular at  $(C, S)$  of expected dimension

$$\chi(N_{(C,S)/\mathbb{P}^n}) = d + g + n^2 + 9,$$

and  $H^i(N_{(C,S)/\mathbb{P}^n}) = 0$  for all  $i > 0$ .

In fact,  $H^i(N_{C/S}) = H^i(N_{S/\mathbb{P}^n}) = 0$  for  $i > 0$ , which implies  $C \subset S$  and  $S \subset \mathbb{P}^n$  have nice (unobstructed) deformations and hence so does  $(C, S)$  in  $\mathbb{P}^n$ .

## Why on del Pezzo?

### Proposition 5 (smoothness of flag-scheme)

Let  $C \subset S = S_n \subset \mathbb{P}^n$  be a smooth curve of degree  $d$  and genus  $g$  lying on a del Pezzo surface  $S_n$  ( $n \geq 3$ ). Then the Hilbert-flag scheme  $\text{HF } \mathbb{P}^n$  is nonsingular at  $(C, S)$  of expected dimension

$$\chi(N_{(C,S)/\mathbb{P}^n}) = d + g + n^2 + 9,$$

and  $H^i(N_{(C,S)/\mathbb{P}^n}) = 0$  for all  $i > 0$ .

In fact,  $H^i(N_{C/S}) = H^i(N_{S/\mathbb{P}^n}) = 0$  for  $i > 0$ , which implies  $C \subset S$  and  $S \subset \mathbb{P}^n$  have nice (unobstructed) deformations and hence so does  $(C, S)$  in  $\mathbb{P}^n$ .

## Why on del Pezzo?

### Proposition 5 (smoothness of flag-scheme)

Let  $C \subset S = S_n \subset \mathbb{P}^n$  be a smooth curve of degree  $d$  and genus  $g$  lying on a del Pezzo surface  $S_n$  ( $n \geq 3$ ). Then the Hilbert-flag scheme  $\mathbf{HF} \mathbb{P}^n$  is nonsingular at  $(C, S)$  of expected dimension

$$\chi(N_{(C,S)/\mathbb{P}^n}) = d + g + n^2 + 9,$$

and  $H^i(N_{(C,S)/\mathbb{P}^n}) = 0$  for all  $i > 0$ .

In fact,  $H^i(N_{C/S}) = H^i(N_{S/\mathbb{P}^n}) = 0$  for  $i > 0$ , which implies  $C \subset S$  and  $S \subset \mathbb{P}^n$  have nice (unobstructed) deformations and hence so does  $(C, S)$  in  $\mathbb{P}^n$ .

## Why on del Pezzo?

### Proposition 5 (smoothness of flag-scheme)

Let  $C \subset S = S_n \subset \mathbb{P}^n$  be a smooth curve of degree  $d$  and genus  $g$  lying on a del Pezzo surface  $S_n$  ( $n \geq 3$ ). Then the Hilbert-flag scheme  $\mathbf{HF} \mathbb{P}^n$  is nonsingular at  $(C, S)$  of expected dimension

$$\chi(N_{(C,S)/\mathbb{P}^n}) = d + g + n^2 + 9,$$

and  $H^i(N_{(C,S)/\mathbb{P}^n}) = 0$  for all  $i > 0$ .

In fact,  $H^i(N_{C/S}) = H^i(N_{S/\mathbb{P}^n}) = 0$  for  $i > 0$ , which implies  $C \subset S$  and  $S \subset \mathbb{P}^n$  have nice (unobstructed) deformations and hence so does  $(C, S)$  in  $\mathbb{P}^n$ .

## Why on del Pezzo?

### Proposition 5 (smoothness of flag-scheme)

Let  $C \subset S = S_n \subset \mathbb{P}^n$  be a smooth curve of degree  $d$  and genus  $g$  lying on a del Pezzo surface  $S_n$  ( $n \geq 3$ ). Then the Hilbert-flag scheme  $\mathbf{HF} \mathbb{P}^n$  is nonsingular at  $(C, S)$  of expected dimension

$$\chi(N_{(C,S)/\mathbb{P}^n}) = d + g + n^2 + 9,$$

and  $H^i(N_{(C,S)/\mathbb{P}^n}) = 0$  for all  $i > 0$ .

In fact,  $H^i(N_{C/S}) = H^i(N_{S/\mathbb{P}^n}) = 0$  for  $i > 0$ , which implies  $C \subset S$  and  $S \subset \mathbb{P}^n$  have nice (unobstructed) deformations and hence so does  $(C, S)$  in  $\mathbb{P}^n$ .

## Generalized Kleppe-Ellia conjecture

Toward a generalization, we study the deformations of space curves lying on a del Pezzo surface of degree.

Conjecture (generalized K-E conj.)

Let  $C \subset S_n \subset \mathbb{P}^n$  be a smooth connected curve lying on a smooth del Pezzo surface  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 3$ . Then  $C$  is stably degenerate if

- ①  $\chi(N_{S/\mathbb{P}^n}(-C)) \geq 0$ ,
- ②  $C$  is linearly normal,
- ③  $\deg(C) > 9$  for  $n = 3$  and  $\deg(C) > 2n$  for  $n \geq 4$ , and  $C$  is general in  $[C] \in \text{Pic } S_n$ .

Remark 6

$$(1) \iff \chi(N_{(C,S)/\mathbb{P}^n}) \geq \chi(N_{C/\mathbb{P}^n}).$$

## Generalized Kleppe-Ellia conjecture

Toward a generalization, we study the deformations of space curves lying on a del Pezzo surface of degree.

Conjecture (generalized K-E conj.)

Let  $C \subset S_n \subset \mathbb{P}^n$  be a smooth connected curve lying on a smooth del Pezzo surface  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 3$ . Then  $C$  is stably degenerate if

- ①  $\chi(N_{S/\mathbb{P}^n}(-C)) \geq 0$ ,
- ②  $C$  is linearly normal,
- ③  $\deg(C) > 9$  for  $n = 3$  and  $\deg(C) > 2n$  for  $n \geq 4$ , and  $C$  is general in  $[C] \in \text{Pic } S_n$ .

Remark 6

$$(1) \iff \chi(N_{(C,S)/\mathbb{P}^n}) \geq \chi(N_{C/\mathbb{P}^n}).$$

## Generalized Kleppe-Ellia conjecture

Toward a generalization, we study the deformations of space curves lying on a del Pezzo surface of degree.

Conjecture (generalized K-E conj.)

Let  $C \subset S_n \subset \mathbb{P}^n$  be a smooth connected curve lying on a smooth del Pezzo surface  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 3$ . Then  $C$  is stably degenerate if

- ①  $\chi(N_{S/\mathbb{P}^n}(-C)) \geq 0$ ,
- ②  $C$  is linearly normal,
- ③  $\deg(C) > 9$  for  $n = 3$  and  $\deg(C) > 2n$  for  $n \geq 4$ , and  $C$  is general in  $[C] \in \text{Pic } S_n$ .

Remark 6

$$(1) \iff \chi(N_{(C,S)/\mathbb{P}^n}) \geq \chi(N_{C/\mathbb{P}^n}).$$

## Generalized Kleppe-Ellia conjecture

Toward a generalization, we study the deformations of space curves lying on a del Pezzo surface of degree.

Conjecture (generalized K-E conj.)

Let  $C \subset S_n \subset \mathbb{P}^n$  be a smooth connected curve lying on a smooth del Pezzo surface  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 3$ . Then  $C$  is stably degenerate if

- ①  $\chi(N_{S/\mathbb{P}^n}(-C)) \geq 0$ ,
- ②  $C$  is linearly normal,
- ③  $\deg(C) > 9$  for  $n = 3$  and  $\deg(C) > 2n$  for  $n \geq 4$ , and  $C$  is general in  $[C] \in \text{Pic } S_n$ .

Remark 6

$$(1) \iff \chi(N_{(C,S)/\mathbb{P}^n}) \geq \chi(N_{C/\mathbb{P}^n}).$$

## Generalized Kleppe-Ellia conjecture

Toward a generalization, we study the deformations of space curves lying on a del Pezzo surface of degree.

Conjecture (generalized K-E conj.)

Let  $C \subset S_n \subset \mathbb{P}^n$  be a smooth connected curve lying on a smooth del Pezzo surface  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 3$ . Then  $C$  is stably degenerate if

- ①  $\chi(N_{S/\mathbb{P}^n}(-C)) \geq 0$ ,
- ②  $C$  is linearly normal,
- ③  $\deg(C) > 9$  for  $n = 3$  and  $\deg(C) > 2n$  for  $n \geq 4$ , and  $C$  is general in  $[C] \in \text{Pic } S_n$ .

Remark 6

$$(1) \iff \chi(N_{(C,S)/\mathbb{P}^n}) \geq \chi(N_{C/\mathbb{P}^n}).$$

## Generalized Kleppe-Ellia conjecture

Toward a generalization, we study the deformations of space curves lying on a del Pezzo surface of degree.

Conjecture (generalized K-E conj.)

Let  $C \subset S_n \subset \mathbb{P}^n$  be a smooth connected curve lying on a smooth del Pezzo surface  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 3$ . Then  $C$  is stably degenerate if

- ①  $\chi(N_{S/\mathbb{P}^n}(-C)) \geq 0$ ,
- ②  $C$  is linearly normal,
- ③  $\deg(C) > 9$  for  $n = 3$  and  $\deg(C) > 2n$  for  $n \geq 4$ , and  $C$  is general in  $[C] \in \text{Pic } S_n$ .

Remark 6

$$(1) \iff \chi(N_{(C,S)/\mathbb{P}^n}) \geq \chi(N_{C/\mathbb{P}^n}).$$

## Generalized Kleppe-Ellia conjecture

Toward a generalization, we study the deformations of space curves lying on a del Pezzo surface of degree.

Conjecture (generalized K-E conj.)

Let  $C \subset S_n \subset \mathbb{P}^n$  be a smooth connected curve lying on a smooth del Pezzo surface  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 3$ . Then  $C$  is stably degenerate if

- ①  $\chi(N_{S/\mathbb{P}^n}(-C)) \geq 0$ ,
- ②  $C$  is linearly normal,
- ③  $\deg(C) > 9$  for  $n = 3$  and  $\deg(C) > 2n$  for  $n \geq 4$ , and  $C$  is general in  $[C] \in \text{Pic } S_n$ .

Remark 6

$$(1) \iff \chi(N_{(C,S)/\mathbb{P}^n}) \geq \chi(N_{C/\mathbb{P}^n}).$$

## Main result

We focus on the case  $n = 4$ , i.e.  $S \simeq S_4$  is a smooth complete intersection

$$S_{2,2} = (2) \cap (2) \subset \mathbb{P}^4.$$

We see that  $N_{S/\mathbb{P}^4} \simeq \mathcal{O}_S(2)^{\oplus 2} \simeq \mathcal{O}_S(-2K_S)^{\oplus 2}$  and hence

$$H^1(N_{S/\mathbb{P}^4}(-C)) \simeq H^1(\mathcal{O}_S(-C - 2K_S))^{\oplus 2} \simeq H^1(\mathcal{I}_C(2))^{\oplus 2}.$$

Then it follows from a general theory that  $C$  is stably degenerate if  $C$  is 2-normal.

### Theorem 7 (N'25)

Let  $C \subset \mathbb{P}^4$  be a smooth connected curve of degree  $d > 8$  contained in a smooth c.i.  $S = S_{2,2} \subset \mathbb{P}^4$ . Then

- ① If  $C$  is 2-normal, then  $C$  is unobstructed and stably degenerate.
- ② If  $\mathcal{O}_C(2)$  is non-special and  $C$  is not 2-normal, then  $C$  is unobstructed, but not stably degenerate.
- ③ If  $h^1(\mathcal{O}_C(2)) \geq h^1(\mathcal{I}_C(2)) = 1$  and  $C$  is general in  $[C] \in \text{Pic } S$ , then  $C$  is obstructed and stably degenerate.

## Main result

We focus on the case  $n = 4$ , i.e.  $S \simeq S_4$  is a smooth complete intersection

$$S_{2,2} = (2) \cap (2) \subset \mathbb{P}^4.$$

We see that  $N_{S/\mathbb{P}^4} \simeq \mathcal{O}_S(2)^{\oplus 2} \simeq \mathcal{O}_S(-2K_S)^{\oplus 2}$  and hence

$$H^1(N_{S/\mathbb{P}^4}(-C)) \simeq H^1(\mathcal{O}_S(-C - 2K_S))^{\oplus 2} \simeq H^1(\mathcal{I}_C(2))^{\oplus 2}.$$

Then it follows from a general theory that  $C$  is stably degenerate if  $C$  is 2-normal.

### Theorem 7 (N'25)

Let  $C \subset \mathbb{P}^4$  be a smooth connected curve of degree  $d > 8$  contained in a smooth c.i.  $S = S_{2,2} \subset \mathbb{P}^4$ . Then

- ① If  $C$  is 2-normal, then  $C$  is unobstructed and stably degenerate.
- ② If  $\mathcal{O}_C(2)$  is non-special and  $C$  is not 2-normal, then  $C$  is unobstructed, but not stably degenerate.
- ③ If  $h^1(\mathcal{O}_C(2)) \geq h^1(\mathcal{I}_C(2)) = 1$  and  $C$  is general in  $[C] \in \text{Pic } S$ , then  $C$  is obstructed and stably degenerate.

## Main result

We focus on the case  $n = 4$ , i.e.  $S \simeq S_4$  is a smooth complete intersection

$$S_{2,2} = (2) \cap (2) \subset \mathbb{P}^4.$$

We see that  $N_{S/\mathbb{P}^4} \simeq \mathcal{O}_S(2)^{\oplus 2} \simeq \mathcal{O}_S(-2K_S)^{\oplus 2}$  and hence

$$H^1(N_{S/\mathbb{P}^4}(-C)) \simeq H^1(\mathcal{O}_S(-C - 2K_S))^{\oplus 2} \simeq H^1(\mathcal{I}_C(2))^{\oplus 2}.$$

Then it follows from a general theory that  $C$  is stably degenerate if  $C$  is 2-normal.

### Theorem 7 (N'25)

Let  $C \subset \mathbb{P}^4$  be a smooth connected curve of degree  $d > 8$  contained in a smooth c.i.  $S = S_{2,2} \subset \mathbb{P}^4$ . Then

- ① If  $C$  is 2-normal, then  $C$  is unobstructed and stably degenerate.
- ② If  $\mathcal{O}_C(2)$  is non-special and  $C$  is not 2-normal, then  $C$  is unobstructed, but not stably degenerate.
- ③ If  $h^1(\mathcal{O}_C(2)) \geq h^1(\mathcal{I}_C(2)) = 1$  and  $C$  is general in  $[C] \in \text{Pic } S$ , then  $C$  is obstructed and stably degenerate.

## Main result

We focus on the case  $n = 4$ , i.e.  $S \simeq S_4$  is a smooth complete intersection

$$S_{2,2} = (2) \cap (2) \subset \mathbb{P}^4.$$

We see that  $N_{S/\mathbb{P}^4} \simeq \mathcal{O}_S(2)^{\oplus 2} \simeq \mathcal{O}_S(-2K_S)^{\oplus 2}$  and hence

$$H^1(N_{S/\mathbb{P}^4}(-C)) \simeq H^1(\mathcal{O}_S(-C - 2K_S))^{\oplus 2} \simeq H^1(\mathcal{I}_C(2))^{\oplus 2}.$$

Then it follows from a general theory that  $C$  is stably degenerate if  $C$  is 2-normal.

### Theorem 7 (N'25)

Let  $C \subset \mathbb{P}^4$  be a smooth connected curve of degree  $d > 8$  contained in a smooth c.i.  $S = S_{2,2} \subset \mathbb{P}^4$ . Then

- ① If  $C$  is 2-normal, then  $C$  is unobstructed and stably degenerate.
- ② If  $\mathcal{O}_C(2)$  is non-special and  $C$  is not 2-normal, then  $C$  is unobstructed, but not stably degenerate.
- ③ If  $h^1(\mathcal{O}_C(2)) \geq h^1(\mathcal{I}_C(2)) = 1$  and  $C$  is general in  $[C] \in \text{Pic } S$ , then  $C$  is obstructed and stably degenerate.

## Main result

We focus on the case  $n = 4$ , i.e.  $S \simeq S_4$  is a smooth complete intersection

$$S_{2,2} = (2) \cap (2) \subset \mathbb{P}^4.$$

We see that  $N_{S/\mathbb{P}^4} \simeq \mathcal{O}_S(2)^{\oplus 2} \simeq \mathcal{O}_S(-2K_S)^{\oplus 2}$  and hence

$$H^1(N_{S/\mathbb{P}^4}(-C)) \simeq H^1(\mathcal{O}_S(-C - 2K_S))^{\oplus 2} \simeq H^1(\mathcal{I}_C(2))^{\oplus 2}.$$

Then it follows from a general theory that  $C$  is stably degenerate if  $C$  is 2-normal.

### Theorem 7 (N'25)

Let  $C \subset \mathbb{P}^4$  be a smooth connected curve of degree  $d > 8$  contained in a smooth c.i.  $S = S_{2,2} \subset \mathbb{P}^4$ . Then

- ① If  $C$  is 2-normal, then  $C$  is unobstructed and stably degenerate.
- ② If  $\mathcal{O}_C(2)$  is non-special and  $C$  is not 2-normal, then  $C$  is unobstructed, but not stably degenerate.
- ③ If  $h^1(\mathcal{O}_C(2)) \geq h^1(\mathcal{I}_C(2)) = 1$  and  $C$  is general in  $[C] \in \text{Pic } S$ , then  $C$  is obstructed and stably degenerate.

## Main result

We focus on the case  $n = 4$ , i.e.  $S \simeq S_4$  is a smooth complete intersection

$$S_{2,2} = (2) \cap (2) \subset \mathbb{P}^4.$$

We see that  $N_{S/\mathbb{P}^4} \simeq \mathcal{O}_S(2)^{\oplus 2} \simeq \mathcal{O}_S(-2K_S)^{\oplus 2}$  and hence

$$H^1(N_{S/\mathbb{P}^4}(-C)) \simeq H^1(\mathcal{O}_S(-C - 2K_S))^{\oplus 2} \simeq H^1(\mathcal{I}_C(2))^{\oplus 2}.$$

Then it follows from a general theory that  $C$  is stably degenerate if  $C$  is 2-normal.

### Theorem 7 (N'25)

Let  $C \subset \mathbb{P}^4$  be a smooth connected curve of degree  $d > 8$  contained in a smooth c.i.  $S = S_{2,2} \subset \mathbb{P}^4$ . Then

- ① If  $C$  is 2-normal, then  $C$  is unobstructed and stably degenerate.
- ② If  $\mathcal{O}_C(2)$  is non-special and  $C$  is not 2-normal, then  $C$  is unobstructed, but not stably degenerate.
- ③ If  $h^1(\mathcal{O}_C(2)) \geq h^1(\mathcal{I}_C(2)) = 1$  and  $C$  is general in  $[C] \in \text{Pic } S$ , then  $C$  is obstructed and stably degenerate.

## Main result

We focus on the case  $n = 4$ , i.e.  $S \simeq S_4$  is a smooth complete intersection

$$S_{2,2} = (2) \cap (2) \subset \mathbb{P}^4.$$

We see that  $N_{S/\mathbb{P}^4} \simeq \mathcal{O}_S(2)^{\oplus 2} \simeq \mathcal{O}_S(-2K_S)^{\oplus 2}$  and hence

$$H^1(N_{S/\mathbb{P}^4}(-C)) \simeq H^1(\mathcal{O}_S(-C - 2K_S))^{\oplus 2} \simeq H^1(\mathcal{I}_C(2))^{\oplus 2}.$$

Then it follows from a general theory that  $C$  is stably degenerate if  $C$  is 2-normal.

### Theorem 7 (N'25)

Let  $C \subset \mathbb{P}^4$  be a smooth connected curve of degree  $d > 8$  contained in a smooth c.i.  $S = S_{2,2} \subset \mathbb{P}^4$ . Then

- ① If  $C$  is 2-normal, then  $C$  is unobstructed and stably degenerate.
- ② If  $\mathcal{O}_C(2)$  is non-special and  $C$  is not 2-normal, then  $C$  is unobstructed, but not stably degenerate.
- ③ If  $h^1(\mathcal{O}_C(2)) \geq h^1(\mathcal{I}_C(2)) = 1$  and  $C$  is general in  $[C] \in \text{Pic } S$ , then  $C$  is obstructed and stably degenerate.

## Main result

We focus on the case  $n = 4$ , i.e.  $S \simeq S_4$  is a smooth complete intersection

$$S_{2,2} = (2) \cap (2) \subset \mathbb{P}^4.$$

We see that  $N_{S/\mathbb{P}^4} \simeq \mathcal{O}_S(2)^{\oplus 2} \simeq \mathcal{O}_S(-2K_S)^{\oplus 2}$  and hence

$$H^1(N_{S/\mathbb{P}^4}(-C)) \simeq H^1(\mathcal{O}_S(-C - 2K_S))^{\oplus 2} \simeq H^1(\mathcal{I}_C(2))^{\oplus 2}.$$

Then it follows from a general theory that  $C$  is stably degenerate if  $C$  is 2-normal.

### Theorem 7 (N'25)

Let  $C \subset \mathbb{P}^4$  be a smooth connected curve of degree  $d > 8$  contained in a smooth c.i.  $S = S_{2,2} \subset \mathbb{P}^4$ . Then

- ① If  $C$  is 2-normal, then  $C$  is unobstructed and stably degenerate.
- ② If  $\mathcal{O}_C(2)$  is non-special and  $C$  is not 2-normal, then  $C$  is unobstructed, but not stably degenerate.
- ③ If  $h^1(\mathcal{O}_C(2)) \geq h^1(\mathcal{I}_C(2)) = 1$  and  $C$  is general in  $[C] \in \text{Pic } S$ , then  $C$  is obstructed and stably degenerate.

## Main result

We focus on the case  $n = 4$ , i.e.  $S \simeq S_4$  is a smooth complete intersection

$$S_{2,2} = (2) \cap (2) \subset \mathbb{P}^4.$$

We see that  $N_{S/\mathbb{P}^4} \simeq \mathcal{O}_S(2)^{\oplus 2} \simeq \mathcal{O}_S(-2K_S)^{\oplus 2}$  and hence

$$H^1(N_{S/\mathbb{P}^4}(-C)) \simeq H^1(\mathcal{O}_S(-C - 2K_S))^{\oplus 2} \simeq H^1(\mathcal{I}_C(2))^{\oplus 2}.$$

Then it follows from a general theory that  $C$  is stably degenerate if  $C$  is 2-normal.


### Theorem 7 (N'25)

Let  $C \subset \mathbb{P}^4$  be a smooth connected curve of degree  $d > 8$  contained in a smooth c.i.  $S = S_{2,2} \subset \mathbb{P}^4$ . Then

- ① If  $C$  is 2-normal, then  $C$  is unobstructed and stably degenerate.
- ② If  $\mathcal{O}_C(2)$  is non-special and  $C$  is not 2-normal, then  $C$  is unobstructed, but not stably degenerate.
- ③ If  $h^1(\mathcal{O}_C(2)) \geq h^1(\mathcal{I}_C(2)) = 1$  and  $C$  is general in  $[C] \in \text{Pic } S$ , then  $C$  is obstructed and stably degenerate.


## Main result (continued)

$C$  is **2-normal** (resp.  $O_C(2)$  is **nonspecial**) if  $(d, g)$  belongs to the region [I] (resp. [IV] except the 6 pairs corresponding to  $\circ$ ).




## Main result (continued)

$C$  is **2-normal** (resp.  $O_C(2)$  is **nonspecial**) if  $(d, g)$  belongs to the region [I] (resp. [IV] except the 6 pairs corresponding to  $\circ$ ).




## Main result (continued)

$C$  is **2-normal** (resp.  $O_C(2)$  is **nonspecial**) if  $(d, g)$  belongs to the region [I] (resp. [IV] except the 6 pairs corresponding to  $\circ$ ).




## Main result (continued)

$C$  is **2-normal** (resp.  $\mathcal{O}_C(2)$  is **nonspecial**) if  $(d, g)$  belongs to the region [I] (resp. [IV] except the 6 pairs corresponding to  $\circ$ ).




## Main result (continued)

$C$  is **2-normal** (resp.  $O_C(2)$  is **nonspecial**) if  $(d, g)$  belongs to the region [I] (resp. [IV] except the 6 pairs corresponding to  $\circ$ ).



## Main result (continued)

$C$  is **2-normal** (resp.  $\mathcal{O}_C(2)$  is **nonspecial**) if  $(d, g)$  belongs to the region [I] (resp. [IV] except the 6 pairs corresponding to  $\circ$ ).



## Applications

### Corollary 8

Generalized K-E conjecture holds to be true, if  $n = 4$  and  $h^1(\mathcal{I}_C(2)) = 1$ .

Let  $C \subset S = S_{2,2} \subset \mathbb{P}^4$  be as in Theorem 7, and  $\mathcal{W}_{C,S}$  the irreducible component of  $\text{HF } \mathbb{P}^4$  passing through  $(C, S)$ , and put

$$W_{C,S} := \text{pr}_1(\mathcal{W}_{C,S}) \cap \text{Hilb}^{sc} \mathbb{P}^4.$$

Then we have the following 3 possibilities for  $W_{C,S}$ :

degeneration of $C$	Is $C$ obstructed?	$W_{C,S} \subset \text{Hilb}^{sc} \mathbb{P}^4$
stable	NO	gen.smooth component
stable	YES	gen.non-reduced component
unstable	YES/NO	not a component

## Applications

### Corollary 8

Generalized K-E conjecture holds to be true, if  $n = 4$  and  $h^1(\mathcal{I}_C(2)) = 1$ .

Let  $C \subset S = S_{2,2} \subset \mathbb{P}^4$  be as in Theorem 7, and  $\mathcal{W}_{C,S}$  the irreducible component of  $\text{HF } \mathbb{P}^4$  passing through  $(C, S)$ , and put

$$W_{C,S} := pr_1(\mathcal{W}_{C,S}) \cap \text{Hilb}^{sc} \mathbb{P}^4.$$

Then we have the following 3 possibilities for  $W_{C,S}$ :

degeneration of $C$	Is $C$ obstructed?	$W_{C,S} \subset \text{Hilb}^{sc} \mathbb{P}^4$
stable	NO	gen.smooth component
stable	YES	gen.non-reduced component
unstable	YES/NO	not a component

## Applications

### Corollary 8

Generalized K-E conjecture holds to be true, if  $n = 4$  and  $h^1(\mathcal{I}_C(2)) = 1$ .

Let  $C \subset S = S_{2,2} \subset \mathbb{P}^4$  be as in Theorem 7, and  $\mathcal{W}_{C,S}$  the irreducible component of  $\text{HF } \mathbb{P}^4$  passing through  $(C, S)$ , and put

$$W_{C,S} := \text{pr}_1(\mathcal{W}_{C,S}) \cap \text{Hilb}^{sc} \mathbb{P}^4.$$

Then we have the following 3 possibilities for  $W_{C,S}$ :

degeneration of $C$	Is $C$ obstructed?	$W_{C,S} \subset \text{Hilb}^{sc} \mathbb{P}^4$
stable	NO	gen.smooth component
stable	YES	gen.non-reduced component
unstable	YES/NO	not a component

## Applications

### Corollary 8

Generalized K-E conjecture holds to be true, if  $n = 4$  and  $h^1(\mathcal{I}_C(2)) = 1$ .

Let  $C \subset S = S_{2,2} \subset \mathbb{P}^4$  be as in Theorem 7, and  $\mathcal{W}_{C,S}$  the irreducible component of  $\mathbf{HF} \mathbb{P}^4$  passing through  $(C, S)$ , and put

$$W_{C,S} := pr_1(\mathcal{W}_{C,S}) \cap \mathbf{Hilb}^{sc} \mathbb{P}^4.$$

Then we have the following 3 possibilities for  $W_{C,S}$ :

degeneration of $C$	Is $C$ obstructed?	$W_{C,S} \subset \mathbf{Hilb}^{sc} \mathbb{P}^4$
stable	NO	gen.smooth component
stable	YES	gen.non-reduced component
unstable	YES/NO	not a component

## Applications

### Corollary 8

Generalized K-E conjecture holds to be true, if  $n = 4$  and  $h^1(\mathcal{I}_C(2)) = 1$ .

Let  $C \subset S = S_{2,2} \subset \mathbb{P}^4$  be as in Theorem 7, and  $\mathcal{W}_{C,S}$  the irreducible component of  $\text{HF } \mathbb{P}^4$  passing through  $(C, S)$ , and put

$$W_{C,S} := pr_1(\mathcal{W}_{C,S}) \cap \text{Hilb}^{sc} \mathbb{P}^4.$$

Then we have the following 3 possibilities for  $W_{C,S}$ :

degeneration of $C$	Is $C$ obstructed?	$W_{C,S} \subset \text{Hilb}^{sc} \mathbb{P}^4$
stable	NO	gen.smooth component
stable	YES	gen.non-reduced component
unstable	YES/NO	not a component

## Analogy of Mumford's example

### Example 2

**Hilb<sup>sc</sup>  $\mathbb{P}^4$**  contains a **generically non-reduced** irreducible component  $W$  whose general member  $C$  satisfies

- ①  $C$  is contained in a **smooth c.i.**  $S = S_{2,2} \subset \mathbb{P}^4$ ,
- ② there exists a **line  $E$**  on  $S$  such that  $C$  belongs to a complete linear system  $\Lambda := |-3K_S + 2E| (\simeq \mathbb{P}^{29})$  on  $S$ , and
- ③  $\dim W = 55$ ,  $h^0(C, N_{C/\mathbb{P}^4}) = 57$ , and  $C$  is of degree **14** and genus **16**.

$$\begin{array}{ccccccc}
 C & \in & W^{(55)} & \subset & \text{Hilb}^{\text{sc}} \mathbb{P}^4 \\
 \downarrow & & \downarrow \text{${\mathbb{P}}^{29}$-b'dle} & & \\
 (E, S) & \in & U^{(26)} & \subset & G \times G(2, H^0(O_{\mathbb{P}^4}(2))) \\
 \downarrow & & \downarrow G(2, 12)\text{-b'dle} & & \downarrow \\
 E & \in & G(2, 5)^{(6)} & = & \{\text{lines in } \mathbb{P}^4\}.
 \end{array}$$

## Analogy of Mumford's example

### Example 2

**Hilb<sup>sc</sup>  $\mathbb{P}^4$**  contains a **generically non-reduced** irreducible component  $W$  whose general member  $C$  satisfies

- ①  $C$  is contained in a **smooth c.i.**  $S = S_{2,2} \subset \mathbb{P}^4$ ,
- ② there exists a **line  $E$**  on  $S$  such that  $C$  belongs to a complete linear system  $\Lambda := |-3K_S + 2E| (\simeq \mathbb{P}^{29})$  on  $S$ , and
- ③  $\dim W = 55$ ,  $h^0(C, N_{C/\mathbb{P}^4}) = 57$ , and  $C$  is of degree **14** and genus **16**.

$$\begin{array}{ccccccc}
 C & \in & W^{(55)} & \subset & \text{Hilb}^{\text{sc}} \mathbb{P}^4 \\
 \downarrow & & \downarrow \text{ } \mathbb{P}^{29}\text{-b'dle} & & \\
 (E, S) & \in & U^{(26)} & \subset & G \times G(2, H^0(O_{\mathbb{P}^4}(2))) \\
 \downarrow & & \downarrow \text{ } G(2, 12)\text{-b'dle} & & \downarrow \\
 E & \in & G(2, 5)^{(6)} & = & \{ \text{lines in } \mathbb{P}^4 \}.
 \end{array}$$

## Analogy of Mumford's example

### Example 2

**Hilb<sup>sc</sup>  $\mathbb{P}^4$**  contains a **generically non-reduced** irreducible component  $W$  whose general member  $C$  satisfies

- ①  $C$  is contained in a **smooth c.i.**  $S = S_{2,2} \subset \mathbb{P}^4$ ,
- ② there exists a **line  $E$**  on  $S$  such that  $C$  belongs to a complete linear system  $\Lambda := |-3K_S + 2E| (\simeq \mathbb{P}^{29})$  on  $S$ , and
- ③  $\dim W = 55$ ,  $h^0(C, N_{C/\mathbb{P}^4}) = 57$ , and  $C$  is of degree 14 and genus 16.

$$\begin{array}{ccccccc}
 C & \in & W^{(55)} & \subset & \text{Hilb}^{\text{sc}} \mathbb{P}^4 \\
 \downarrow & & \downarrow \text{P}^{29}\text{-b'dle} & & \\
 (E, S) & \in & U^{(26)} & \subset & G \times G(2, H^0(O_{\mathbb{P}^4}(2))) \\
 \downarrow & & \downarrow G(2, 12)\text{-b'dle} & & \downarrow \\
 E & \in & G(2, 5)^{(6)} & = & \{\text{lines in } \mathbb{P}^4\}.
 \end{array}$$

## Analogy of Mumford's example

### Example 2

**Hilb<sup>sc</sup> P<sup>4</sup>** contains a **generically non-reduced** irreducible component **W** whose general member **C** satisfies

- ① **C** is contained in a **smooth c.i.**  $S = S_{2,2} \subset \mathbb{P}^4$ ,
- ② there exists a **line E** on  $S$  such that **C** belongs to a complete linear system  $\Lambda := |-3K_S + 2E| (\simeq \mathbb{P}^{29})$  on  $S$ , and
- ③ **dim W = 55**,  $h^0(C, N_{C/\mathbb{P}^4}) = 57$ , and **C** is of degree **14** and genus **16**.

$$\begin{array}{ccccccc}
 C & \in & W^{(55)} & \subset & \text{Hilb}^{\text{sc}} \mathbb{P}^4 \\
 \downarrow & & \downarrow \text{P}^{29}\text{-b'dle} & & \\
 (E, S) & \in & U^{(26)} & \subset & G \times G(2, H^0(O_{\mathbb{P}^4}(2))) \\
 \downarrow & & \downarrow G(2, 12)\text{-b'dle} & & \downarrow \\
 E & \in & G(2, 5)^{(6)} & = & \{\text{lines in } \mathbb{P}^4\}.
 \end{array}$$

## Analogy of Mumford's example


### Example 2

$\text{Hilb}^{sc} \mathbb{P}^4$  contains a **generically non-reduced** irreducible component  $W$  whose general member  $C$  satisfies


- ①  $C$  is contained in a smooth c.i.  $S = S_{2,2} \subset \mathbb{P}^4$ ,
- ② there exists a line  $E$  on  $S$  such that  $C$  belongs to a complete linear system  $\Lambda := |-3K_S + 2E| (\simeq \mathbb{P}^{29})$  on  $S$ , and
- ③  $\dim W = 55$ ,  $h^0(C, N_{C/\mathbb{P}^4}) = 57$ , and  $C$  is of degree 14 and genus 16.

$$\begin{array}{ccccccc}
 C & \in & W^{(55)} & \subset & \text{Hilb}^{sc} \mathbb{P}^4 \\
 \downarrow & & \downarrow \text{$\mathbb{P}^{29}$-b'dle} & & \\
 (E, S) & \in & U^{(26)} & \subset & G \times G(2, H^0(O_{\mathbb{P}^4}(2))) \\
 \downarrow & & \downarrow \text{$G(2, 12)$-b'dle} & & \downarrow \\
 E & \in & G(2, 5)^{(6)} & = & \{\text{lines in } \mathbb{P}^4\}.
 \end{array}$$

## Curves on $S_{2,2} \subset \mathbb{P}^4$ (analogy of Mumford's ex.)



$$\begin{aligned}
 S_{2,2} &= \text{Bl}_{p_1, \dots, p_5} \mathbf{P}^2 \\
 [C] &= (9; 3, 3, 3, 3, 1) \in \text{Pic} S_{2,2} = \mathbf{Z}^6 \\
 d &= 3 \cdot 9 - 3 - 3 - 3 - 3 - 1 = 14 \\
 g &= \frac{(9-1)(9-2)}{2} - 3 - 3 - 3 - 3 = 16
 \end{aligned}$$



## Standard coordinate

Let  $D$  be a divisor on a smooth c.i.  $S_{2,2} \subset \mathbb{P}^4$ , i.e., a quartic del Pezzo surface. Then there exists a suitable blow-up  $\varepsilon : S_{2,2} \rightarrow \mathbb{P}^2$  such that

$$[D] = a - \sum_{i=1}^5 b_i e_i,$$

in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ , where  $l = [\varepsilon^* \mathcal{O}_{\mathbb{P}^2}(1)]$  and  $e_i$ 's are 5 exceptionals, and such that

$$b_1 \geq \dots \geq b_5 \quad \text{and} \quad a \geq b_1 + b_2 + b_3. \tag{1}$$

Then the set of 6-tuples  $(a; b_1, \dots, b_5)$  of integers is called standard coordinate of  $[D]$  in  $\text{Pic } S_{2,2}$ .

## Standard coordinate

Let  $D$  be a divisor on a smooth c.i.  $S_{2,2} \subset \mathbb{P}^4$ , i.e., a quartic del Pezzo surface. Then there exists a suitable blow-up  $\varepsilon : S_{2,2} \rightarrow \mathbb{P}^2$  such that

$$[D] = a - \sum_{i=1}^5 b_i e_i,$$

in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ , where  $l = [\varepsilon^* \mathcal{O}_{\mathbb{P}^2}(1)]$  and  $e_i$ 's are 5 exceptionals, and such that

$$b_1 \geq \dots \geq b_5 \quad \text{and} \quad a \geq b_1 + b_2 + b_3. \tag{1}$$

Then the set of 6-tuples  $(a; b_1, \dots, b_5)$  of integers is called **standard coordinate** of  $[D]$  in  $\text{Pic } S_{2,2}$ .

## Standard coordinate

Let  $D$  be a divisor on a smooth c.i.  $S_{2,2} \subset \mathbb{P}^4$ , i.e., a quartic del Pezzo surface. Then there exists a suitable blow-up  $\varepsilon : S_{2,2} \rightarrow \mathbb{P}^2$  such that

$$[D] = a - \sum_{i=1}^5 b_i e_i,$$

in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ , where  $\mathbf{l} = [\varepsilon^* \mathcal{O}_{\mathbb{P}^2}(1)]$  and  $e_i$ 's are 5 exceptionals, and such that

$$b_1 \geq \dots \geq b_5 \quad \text{and} \quad a \geq b_1 + b_2 + b_3. \tag{1}$$

Then the set of 6-tuples  $(a; b_1, \dots, b_5)$  of integers is called **standard coordinate** of  $[D]$  in  $\text{Pic } S_{2,2}$ .

## Standard coordinate

Let  $D$  be a divisor on a smooth c.i.  $S_{2,2} \subset \mathbb{P}^4$ , i.e., a quartic del Pezzo surface. Then there exists a suitable blow-up  $\varepsilon : S_{2,2} \rightarrow \mathbb{P}^2$  such that

$$[D] = a - \sum_{i=1}^5 b_i e_i,$$

in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ , where  $l = [\varepsilon^* \mathcal{O}_{\mathbb{P}^2}(1)]$  and  $e_i$ 's are 5 exceptionals, and such that

$$b_1 \geq \dots \geq b_5 \quad \text{and} \quad a \geq b_1 + b_2 + b_3. \tag{1}$$

Then the set of 6-tuples  $(a; b_1, \dots, b_5)$  of integers is called **standard coordinate** of  $[D]$  in  $\text{Pic } S_{2,2}$ .

## Standard coordinate

Let  $D$  be a divisor on a smooth c.i.  $S_{2,2} \subset \mathbb{P}^4$ , i.e., a quartic del Pezzo surface. Then there exists a suitable blow-up  $\varepsilon : S_{2,2} \rightarrow \mathbb{P}^2$  such that

$$[D] = a - \sum_{i=1}^5 b_i e_i,$$

in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ , where  $l = [\varepsilon^* \mathcal{O}_{\mathbb{P}^2}(1)]$  and  $e_i$ 's are 5 exceptionals, and such that

$$b_1 \geq \dots \geq b_5 \quad \text{and} \quad a \geq b_1 + b_2 + b_3. \tag{1}$$

Then the set of 6-tuples  $(a; b_1, \dots, b_5)$  of integers is called standard coordinate of  $[D]$  in  $\text{Pic } S_{2,2}$ .

## Standard coordinate

Let  $D$  be a divisor on a smooth c.i.  $S_{2,2} \subset \mathbb{P}^4$ , i.e., a quartic del Pezzo surface. Then there exists a suitable blow-up  $\varepsilon : S_{2,2} \rightarrow \mathbb{P}^2$  such that

$$[D] = a - \sum_{i=1}^5 b_i e_i,$$

in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ , where  $l = [\varepsilon^* \mathcal{O}_{\mathbb{P}^2}(1)]$  and  $e_i$ 's are 5 exceptionals, and such that

$$b_1 \geq \dots \geq b_5 \quad \text{and} \quad a \geq b_1 + b_2 + b_3. \tag{1}$$

Then the set of 6-tuples  $(a; b_1, \dots, b_5)$  of integers is called **standard coordinate** of  $[D]$  in  $\text{Pic } S_{2,2}$ .

## A criterion

### Theorem 9 (N'25)

Let  $C \subset S_{2,2} \subset \mathbb{P}^4$  be a smooth curve of degree  $d \geq 10$  of genus  $g \geq 2d - 12$ , contained in a smooth c.i.  $S_{2,2}$  in  $\mathbb{P}^4$ . Let  $(a; b_1, \dots, b_5)$  be the standard coordinate of  $[C]$  in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ . Then

- ① If  $b_5 \geq 2$ , then  $C$  is unobstructed and stably degenerate.
- ② If  $b_5 = 1$  and  $b_4 \geq 2$ , then  $C$  is obstructed and stably degenerate.
- ③ If  $b_5 = 0$ , then  $C$  is not stably degenerate.

## A criterion

### Theorem 9 (N'25)

Let  $C \subset S_{2,2} \subset \mathbb{P}^4$  be a smooth curve of degree  $d \geq 10$  of genus  $g \geq 2d - 12$ , contained in a smooth c.i.  $S_{2,2}$  in  $\mathbb{P}^4$ . Let  $(a; b_1, \dots, b_5)$  be the standard coordinate of  $[C]$  in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ . Then

- ① If  $b_5 \geq 2$ , then  $C$  is unobstructed and stably degenerate.
- ② If  $b_5 = 1$  and  $b_4 \geq 2$ , then  $C$  is obstructed and stably degenerate.
- ③ If  $b_5 = 0$ , then  $C$  is not stably degenerate.

## A criterion

### Theorem 9 (N'25)

Let  $C \subset S_{2,2} \subset \mathbb{P}^4$  be a smooth curve of degree  $d \geq 10$  of genus  $g \geq 2d - 12$ , contained in a smooth c.i.  $S_{2,2}$  in  $\mathbb{P}^4$ . Let  $(a; b_1, \dots, b_5)$  be the standard coordinate of  $[C]$  in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ . Then

- ① If  $b_5 \geq 2$ , then  $C$  is unobstructed and stably degenerate.
- ② If  $b_5 = 1$  and  $b_4 \geq 2$ , then  $C$  is obstructed and stably degenerate.
- ③ If  $b_5 = 0$ , then  $C$  is not stably degenerate.

## A criterion

### Theorem 9 (N'25)

Let  $C \subset S_{2,2} \subset \mathbb{P}^4$  be a smooth curve of degree  $d \geq 10$  of genus  $g \geq 2d - 12$ , contained in a smooth c.i.  $S_{2,2}$  in  $\mathbb{P}^4$ . Let  $(a; b_1, \dots, b_5)$  be the standard coordinate of  $[C]$  in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ . Then

- ① If  $b_5 \geq 2$ , then  $C$  is unobstructed and stably degenerate.
- ② If  $b_5 = 1$  and  $b_4 \geq 2$ , then  $C$  is obstructed and stably degenerate.
- ③ If  $b_5 = 0$ , then  $C$  is not stably degenerate.

# A criterion

## Theorem 9 (N'25)

Let  $C \subset S_{2,2} \subset \mathbb{P}^4$  be a smooth curve of degree  $d \geq 10$  of genus  $g \geq 2d - 12$ , contained in a smooth c.i.  $S_{2,2}$  in  $\mathbb{P}^4$ . Let  $(a; b_1, \dots, b_5)$  be the standard coordinate of  $[C]$  in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ . Then

- ① If  $b_5 \geq 2$ , then  $C$  is unobstructed and stably degenerate.
- ② If  $b_5 = 1$  and  $b_4 \geq 2$ , then  $C$  is obstructed and stably degenerate.
- ③ If  $b_5 = 0$ , then  $C$  is not stably degenerate.

## A criterion

### Theorem 9 (N'25)

Let  $C \subset S_{2,2} \subset \mathbb{P}^4$  be a smooth curve of degree  $d \geq 10$  of genus  $g \geq 2d - 12$ , contained in a smooth c.i.  $S_{2,2}$  in  $\mathbb{P}^4$ . Let  $(a; b_1, \dots, b_5)$  be the standard coordinate of  $[C]$  in  $\text{Pic } S_{2,2} \simeq \mathbb{Z}^6$ . Then

- ① If  $b_5 \geq 2$ , then  $C$  is unobstructed and stably degenerate.
- ② If  $b_5 = 1$  and  $b_4 \geq 2$ , then  $C$  is obstructed and stably degenerate.
- ③ If  $b_5 = 0$ , then  $C$  is not stably degenerate.

# Examples

Table: curves on  $S_{2,2}$  and stable degeneration

$(d, g)$	$(a; b_1, b_2, b_3, b_4, b_5)$	$W(a; b_1, b_2, b_3, b_4, b_5)$
(14, 16)	(8; 2, 2, 2, 2, 2)	unobstructed and stab.degnerate
(14, 16)	(9; 4, 3, 2, 2, 2)	unobstructed and stab.degnerate
(14, 16)	(9; 3, 3, 3, 3, 1)	obstructed and stab.degnerate
(15, 18)	(9; 4, 2, 2, 2, 2)	unobstructed and stab.degnerate
(15, 18)	(9; 3, 3, 3, 2, 1)	obstructed and stab.degnerate
⋮	⋮	⋮
(18, 24)	(9; 2, 2, 2, 2, 1)	obstructed and stab.degnerate
(18, 24)	(10; 4, 3, 3, 1, 1)	unknown ( $h^1(\mathcal{I}_C(2)) = 2$ )
(18, 24)	(10; 3, 3, 3, 3, 0)	not stab.degnerate ( $h^1(\mathcal{I}_C(2)) = 3$ )
(18, 24)	(11; 6, 3, 2, 2, 2)	unobstructed and stab.degnerate
⋮	⋮	⋮

## further questions

- ① Deformations of curves lying on del Pezzo  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 5$ .
- ② Deformation of degenerate curves on del Pezzo manifold of higher dimension ( $> 3$ ).
- ③ Study the relation to other examples of obstructed curves  $C \subset \mathbb{P}^n$  (or non-reduced components of  $\text{Hilb}^{sc} \mathbb{P}^n$ ).  
[Y. Choi–H. Iliev–S. Kim'24] have recently proved the existence of many non-reduced components of  $\text{Hilb}^{sc} \mathbb{P}^n$  of higher dimensional projective space  $\mathbb{P}^n$  by using ruled surfaces.
- ④ ...

## further questions

- ① Deformations of curves lying on del Pezzo  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 5$ .
- ② Deformation of degenerate curves on del Pezzo manifold of higher dimension ( $> 3$ ).
- ③ Study the relation to other examples of obstructed curves  $C \subset \mathbb{P}^n$  (or non-reduced components of  $\text{Hilb}^{sc} \mathbb{P}^n$ ).  
[Y. Choi–H. Iliev–S. Kim'24] have recently proved the existence of many non-reduced components of  $\text{Hilb}^{sc} \mathbb{P}^n$  of higher dimensional projective space  $\mathbb{P}^n$  by using ruled surfaces.
- ④ ...

## further questions

- ① Deformations of curves lying on del Pezzo  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 5$ .
- ② Deformation of degenerate curves on del Pezzo manifold of higher dimension ( $> 3$ ).
- ③ Study the relation to other examples of obstructed curves  $C \subset \mathbb{P}^n$  (or non-reduced components of  $\mathbf{Hilb}^{sc} \mathbb{P}^n$ ).

[Y. Choi–H. Iliev–S. Kim'24] have recently proved the existence of many non-reduced components of  $\mathbf{Hilb}^{sc} \mathbb{P}^n$  of higher dimensional projective space  $\mathbb{P}^n$  by using ruled surfaces.

4 ...

## further questions

- ① Deformations of curves lying on del Pezzo  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 5$ .
- ② Deformation of degenerate curves on del Pezzo manifold of higher dimension ( $> 3$ ).
- ③ Study the relation to other examples of obstructed curves  $C \subset \mathbb{P}^n$  (or non-reduced components of  $\mathbf{Hilb}^{sc} \mathbb{P}^n$ ).

[Y. Choi–H. Iliev–S. Kim'24] have recently proved the existence of many non-reduced components of  $\mathbf{Hilb}^{sc} \mathbb{P}^n$  of higher dimensional projective space  $\mathbb{P}^n$  by using ruled surfaces.

## further questions

- ① Deformations of curves lying on del Pezzo  $S_n \subset \mathbb{P}^n$  of degree  $n \geq 5$ .
- ② Deformation of degenerate curves on del Pezzo manifold of higher dimension ( $> 3$ ).
- ③ Study the relation to other examples of obstructed curves  $C \subset \mathbb{P}^n$  (or non-reduced components of  $\mathbf{Hilb}^{sc} \mathbb{P}^n$ ).  
[Y. Choi–H. Iliev–S. Kim'24] have recently proved the existence of many non-reduced components of  $\mathbf{Hilb}^{sc} \mathbb{P}^n$  of higher dimensional projective space  $\mathbb{P}^n$  by using ruled surfaces.
- ④ ...

Thank you very much for listening! and Happy birthday to Prof. Youngook Choi.



## References



D. Mumford.

Further pathologies in algebraic geometry.

*Amer. J. Math.*, 84:642–648, 1962.



S. Mukai and H. Nasu.

Obstructions to deforming curves on a 3-fold, I: A generalization of Mumford's example and an application to Hom schemes.

*J. Algebraic Geom.*, 18(4):691–709, 2009.



H. Nasu,

Obstructions to deforming curves on a 3-fold, II: Deformations of degenerate curves on a del Pezzo 3-fold,

*Annales de L'Institut Fourier*, 60(2010), no. 4, 1289–1316.



H. Nasu.

Obstructions to deforming curves on a 3-fold, III: Deformations of curves lying on a **K3** surface.

*Internat. J. Math.*, 28(13):1750099, 30, 2017.



H. Nasu.

Obstructions to deforming space curves lying on a smooth cubic surface.

*Manuscripta Math.*, 172 (2023), pp.31-55.