

# *Approximation of a System of Rational Functional Equations of Three Variables*

**B. V. Senthil Kumar, Hemen Dutta & S. Sabarinathan**

**International Journal of Applied and Computational Mathematics**


ISSN 2349-5103

Volume 5

Number 2

Int. J. Appl. Comput. Math (2019) 5:1-16

DOI 10.1007/s40819-019-0626-y



**Your article is protected by copyright and all rights are held exclusively by Springer Nature India Private Limited. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at [link.springer.com](http://link.springer.com)".**



# Approximation of a System of Rational Functional Equations of Three Variables

B. V. Senthil Kumar<sup>1</sup> · Hemen Dutta<sup>2</sup> · S. Sabarinathan<sup>3</sup>

© Springer Nature India Private Limited 2019

## Abstract

This study aims at substantiating the validity of stability results of a system of rational functional equations involving three variables connected with the Ulam stability theory of functional equations. There are some functional equations identified with real-time occurrences. This system of functional equations is related with the intensive properties of substances such as density and volume.

**Keywords** Hyperreal · Reciprocal function · Bi-reciprocal functional equation · Generalized Hyers–Ulam–Rassias stability

**Mathematics Subject Classification** 39B82 · 39B72

## Introduction

The set of real numbers is extended to a set comprising indeterminable and minuscule numbers, which is called as the set of hyperreal numbers. If there exists a positive integer  $k$  such that for a hyperreal number  $v$ , if  $|v| < k$ , then  $v$  is said to be finite and minuscule if  $|v| < \frac{1}{k}$ . The symbol  ${}^*\mathbb{R}$  is used to denote the set of hyperreals. The conjecture of real numbers is employed in the standard analysis. The study of standard analysis is calculus. In the same manner, the hypothesis of hyperreals are utilized for non-standard analysis (NSA). The hypothesis of NSA was extended by Robinson in 1960's and it was considered as attentive

---

✉ B. V. Senthil Kumar  
 bvsukumarmaths@gmail.com; senthilkumar@nct.edu.om

Hemen Dutta  
 hemen\_dutta08@rediffmail.com

S. Sabarinathan  
 ssabarimaths@gmail.com

<sup>1</sup> Section of Mathematics, Department of Information Technology, Nizwa College of Technology, 611 Nizwa, Oman

<sup>2</sup> Department of Mathematics, Gauhati University, Gawahati, Assam 781 014, India

<sup>3</sup> Department of Mathematics, SRM Institute of Science and Technology, Kattankulathur, Tamil Nadu 603 203, India

fundamentals for intuitions about minuscules. It can be found that there are plenty of applications of hyperreals in many divisions of science, often with an interesting philosophical dimension. Hyperreals could be used to reevaluate the archive of calculus. A remarkable implementation of NSA is in the didactics of calculus. There are many other exertions of NSA in physics that are related to differential and stochastic equations and also in Markov processes, Lévy Brownian motion, and Sturm-Liouville problems. NSA also finds major role in quantum mechanics which includes quantum field theory and Feynman path integrals. For details of many fascinating and interesting features of hyperreals, one can refer to [13,40].

The conjecture of stability of functional equations is pioneered via the query submitted by Ulam [39] and subsequently his problem was responded by Hyers [18]. Eventually Ulam's query was postulated by Rassias [28] and Gavruta [12] in different adaptations. There are many novel, significant, motivating, interesting problems connected with the stability problem of various forms of functional equations and system of functional equations in different spaces and domains (see [1,2,10,14,15,21–23,25,26]).

The reciprocal, quadratic reciprocal and cubic reciprocal functional equations are studied for the first time in [5,31] and [19], respectively.

Several other type of rational form of functional equations and their stability results are discussed in [3,4,6–9,16,17,20,24,30,33,34,37,38].

Moreover, the mathematical formulation of several multiplicative inverse functional equations and the classical investigation of various stability results of multiplicative inverse undecic and duodecic functional equations can be found in [36].

In the recent times, the authors of this paper [35] have dealt with a new rational functional equation involving two variables of the form

$$r(u_1 + u_2, v_1 + v_2) = \frac{\prod_{j=1}^2 \prod_{k=1}^2 r(u_j, v_k)}{\sum_{j=1}^2 \sum_{k=1}^2 \left( \frac{1}{r(u_j, v_k)} \prod_{m=1}^2 \prod_{n=1}^2 r(u_m, v_n) \right)} \quad (1)$$

to accomplish the solution of (1), examine its various stabilities and illustrate its relevance to designing some systems appearing in physics.

Ravi and first author of this paper [32] dealt the following system of bi-reciprocal functional equations

$$\left. \begin{aligned} r(x + u, y) &= \frac{r(x, y)r(u, y)}{r(x, y) + r(u, y)} \\ r(x, y + v) &= \frac{r(x, y)r(x, v)}{r(x, y) + r(x, v)} \end{aligned} \right\} \quad (2)$$

and validated the existence of various stabilities in Fréchet spaces. It is easy to see that  $r(x, y) = \frac{1}{xy}$  is a solution of (2).

In the sequel, we extend the system of bi-reciprocal functional equations (2) of two variables into a system of functional equations involving three variables with the following definition.

**Definition 1** A function  $T_r : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \longrightarrow \mathbb{R}$  is called as tri-reciprocal if  $T_r$  fulfills the ensuing system of functional equations

$$\left. \begin{aligned} T_r(u_1 + u_2, v, w) &= \frac{T_r(u_1, v, w)T_r(u_2, v, w)}{T_r(u_1, v, w) + T_r(u_2, v, w)}, \\ T_r(u, v_1 + v_2, w) &= \frac{T_r(u, v_1, w)T_r(u, v_2, w)}{T_r(u, v_1, w) + T_r(u, v_2, w)}, \\ T_r(u, v, w_1 + w_2) &= \frac{T_r(u, v, w_1)T_r(u, v, w_2)}{T_r(u, v, w_1) + T_r(u, v, w_2)}. \end{aligned} \right\} \quad (3)$$

It can be easily substantiated that the rational function  $T_r(u, v, w) = \frac{c}{uvw}$  is a solution of the system (3), where  $c$  is a constant.

In this paper, we prove the existence of the stability results of the system of equations (3) in the setting of hyperreals, associated with the stability theory originated from Ulam, Hyers, Rassias and Gavruta. We also discuss an application of the solution of system of equations (3) with the basic properties of an object.

For the convenience of manipulation, let us symbolize for a given mapping  $T_r : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$ , the difference operators  $\Delta_1, \Delta_2, \Delta_3 : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$  by

$$\begin{aligned} \Delta_1 T_r(u_1, u_2, v, w) &= T_r(u_1 + u_2, v, w) - \frac{T_r(u_1, v, w)T_r(u_2, v, w)}{T_r(u_1, v, w) + T_r(u_2, v, w)}, \\ \Delta_2 T_r(u, v_1, v_2, w) &= T_r(u, v_1 + v_2, w) - \frac{T_r(u, v_1, w)T_r(u, v_2, w)}{T_r(u, v_1, w) + T_r(u, v_2, w)}, \\ \Delta_3 T_r(u, v, w_1, w_2) &= T_r(u, v, w_1 + w_2) - \frac{T_r(u, v_1, w)T_r(u, v_2, w)}{T_r(u, v_1, w) + T_r(u, v_2, w)} \end{aligned}$$

for all  $u_1, u_2, v_1, v_2, w_1, w_2 \in {}^*\mathbb{R}$ .

## Stability of the System of Equation (3)

In the sequel, we validate the existence of stabilities of the system of equations (3) pertinent to Hyers, Rassias and Gavruta by considering the set of hyperreals as domain. The proceeding result includes the investigation of the stability of system of equations (3) relevant to Gavruta's repercussion.

**Theorem 1** Assume that the functions  $P, Q, R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow [0, \infty)$  satisfy the following conditions:

$$\left. \begin{aligned} \sum_{j=0}^{\infty} 8^j P(2^j u, 2^j v, 2^j w) &< \infty, \\ \sum_{j=0}^{\infty} 8^j Q(2^{j+1} u, 2^j v, 2^j w) &< \infty, \\ \sum_{j=0}^{\infty} 8^j R(2^{j+1} u, 2^{j+1} v, 2^j w) &< \infty \end{aligned} \right\} \quad (4)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Also, let us presume that for a given positive real number  $\epsilon$ , the above functions satisfy the ensuing inequalities:

$$\left. \begin{aligned} \sum_{j=0}^{\infty} 8^j P(2^j u, 2^j v, 2^j w) &< \frac{\epsilon}{6}, \\ \sum_{j=0}^{\infty} 8^j Q(2^{j+1} u, 2^j v, 2^j w) &< \frac{\epsilon}{12} \\ \sum_{j=0}^{\infty} 8^j R(2^{j+1} u, 2^{j+1} v, 2^j w) &< \frac{\epsilon}{24} \end{aligned} \right\} \quad (5)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Suppose a function  $T_r^* : \mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \longrightarrow \mathbb{R}$  fulfills

$$|\Delta_1 T_r(u_1, u_2, v, w)| \leq P(u_1, u_2, v, w) \quad (6)$$

$$|\Delta_2 T_r(u, v_1, v_2, w)| \leq Q(u, v_1, v_2, w) \quad (7)$$

$$|\Delta_3 T_r(u, v, w_1, w_2)| \leq R(u, v, w_1, w_2) \quad (8)$$

for all  $u_1, u_2, v_1, v_2, w_1, w_2 \in {}^*\mathbb{R}$ . Then, there exists a unique tri-reciprocal function  $T_R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \longrightarrow \mathbb{R}$  which satisfies (3) and

$$\begin{aligned} &|T_r(u, v, w) - T_R(u, v, w)| \\ &\leq 2 \sum_{j=0}^{\infty} 8^j P(2^j u, 2^j v, 2^j w) + 4 \sum_{j=0}^{\infty} 8^j Q(2^{j+1} u, 2^j v, 2^j w) \\ &\quad + 8 \sum_{j=0}^{\infty} 8^j R(2^{j+1} u, 2^{j+1} v, 2^j w) \\ &\leq \epsilon \end{aligned} \quad (9)$$

for all  $u, v, w \in {}^*\mathbb{R}$ .

**Proof** First, let us substitute  $u_1 = u_2 = u$  in (6) and then multiplying by 2 on both sides to obtain

$$|2T_r(2u, v, w) - T_r(u, v, w)| \leq 2P(u, u, v, w) \quad (10)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Now, letting  $v_1 = v_2 = v$  in (7) and then multiplying by 2 on both sides, we obtain

$$|2T_r(u, 2v, w) - T_r(u, v, w)| \leq 2Q(u, v, v, w) \quad (11)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Now, plugging  $(w_1, w_2)$  by  $(w, w)$  in (8) and then multiplying by 2 on both sides, we get

$$|2T_r(u, v, 2w) - T_r(u, v, w)| \leq 2R(u, v, w, w) \quad (12)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Replacing  $v$  by  $2v$  in (12) and then multiplying by 2 on both sides, yields

$$|4T_r(u, 2v, 2w) - 2T_r(u, 2v, w)| \leq 4R(u, 2v, w, w) \quad (13)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . By employing triangle inequality law on (11) and (13), we find

$$|4T_r(u, 2v, 2w) - T_r(u, v, w)| \leq 2Q(u, v, v, w) + 4R(u, 2v, w, w) \quad (14)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Now, swapping  $u$  by  $2u$  in (14) and then multiplying by 2, we obtain

$$|8T_r(2u, 2v, 2w) - 2T_r(2u, v, w)| \leq 4Q(2u, v, v, w) + 8R(2u, 2v, w, w) \quad (15)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Now, again in lieu of triangle inequality law on (10) and (15), we lead to

$$\begin{aligned} & |8T_r(2u, 2v, 2w) - T_r(u, v, w)| \\ & \leq 2P(u, u, v, w) + 4Q(2u, v, v, w) + 8R(2u, 2v, w, w) \end{aligned} \quad (16)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Now, plugging  $(u, v, w)$  into  $(2u, 2v, 2w)$  in (16) and then multiplying by 8 on both sides, we obtain

$$\begin{aligned} & |8^2T_r(4u, 4v, 4w) - 8T_r(2u, 2v, 2w)| \\ & \leq 16P(2u, 2u, 2v, 2w) + 32Q(4u, 2v, 2v, 2w) + 64R(4u, 4v, 2w, 2w) \end{aligned} \quad (17)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Further by the application of triangle inequality law on (16) and (17), we get

$$\begin{aligned} & |8^2T_r(4u, 4v, 4w) - T_r(u, v, w)| \\ & \leq 2P(u, u, v, w) + 16P(2u, 2u, 2v, 2w) + 4Q(2u, v, v, w) + 32Q(4u, 2v, 2v, 2w) \\ & \quad + 8R(2u, 2v, w, w) + 64R(4u, 4v, 2w, 2w) \\ & \leq 2 \sum_{j=0}^1 8^j P(2^j u, 2^j u, 2^j v, 2^j w) + 4 \sum_{j=0}^1 8^j Q(2^{j+1} u, 2^j v, 2^j v, 2^j w) \\ & \quad + 8 \sum_{j=0}^1 8^j R(2^{j+1} u, 2^{j+1} v, 2^j w, 2^j w) \end{aligned}$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Continuing further in this fashion and utilizing induction arguments on the above inequality, we conclude that

$$\begin{aligned} & |8^n T_r(4u, 4v, 4w) - T_r(u, v, w)| \\ & \leq 2 \sum_{j=0}^{n-1} 8^j P(2^j u, 2^j u, 2^j v, 2^j w) + 4 \sum_{j=0}^{n-1} 8^j Q(2^{j+1} u, 2^j v, 2^j v, 2^j w) \\ & \quad + 8 \sum_{j=0}^{n-1} 8^j R(2^{j+1} u, 2^{j+1} v, 2^j w, 2^j w) \\ & \leq 2 \sum_{j=0}^{\infty} 8^j P(2^j u, 2^j u, 2^j v, 2^j w) + 4 \sum_{j=0}^{\infty} 8^j Q(2^{j+1} u, 2^j v, 2^j v, 2^j w) \\ & \quad + 8 \sum_{j=0}^{\infty} 8^j R(2^{j+1} u, 2^{j+1} v, 2^j w, 2^j w) \end{aligned} \quad (18)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Next is to prove the sequence  $\{8^n r(2^n u, 2^n v, 2^n w)\}$  is convergent. For this, let us plug  $(u, v, w)$  into  $(2^k u, 2^k v, 2^k w)$  in (18) and then multiply by  $8^k$ , then we find for any  $n, k > 0$ ,

$$\begin{aligned} & |8^{n+k} T_r(2^{n+k} u, 2^{n+k} v, 2^{n+k} w) - 8^k T_r(2^k u, 2^k v, 2^k w)| \\ & = 8^k |8^n T_r(2^{n+k} u, 2^{n+k} v, 2^{n+k} w) - T_r(2^k u, 2^k v, 2^k w)| \end{aligned}$$

$$\begin{aligned}
 &\leq 2 \sum_{j=0}^{\infty} 8^{k+j} T_r P \left( 2^{k+j} u, 2^{k+j} u, 2^{k+j} v, 2^{k+j} w \right) \\
 &\quad + 4 \sum_{j=0}^{\infty} 8^{k+j} Q \left( 2^{k+j+1} u, 2^{k+j} u, 2^{k+j} v, 2^{k+j} w \right) \\
 &\quad + 8 \sum_{j=0}^{\infty} 8^{k+j} R \left( 2^{k+j+1} u, 2^{k+j+1} v, 2^{k+j} w, 2^{k+j} w \right) \\
 &\longrightarrow 0 \text{ as } k \longrightarrow \infty.
 \end{aligned}$$

From the above result, we conclude that the sequence  $\{8^n r(2^n u, 2^n v, 2^n w)\}$  is Cauchy. Since  $\mathbb{R}$  is complete, it is obvious that there exists a function  $T_R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \longrightarrow \mathbb{R}$  defined by the existence of the limit  $T_R(u, v, w) = \lim_{n \rightarrow \infty} 8^n T_r(2^n u, 2^n v, 2^n w)$  for all  $u, v, w \in X$ . It is clear that (9) is valid for all  $u, v, w \in {}^*\mathbb{R}$  by letting  $n \longrightarrow \infty$  in (18). Now, let us show that  $T_R$  satisfies (3). In fact, switching  $(u, v, w)$  to  $(2^n u, 2^n v, 2^n w)$  and then multiplying by  $8^n$  in (6), (7) and (8), respectively, we lead to

$$\begin{aligned}
 |\Delta_1 T_R(u_1, u_2, v, w)| &= \lim_{n \rightarrow \infty} 8^n \Delta_1 T_r(2^n u_1, 2^n u_2, 2^n v, 2^n w) \\
 &\leq \lim_{n \rightarrow \infty} 8^n P(2^n u_1, 2^n u_2, 2^n v, 2^n w) = 0, \\
 |\Delta_2 T_R(u, v_1, v_2, w)| &= \lim_{n \rightarrow \infty} 8^n \Delta_2 T_r(2^n u, 2^n v_1, 2^n v_2, 2^n w) \\
 &\leq \lim_{n \rightarrow \infty} 8^n P(2^n u, 2^n v_1, 2^n v_2, 2^n w) = 0, \\
 |\Delta_3 T_R(u, v, w_1, w_2)| &= \lim_{n \rightarrow \infty} 8^n \Delta_3 T_r(2^n u, 2^n v, 2^n w_1, 2^n w_2) \\
 &\leq \lim_{n \rightarrow \infty} 8^n P(2^n u, 2^n v, 2^n w_1, 2^n w_2) = 0
 \end{aligned}$$

for all  $u_1, u_2, v_1, v_2, w_1, w_2 \in {}^*\mathbb{R}$ , which shows that  $T_R$  is tri-reciprocal. In order to show the uniqueness of  $T_R$ , let us presume that  $T'_R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \longrightarrow \mathbb{R}$  is another tri-reciprocal function satisfying (3) and (9), then

$$\begin{aligned}
 &|T_R(u, v, w) - T'_R(u, v, w)| \\
 &= 8^n |T_R(2^n u, 2^n v, 2^n w) - T'_R(2^n u, 2^n v, 2^n w)| \\
 &= 8^n \left\{ |T_R(2^n u, 2^n v, 2^n w) - T_r(2^n u, 2^n v, 2^n w)| \right. \\
 &\quad \left. + |T_r(2^n u, 2^n v, 2^n w) - T'_R(2^n u, 2^n v, 2^n w)| \right\} \\
 &\leq 4 \sum_{j=0}^{\infty} 8^{n+j} P(2^{n+j} u, 2^{n+j} u, 2^{n+j} v, 2^{n+j} w) \\
 &\quad + 8 \sum_{j=0}^{\infty} 8^{n+j} Q(2^{n+j+1} u, 2^{n+j} v, 2^{n+j} v, 2^{n+j} w) \\
 &\quad + 16 \sum_{j=0}^{\infty} 8^{n+j} R(2^{n+j+1} u, 2^{n+j+1} v, 2^{n+j} w, 2^{n+j} w)
 \end{aligned}$$

for all  $u, v, w \in {}^*\mathbb{R}$ . This implies that  $T_R(u, v, w) = T'_R(u, v, w)$  for all  $u, v, w \in {}^*\mathbb{R}$ . Hence, we conclude that  $T_R$  is unique, which completes the proof.  $\square$



In the upcoming result, we illustrate the validity of the existence of stability of the system of equations (3) incorporated with addition of exponents of norms.

**Corollary 1** Consider a fixed constant  $\theta_1 > 0$ . Suppose  $\alpha < -3$  is a real number. Assume a function  $T_r : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \longrightarrow \mathbb{R}$  satisfies the ensuing inequalities

$$\left. \begin{aligned} |\Delta_1 T_r(u_1, u_2, v, w)| &\leq \theta_1 (\|u_1\|^\alpha + \|u_2\|^\alpha + \|v\|^\alpha + \|w\|^\alpha), \\ |\Delta_2 T_r(u, v_1, v_2, w)| &\leq \theta_1 (\|u\|^\alpha + \|v_1\|^\alpha + \|v_2\|^\alpha + \|w\|^\alpha), \\ |\Delta_3 T_r(u, v, w_1, w_2)| &\leq \theta_1 (\|u\|^\alpha + \|v\|^\alpha + \|w_1\|^\alpha + \|w_2\|^\alpha), \end{aligned} \right\} \quad (19)$$

for all  $u_1, u_2, v_2, w_1, w_2 \in {}^*\mathbb{R}$ . Then, there exists a unique tri-reciprocal function  $T_R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \longrightarrow \mathbb{R}$  which satisfies (3) and

$$\begin{aligned} &|T_r(u, v, w) - T_R(u, v, w)| \\ &\leq \left( \frac{2\theta_1}{1 - 2^{\alpha+3}} \right) \{ (2 + 6 \cdot 2^\alpha) |u|^\alpha + (5 + 4 \cdot 2^\alpha) |v|^\alpha + 11 |w|^\alpha \} \end{aligned}$$

for all  $u, v, w \in {}^*\mathbb{R}$ .

**Proof** Considering

$$P(u_1, u_2, v, w) = \theta_1 (\|u_1\|^\alpha + \|u_2\|^\alpha + \|v\|^\alpha + \|w\|^\alpha),$$

$$Q(u, v_1, v_2, w) = \theta_1 (\|u\|^\alpha + \|v_1\|^\alpha + \|v_2\|^\alpha + \|w\|^\alpha),$$

and

$$R(u, v, w_1, w_2) = \theta_1 (\|u\|^\alpha + \|v\|^\alpha + \|w_1\|^\alpha + \|w_2\|^\alpha)$$

for all  $u, u_1, u_2, v, v_1, v_2, w, w_1, w_2 \in {}^*\mathbb{R}$  in Theorem 1, we get

$$\begin{aligned} &|T_r(u, v, w) - T_R(u, v, w)| \\ &\leq 2\theta_1 \sum_{j=0}^{\infty} 8^j \left( |2^j u|^\alpha + |2^j v|^\alpha + |2^j w|^\alpha \right) \\ &\quad + 4\theta_1 \sum_{j=0}^{\infty} 8^j \left( |2^{j+1} u|^\alpha + |2^j v|^\alpha + |2^j v|^\alpha + |2^j w|^\alpha \right) \\ &\quad + 8\theta_1 \sum_{j=0}^{\infty} 8^j \left( |2^{j+1} u|^\alpha + |2^{j+1} v|^\alpha + |2^j w|^\alpha + |2^j w|^\alpha \right) \\ &\leq 2\theta_1 \sum_{j=0}^{\infty} \left( 2 \cdot 2^{(\alpha+3)j} |u|^\alpha + 2^{(\alpha+3)j} |v|^\alpha + 2^{(\alpha+3)j} |w|^\alpha \right) \\ &\quad + 4\theta_1 \sum_{j=0}^{\infty} \left( 2^\alpha \cdot 2^{(\alpha+3)j} |u|^\alpha + 2 \cdot 2^{(\alpha+3)j} |v|^\alpha + 2^{(\alpha+3)j} |w|^\alpha \right) \\ &\quad + 8\theta_1 \sum_{j=0}^{\infty} \left( 2^\alpha \cdot 2^{(\alpha+3)j} |u|^\alpha + 2^\alpha \cdot 2^{(\alpha+3)j} |v|^\alpha + 2 \cdot 2^{(\alpha+3)j} |w|^\alpha \right) \\ &\leq \left( \frac{2\theta_1}{1 - 2^{\alpha+3}} \right) [(2 + 6 \cdot 2^\alpha) |u|^\alpha + (5 + 4 \cdot 2^\alpha) |v|^\alpha + 11 |w|^\alpha] \end{aligned}$$

for all  $u, v, w \in {}^*\mathbb{R}$ . □

The following corollaries are the stability results of system of equations (3) associated with UGR stability (involving product of powers of norms) [27] and J. Rassias stability [29] (involving mixed product-sum of powers of norms). Since the arguments of arriving at the desired results are similar to Corollary 1, we provide only the results.

**Corollary 2** Let  $\theta_2 > 0$  be a fixed constant and  $\alpha < -3$  is a real number. Let a mapping  $T_r : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$  satisfies the following inequalities

$$\left. \begin{aligned} |\Delta_1 T_r(u_1, u_2, v, w)| &\leq \theta_3 |u_1|^{\frac{\alpha}{6}} |u_2|^{\frac{\alpha}{6}} |v|^{\frac{\alpha}{3}} |w|^{\frac{\alpha}{3}}, \\ |\Delta_2 T_r(u, v_1, v_2, w)| &\leq \theta_3 |u|^{\frac{\alpha}{3}} |v_1|^{\frac{\alpha}{6}} |v_2|^{\frac{\alpha}{6}} |w|^{\frac{\alpha}{3}}, \\ |\Delta_3 T_r(u, v, w_1, w_2)| &\leq \theta_3 |u|^{\frac{\alpha}{3}} |v|^{\frac{\alpha}{3}} |w_1|^{\frac{\alpha}{6}} |w_2|^{\frac{\alpha}{6}}, \end{aligned} \right\} \quad (20)$$

for all  $u_1, u_2, v_2, w_1, w_2 \in {}^*\mathbb{R}$ . Then, a unique tri-reciprocal function  $T_R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$  exists and satisfies (3) such that

$$|T_r(u, v, w) - T_R(u, v, w)| \leq \left( \frac{2\theta_2}{1 - 2^{\alpha+3}} \right) \left( 1 + 2 \cdot 2^{\frac{\alpha}{3}} + 4 \cdot 2^{\frac{\alpha}{3}} \right) |u|^{\frac{\alpha}{3}} |v|^{\frac{\alpha}{3}} |w|^{\frac{\alpha}{3}}$$

for all  $u, v, w \in {}^*\mathbb{R}$ .

**Corollary 3** Assume that  $\theta_3 > 0$  is a fixed constant and  $\alpha < -3$  is a real number. If a function  $T_r : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$  satisfies the ensuing inequalities

$$\left. \begin{aligned} |\Delta_1 T_r(u_1, u_2, v, w)| &\leq \theta_3 \left( |u_1|^{\frac{\alpha}{6}} |u_2|^{\frac{\alpha}{6}} |v|^{\frac{\alpha}{3}} |w|^{\frac{\alpha}{3}} + (|u_1|^\alpha + |u_2|^\alpha + |v|^\alpha + |w|^\alpha) \right), \\ |\Delta_2 T_r(u, v_1, v_2, w)| &\leq \theta_3 \left( |u|^{\frac{\alpha}{3}} |v_1|^{\frac{\alpha}{6}} |v_2|^{\frac{\alpha}{6}} |w|^{\frac{\alpha}{3}} + (|u|^\alpha + |v_1|^\alpha + |v_2|^\alpha + |w|^\alpha) \right), \\ |\Delta_3 T_r(u, v, w_1, w_2)| &\leq \theta_3 \left( |u|^{\frac{\alpha}{3}} |v|^{\frac{\alpha}{3}} |w_1|^{\frac{\alpha}{6}} |w_2|^{\frac{\alpha}{6}} + (|u|^\alpha + |v|^\alpha + |w_1|^\alpha + |w_2|^\alpha) \right), \end{aligned} \right\} \quad (21)$$

for all  $u_1, u_2, v_2, w_1, w_2 \in {}^*\mathbb{R}$ , then a unique tri-reciprocal function  $T_R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$  exists and satisfies (3) such that

$$\begin{aligned} |T_r(u, v, w) - T_R(u, v, w)| &\leq \left( \frac{2\theta_3}{1 - 2^{\alpha+3}} \right) \left\{ \left( 1 + 2 \cdot 2^{\frac{\alpha}{6}} + 4 \cdot 2^{\frac{2\alpha}{3}} \right) |u|^{\frac{\alpha}{3}} |v|^{\frac{\alpha}{3}} |w|^{\frac{\alpha}{3}} \right. \\ &\quad \left. + 2 \left( 1 + 2^\alpha + 2^{\alpha+1} \right) |u|^\alpha + \left( 5 + 2^{\alpha+3} \right) |v|^\alpha + 7 |w|^\alpha \right\} \end{aligned}$$

for all  $u, v, w \in {}^*\mathbb{R}$ .

The subsequent theorem is dual result of Theorem 1. Eventhough, the proof of this theorem is analogous to Theorem 1, for the sake of comprehensiveness, we provide the key part of the proof.

**Theorem 2** Assume that the functions  $P, Q, R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \longrightarrow [0, \infty)$  satisfy the following conditions:

$$\left. \begin{aligned} \sum_{j=0}^{\infty} \frac{1}{8^j} P\left(\frac{u}{2^{j+1}}, \frac{u}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) &< \infty, \\ \sum_{j=0}^{\infty} \frac{1}{8^j} Q\left(\frac{u}{2^j}, \frac{v}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) &< \infty, \\ \sum_{j=0}^{\infty} \frac{1}{8^j} R\left(\frac{u}{2^j}, \frac{v}{2^j}, \frac{w}{2^{j+1}}, \frac{w}{2^{j+1}}\right) &< \infty \end{aligned} \right\} \quad (22)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Let us assume that for a given positive real number  $\epsilon$ , the above functions satisfy the following inequalities:

$$\left. \begin{aligned} \sum_{j=0}^{\infty} \frac{1}{8^j} P\left(\frac{u}{2^{j+1}}, \frac{u}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) &< \frac{4\epsilon}{3}, \\ \sum_{j=0}^{\infty} \frac{1}{8^j} Q\left(\frac{u}{2^j}, \frac{v}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) &< \frac{2\epsilon}{3}, \\ \sum_{j=0}^{\infty} \frac{1}{8^j} R\left(\frac{u}{2^j}, \frac{v}{2^j}, \frac{w}{2^{j+1}}, \frac{w}{2^{j+1}}\right) &< \frac{\epsilon}{3} \end{aligned} \right\} \quad (23)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Suppose a function  $T_r : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \longrightarrow \mathbb{R}$  satisfies (6), (7) and (8) for all  $u, u_1, u_2, v, v_1, v_2, w, w_1, w_2 \in {}^*\mathbb{R}$ . Then, there exists a unique tri-reciprocal function  $T_R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \longrightarrow \mathbb{R}$  which satisfies (3) and

$$\begin{aligned} &|T_r(u, v, w) - T_R(u, v, w)| \\ &\leq \frac{1}{4} \sum_{j=0}^{\infty} \frac{1}{8^j} P\left(\frac{u}{2^{j+1}}, \frac{u}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) + \frac{1}{2} \sum_{j=0}^{\infty} \frac{1}{8^j} Q\left(\frac{u}{2^j}, \frac{v}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) \\ &\quad + \sum_{j=0}^{\infty} \frac{1}{8^j} R\left(\frac{u}{2^j}, \frac{v}{2^j}, \frac{w}{2^{j+1}}, \frac{w}{2^{j+1}}\right) \\ &\leq \epsilon \end{aligned} \quad (24)$$

for all  $u, v, w \in {}^*\mathbb{R}$ .

**Proof** First, let us take  $(u_1, u_2, v, w)$  as  $(\frac{u}{2}, \frac{u}{2}, \frac{v}{2}, \frac{w}{2})$  in (6) and then dividing by 4, we get

$$\left| \frac{1}{4} T_r\left(u, \frac{v}{2}, \frac{w}{2}\right) - \frac{1}{8} T_r\left(\frac{u}{2}, \frac{v}{2}, \frac{w}{2}\right) \right| \leq \frac{1}{4} P\left(\frac{u}{2}, \frac{u}{2}, \frac{v}{2}, \frac{w}{2}\right) \quad (25)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Now, plugging  $(u, v_1, v_2, w)$  into  $(u, \frac{v}{2}, \frac{v}{2}, \frac{w}{2})$  in (7) and then dividing by 2, we obtain

$$\left| \frac{1}{2} T_r\left(u, v, \frac{w}{2}\right) - \frac{1}{4} T_r\left(u, \frac{v}{2}, \frac{w}{2}\right) \right| \leq \frac{1}{2} Q\left(u, \frac{v}{2}, \frac{v}{2}, \frac{w}{2}\right) \quad (26)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Swapping  $(u, v, w_1, w_2)$  into  $(u, v, \frac{w}{2}, \frac{w}{2})$  in (8), we have

$$\left| T_r(u, v, w) - \frac{1}{2} T_r\left(u, v, \frac{w}{2}, \frac{w}{2}\right) \right| \leq R\left(u, v, \frac{w}{2}, \frac{w}{2}\right) \quad (27)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . By employing triangle inequality law on (26) and (27), we find

$$\left| T_r(u, v, w) - \frac{1}{4} T_r\left(u, \frac{v}{2}, \frac{v}{2}, \frac{w}{2}\right) \right| \leq \frac{1}{2} Q\left(u, \frac{v}{2}, \frac{v}{2}, \frac{w}{2}\right) + R\left(u, v, \frac{w}{2}, \frac{w}{2}\right) \quad (28)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Once again, utilizing triangle inequality law on (25) and (28), we lead to

$$\begin{aligned} & \left| T_r(u, v, w) - \frac{1}{8} T_r\left(\frac{u}{2}, \frac{v}{2}, \frac{w}{2}\right) \right| \\ & \leq \frac{1}{4} P\left(\frac{u}{2}, \frac{u}{2}, \frac{v}{2}, \frac{w}{2}\right) + \frac{1}{2} Q\left(u, \frac{v}{2}, \frac{v}{2}, \frac{w}{2}\right) + R\left(u, v, \frac{w}{2}, \frac{w}{2}\right) \end{aligned} \quad (29)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Now, swapping  $(u, v, w)$  into  $(\frac{u}{2}, \frac{v}{2}, \frac{w}{2})$  in (29) and then dividing by 8, we obtain

$$\begin{aligned} & \left| \frac{1}{8} T_r\left(\frac{u}{2}, \frac{v}{2}, \frac{w}{2}\right) - \frac{1}{8^2} T_r\left(\frac{u}{2^2}, \frac{v}{2^2}, \frac{w}{2^2}\right) \right| \\ & \leq \frac{1}{32} P\left(\frac{u}{2^2}, \frac{u}{2^2}, \frac{v}{2^2}, \frac{w}{2^2}\right) + \frac{1}{16} Q\left(\frac{u}{2}, \frac{v}{2^2}, \frac{v}{2^2}, \frac{w}{2^2}\right) + \frac{1}{8} R\left(\frac{u}{2}, \frac{v}{2}, \frac{w}{2^2}, \frac{w}{2^2}\right) \end{aligned} \quad (30)$$

for all  $u, v, w \in {}^*\mathbb{R}$ . In lieu of triangle inequality law on (29) and (30), one finds

$$\begin{aligned} & \left| \frac{1}{8} T_r\left(\frac{u}{2}, \frac{v}{2}, \frac{w}{2}\right) - \frac{1}{8^2} T_r\left(\frac{u}{2^2}, \frac{v}{2^2}, \frac{w}{2^2}\right) \right| \\ & \leq \frac{1}{4} \sum_{j=0}^1 \frac{1}{8^j} P\left(\frac{u}{2^{j+1}}, \frac{u}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) + \frac{1}{2} \sum_{j=0}^1 \frac{1}{8^j} Q\left(\frac{u}{2^j}, \frac{v}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) \\ & \quad + \sum_{j=0}^1 \frac{1}{8^j} R\left(\frac{u}{2^j}, \frac{v}{2^j}, \frac{w}{2^{j+1}}, \frac{w}{2^{j+1}}\right) \end{aligned}$$

for all  $u, v, w \in {}^*\mathbb{R}$ . Continuing further in this fashion and making use of induction arguments on a positive integer  $n$ , we have

$$\begin{aligned} & \left| \frac{1}{8} T_r\left(\frac{u}{2}, \frac{v}{2}, \frac{w}{2}\right) - \frac{1}{8^n} T_r\left(\frac{u}{2^n}, \frac{v}{2^n}, \frac{w}{2^n}\right) \right| \\ & \leq \frac{1}{4} \sum_{j=0}^{n-1} \frac{1}{8^j} P\left(\frac{u}{2^{j+1}}, \frac{u}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) + \frac{1}{2} \sum_{j=0}^{n-1} \frac{1}{8^j} Q\left(\frac{u}{2^j}, \frac{v}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) \\ & \quad + \sum_{j=0}^{n-1} \frac{1}{8^j} R\left(\frac{u}{2^j}, \frac{v}{2^j}, \frac{w}{2^{j+1}}, \frac{w}{2^{j+1}}\right) \\ & \leq \frac{1}{4} \sum_{j=0}^{\infty} \frac{1}{8^j} P\left(\frac{u}{2^{j+1}}, \frac{u}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) + \frac{1}{2} \sum_{j=0}^{\infty} \frac{1}{8^j} Q\left(\frac{u}{2^j}, \frac{v}{2^{j+1}}, \frac{v}{2^{j+1}}, \frac{w}{2^{j+1}}\right) \end{aligned}$$

$$+ \sum_{j=0}^{\infty} \frac{1}{8^j} R\left(\frac{u}{2^j}, \frac{v}{2^j}, \frac{w}{2^{j+1}}, \frac{w}{2^{j+1}}\right) \\ \leq \epsilon$$

for all  $u, v, w \in {}^*\mathbb{R}$ . The enduring part of the proof is achieved akin to Theorem 1.  $\square$

The following result comprises the validity of the stability result of the system of equations (3) concerning the sum of powers of norms, product of powers of norms and mixed product-sum of powers of norms. The following corollaries directly follow from Theorem 2 and hence we omit the proof.

**Corollary 4** Consider a fixed constant  $\theta_1 > 0$ . Suppose  $\alpha > -3$  is a real number. Assume a function  $T_r : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$  fulfills the inequalities in (19). Then, a unique tri-reciprocal function  $T_R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$  exists and satisfies (3) such that

$$|T_r(u, v, w) - T_R(u, v, w)| \\ \leq \left(\frac{2^{\alpha+3}\theta_1}{2^{\alpha+3}-1}\right) \left[\left(\frac{3 \cdot 2^\alpha + 4}{2^\alpha + 1}\right) |u|^\alpha + \left(\frac{2^\alpha + 5}{4 \cdot 2^\alpha}\right) |v|^\alpha + \left(\frac{14}{4 \cdot 2^{\alpha+1}}\right) |w|^\alpha\right]$$

for all  $u, v, w \in {}^*\mathbb{R}$ .

**Proof** The proof is similar to that of Corollary 1.  $\square$

**Corollary 5** Let  $\theta_2 > 0$  be a fixed constant and  $\alpha < -3$  is a real number. Let a mapping  $T_r : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$  satisfies the inequalities (20). Then, a unique tri-reciprocal function  $T_R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$  exists and satisfies (3) such that

$$|T_r(u, v, w) - T_R(u, v, w)| \leq \frac{8\theta_2}{2^\alpha(1-2^{\alpha+3})} \left(1 + 2^{\frac{\alpha}{3}} + 2^{\frac{5\alpha}{3}}\right) |u|^{\frac{\alpha}{3}} |v|^{\frac{\alpha}{3}} |w|^{\frac{\alpha}{3}}$$

for all  $u, v, w \in {}^*\mathbb{R}$ .

**Corollary 6** Assume that  $\theta_3 > 0$  is a fixed constant and  $\alpha < -3$  is a real number. If a function  $T_r : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$  satisfies the inequalities (21), then a unique tri-reciprocal function  $T_R : {}^*\mathbb{R} \times {}^*\mathbb{R} \times {}^*\mathbb{R} \rightarrow \mathbb{R}$  exists and satisfies (3) such that

$$|T_r(u, v, w) - T_R(u, v, w)| \\ \leq \left(\frac{\theta_3}{2^{\alpha+3}-1}\right) \left\{ \left(2 + 12 \cdot 2^{\frac{2\alpha}{3}}\right) |u|^{\frac{\alpha}{3}} |v|^{\frac{\alpha}{3}} |w|^{\frac{\alpha}{3}} \right. \\ \left. + \left(4 + 4^\alpha + 2^{\alpha+3}\right) |u|^\alpha + \left(10 + 2^{\alpha+3}\right) |v|^\alpha + 22 |w|^\alpha \right\}$$

for all  $u, v, w \in {}^*\mathbb{R}$ .

## Counter-Examples

In this section, employing the notion of the famous counter-example presented in [11], we illustrate some counter-examples that the system of equations (3) is not stable for  $\alpha = -3$  in Corollaries 1, 3, 4 and 5.

Consider the function

$$\varphi(u, v, w) = \begin{cases} k(|u|^{-3} + |v|^{-3} + |w|^{-3}), & \text{for } u, v, w \in (0, \infty) \\ k, & \text{otherwise} \end{cases} \quad (31)$$

where  $\varphi : \mathbb{R}^* \times \mathbb{R}^* \times \mathbb{R}^* \longrightarrow \mathbb{R}$ . Let  $T_r : \mathbb{R}^* \times \mathbb{R}^* \times \mathbb{R}^* \longrightarrow \mathbb{R}$  be defined by

$$T_r(u, v, w) = \sum_{n=0}^{\infty} 8^{-n} \varphi(2^{-n}u, 2^{-n}v, 2^{-n}w) \quad (32)$$

for all  $u, v, w \in \mathbb{R}^*$ . Then the function  $T_r$  serves as a counter-example for the fact that the system of equations (3) is not stable for  $\alpha = -3$  in Corollary 1 in the following theorem.

**Theorem 3** *If the function  $T_r : \mathbb{R}^* \times \mathbb{R}^* \times \mathbb{R}^* \longrightarrow \mathbb{R}$  defined in (32) satisfies the following functional inequalities*

$$\left. \begin{aligned} |\Delta_1 T_r(u_1, u_2, v, w)| &\leq \frac{12k}{7} (|u_1|^{-3} + |u_2|^{-3} + |v|^{-3} + |w|^{-3}), \\ |\Delta_2 T_r(u, v_1, v_2, w)| &\leq \frac{12k}{7} (|u|^{-3} + |v_1|^{-3} + |v_2|^{-3} + |w|^{-3}), \\ |\Delta_3 T_r(u, v, w_1, w_2)| &\leq \frac{12k}{7} (|u|^{-3} + |v|^{-3} + |w_1|^{-3} + |w_2|^{-3}), \end{aligned} \right\} \quad (33)$$

for all  $u_1, u_2, v_2, w_1, w_2 \in \mathbb{R}^*$ , then there do not exist a tri-reciprocal mapping  $T_R : \mathbb{R}^* \times \mathbb{R}^* \times \mathbb{R}^* \longrightarrow \mathbb{R}$  and a constant  $\mu > 0$  such that

$$|T_r(u, v, w) - T_R(u, v, w)| \leq \mu (|u|^{-3} + |v|^{-3} + |w|^{-3}) \quad (34)$$

for all  $u, v, w \in \mathbb{R}^*$ .

**Proof** Firstly, let us prove that  $T_r$  satisfies (34). By computation, we have

$$|T_r(u, v, w)| = \left| \sum_{n=0}^{\infty} 8^{-n} \varphi(2^{-n}x) \right| \leq \sum_{n=0}^{\infty} \frac{k}{8^n} = \frac{8k}{7}.$$

Therefore, we see that  $T_r$  is bounded by  $\frac{8k}{7}$  on  $\mathbb{R}$ . If  $|u_1|^{-3} + |u_2|^{-3} + |v|^{-3} + |w|^{-3} \geq 1$ ,  $|u|^{-3} + |v_1|^{-3} + |v_2|^{-3} + |w|^{-3} \geq 1$  and  $|u|^{-3} + |v|^{-3} + |w_1|^{-3} + |w_2|^{-3} \geq 1$  then the left hand side of each inequalities in (33) is less than  $\frac{12k}{7}$ . Now, suppose that  $0 < |u_1|^{-3} + |u_2|^{-3} + |v|^{-3} + |w|^{-3} < 1$ . Hence, there exists a positive integer  $k$  such that

$$\frac{1}{8^{k+1}} \leq |u_1|^{-3} + |u_2|^{-3} + |v|^{-3} + |w|^{-3} < \frac{1}{8^k}. \quad (35)$$

The relation (35) implies  $8^k (|u_1|^{-3} + |u_2|^{-3} + |v|^{-3} + |w|^{-3}) < 1$ , or equivalently;  $8^k u_1^{-3} < 1$ ,  $8^k u_2^{-3} < 1$ ,  $8^k v^{-3} < 1$ ,  $8^k w^{-3} < 1$ . So,  $\frac{u_1^3}{8^k} > 1$ ,  $\frac{u_2^3}{8^k} > 1$ ,  $\frac{v^3}{8^k} > 1$ ,  $\frac{w^3}{8^k} > 1$ . The last inequalities imply that  $\frac{u_1^3}{8^{k-1}} > 8 > 1$ ,  $\frac{u_2^3}{8^{k-1}} > 8 > 1$ ,  $\frac{v^3}{8^{k-1}} > 8 > 1$ ,  $\frac{w^3}{8^{k-1}} > 8 > 1$  and consequently

$$\frac{1}{2^{k-1}}(u_1) > 1, \frac{1}{2^{k-1}}(u_2) > 1, \frac{1}{2^{k-1}}(v) > 1, \frac{1}{2^{k-1}}(w) > 1, \frac{1}{2^{k-1}}(u_1 + u_2) > 1.$$

Therefore, for each value of  $n = 0, 1, 2, \dots, k-1$ , we obtain

$$\frac{1}{2^n}(u_1) > 1, \frac{1}{2^n}(u_2) > 1, \frac{1}{2^n}(v) > 1, \frac{1}{2^n}(w) > 1, \frac{1}{2^n}(u_1 + u_2) > 1.$$

and  $\Delta_1 \varphi(2^{-n}u_1, 2^{-n}u_2, 2^{-n}v, 2^{-n}w) = 0$  for  $n = 0, 1, 2, \dots, k-1$ . Using (31) and the definition of  $T_r$ , we obtain

$$\begin{aligned} |\Delta_1 T_r(u_1, u_2, v, w)| &\leq \sum_{n=k}^{\infty} \frac{k}{8^n} + \frac{1}{2} \sum_{n=k}^{\infty} \frac{k}{8^n} \leq \frac{3k}{2} \frac{1}{8^k} \left(1 - \frac{1}{8}\right)^{-1} \\ &\leq \frac{3k}{2} \frac{1}{8^k} \leq \frac{12k}{7} \frac{1}{8^{k+1}} \leq \frac{12k}{7} (|u_1|^{-3} + |u_2|^{-3} + |v|^{-3} + |w|^{-3}) \end{aligned}$$

for all  $u_1, u_2, v, w \in \mathbb{R}^*$ . Therefore, the first inequality in (33) holds good. Similarly, we can show that the second and third inequalities in (33) hold good. Now, we claim that the system of equations (3) is not stable for  $\alpha = -3$  in Corollary 1. Assume that there exists a tri-reciprocal mapping  $T : \mathbb{R}^* \times \mathbb{R}^* \times \mathbb{R}^* \rightarrow \mathbb{R}$  satisfying (34). Therefore, we have

$$|T_r(u, v, w)| \leq (\mu + 1) (|u|^{-3} + |v|^{-3} + |w|^{-3}). \quad (36)$$

However, we can choose a positive integer  $m$  with  $m\delta > \mu + 1$ . If  $u, v, w \in (1, 2^{m-1})$  then  $2^{-n}u, 2^{-n}v, 2^{-n}w \in (1, \infty)$ , for all  $n = 0, 1, 2, \dots, m-1$  and thus

$$\begin{aligned} |T_r(u, v, w)| &= \sum_{n=0}^{\infty} \frac{\varphi(2^{-n}x)}{8^n} \geq \sum_{n=0}^{m-1} \frac{8^n k (u^{-3} + v^{-3} + w^{-3})}{8^n} \\ &= mk (u^{-3} + v^{-3} + w^{-3}) > (\mu + 1) (u^{-3} + v^{-3} + w^{-3}) \end{aligned}$$

which contradicts (36). Therefore, the system of equations (3) is not stable for  $\alpha = -3$  in Corollary 1.  $\square$

The upcoming result indicates that the system of equations (3) is not stable for  $\alpha = -3$  in Corollary 3. Since the method of proof is similar, we omit the proof.

**Theorem 4** *If the function  $T_r : \mathbb{R}^* \times \mathbb{R}^* \times \mathbb{R}^* \rightarrow \mathbb{R}$  defined in (32) satisfies the following functional inequalities*

$$\left. \begin{aligned} |\Delta_1 T_r(u_1, u_2, v, w)| &\leq \theta_3 \left( |u_1|^{-\frac{1}{2}} |u_2|^{-\frac{1}{2}} |v|^{-1} |w|^{-1} + (|u_1|^{-3} + |u_2|^{-3} + |v|^{-3} + |w|^{-3}) \right), \\ |\Delta_2 T_r(u, v_1, v_2, w)| &\leq \theta_3 \left( |u|^{-1} |v_1|^{-\frac{1}{2}} |v_2|^{-\frac{1}{2}} |w|^{-1} + (|u|^{-3} + |v_1|^{-3} + |v_2|^{-3} + |w|^{-3}) \right), \\ |\Delta_3 T_r(u, v, w_1, w_2)| &\leq \theta_3 \left( |u|^{-1} |v|^{-1} |w_1|^{-\frac{1}{2}} |w_2|^{-\frac{1}{2}} + (|u|^{-3} + |v|^{-3} + |w_1|^{-3} + |w_2|^{-3}) \right), \end{aligned} \right\} \quad (37)$$

for all  $u_1, u_2, v_2, w_1, w_2 \in \mathbb{R}^*$ , then there do not exist a tri-reciprocal mapping  $T_R : \mathbb{R}^* \times \mathbb{R}^* \times \mathbb{R}^* \rightarrow \mathbb{R}$  and a constant  $\mu > 0$  such that

$$|T_r(u, v, w) - T_R(u, v, w)| \leq \mu (|u|^{-1} |v|^{-1} |w|^{-1} + (|u|^{-3} + |v|^{-3} + |w|^{-3})) \quad (38)$$

for all  $u, v, w \in \mathbb{R}^*$ .


Counter-examples similar to Theorems 3 and 4 can be considered for the non-stability of system of equations in Corollary 4 and 9.

## An Exertion of the System of Equation (2)

In this section, we associate the system of functional equations (3) with the relationship between density and volume of any object. We know, density is the mass per unit volume of any object. In this section, we assume that the mass of the object to be a constant. Imagine a cuboid shaped sponge of dimensions of length  $u$ , breadth  $v$  and height  $w$  as in Figure 1. Then the rational function  $T_r(u, v, w) = \frac{c}{uvw}$  represents the density of the sponge, which means that if volume increases, then density decreases with constant mass. This can be illustrated in the following occasion. If the sponge is squeezed, then the density of the squeezed sponge is more when compared to the less density of the loosely left sponge.

Now, divide the sponge shown in Figure 1 into two subcuboids along its length as portrayed in Figure 2. Then the densities of the subcuboids are  $T_r(u_1, v, w) = \frac{c}{u_1vw}$  and  $T_r(u_2, v, w) = \frac{c}{u_2vw}$ , respectively, where  $u_1 + u_2 = u$ . Hence, the density of the original cuboid shaped sponge is half of the harmonic mean of the volumes of the two sub cuboids, which satisfies the first member of the system of equations (3) dealt in our study. Similarly, if the original cuboid shaped sponge is divided along its breadth and height, the second and third members of the system of equations (3) satisfy the above fact relating density and volume.

**Fig. 1** Cuboid sponge



**Fig. 2** Division of cuboid sponge into two subcuboids



## Conclusion

In this investigation, we have carried out various stabilities of the system of tri-reciprocal functional equations (3). We illustrated counter-examples that the stability results of system of equations (3) do not hold for singular cases. Since the solution of the system (3) is a rational function, both indeterminable and minuscule numbers could be considered in the domain. This is the first attempt that we have considered the set of hyperreals as domain for the investigation of stability results. Also, the notion of non-standard analysis could be identified via functional equations. We have also exhibited the role of the system of functional equations dealt in this study in connection with volume and density of a cuboid shaped sponge.

## References

- Almahalebi, M., Chahbi, A., Kabbaj, S.: A fixed point approach to the stability of a bi-cubic functional equation in 2-Banach spaces. *Palest. J. Math.* **5**(2), 220–227 (2016)
- Alshybani, S., Vaezpour, S.M., Saadati, R.: Stability of the sextic functional equation in various spaces. *J. Inequ. Spec. Func.* **9**(4), 8–27 (2018)
- Arunkumar, M., Karthikeyan, S.: Solution and stability of a reciprocal functional equation originating from  $n$ -consecutive terms of a harmonic progression: Direct and fixed point methods. *Int. J. Inf. Sci. Intell. Syst.* **3**(1), 151–168 (2014)
- Arunkumar, M., Karthikeyan, S.: Fuzzy Banach algebra stability of reciprocal quadratic functional equation via fixed point approach. *Int. J. Pure Appl. Math.* **119**(3), 31–39 (2018)
- Bodaghi, A., Kim, S.O.: Approximation on the quadratic reciprocal functional equation, *J. Func. Spac. Appl.* Article ID532463 (2014)
- Bodaghi, A., Narasimman, P., Rassias, J.M., Ravi, K.: Ulam stability of the reciprocal functional equation in non-Archimedean fields. *Acta Math. Univ. Comen.* **85**(1), 113–124 (2016)
- Bodaghi, A., Rassias, J.M., Park, C.: Fundamental stabilities of an alternative quadratic reciprocal functional equation in non-Archimedean fields. *Proc. Jangjeon Math. Soc.* **18**(3), 313–320 (2015)
- Bodaghi, A., Senthil Kumar, B.V.: Estimation of inexact reciprocal-quintic and reciprocal-sextic functional equations. *Mathematica*, **49**(82), No. 1-2, 3–14 (2017)
- Ebadian, A., Zolfaghari, S., Ostadbashi, S., Park, C.: Approximation on the reciprocal functional equation in several variables in matrix non-Archimedean random normed spaces. *Adv. Differ. Equ.* **314**, 2015 (2015). <https://doi.org/10.1186/s13662-015-0656-7>
- Fassi, Z.I., Brzdek, J., Chahbi, A., Kabbaj, S.: On hyperstability of the biadditive functional equation. *Acta Mathematica Scientia* **37**(6), 1727–1739 (2017)
- Gajda, Z.: On the stability of additive mappings. *Int. J. Math. Math. Sci.* **14**, 431–434 (1991)
- Gavruta, P.: A generalization of the Hyers–Ulam–Rassias stability of approximately additive mappings. *J. Math. Anal. Appl.* **184**, 431–436 (1994)
- Goldblatt, R.: *Lectures on the Hyperreals; An Introduction to Nonstandard Analysis*, Volume of 188 Graduate Texts in Mathematics. Springer, New York (1998)
- Govindan, V., Balasubramanian, G., Muthamilarasi, C.: Generalized stability of cubic functional equation. *Int. J. Sci. Eng. Sci.* **2**(1), 74–82 (2018)
- Hejmej, B.: Stability of functional equations in dislocated quasi-metric spaces. *Annales Mathematicae Silesianae* **32**, 215–225 (2018)
- Hooda, N., Tomar, S.: Non-Archimedean stability and non-stability of quadratic reciprocal functional equation in several variables. *Bull. Pure Appl. Sci.* **37E**(2), 267–272 (2018)
- Hooda, N., Tomar, S.: Approximation of reciprocal-cubic functional equation in non-Archimedean normed space. *Int. J. Sci. Res. Math. Stat. Sci.* **5**(5), 169–172 (2018)
- Hyers, D.H.: On the stability of the linear functional equation. *Proc. Nat. Acad. Sci. USA* **27**, 222–224 (1941)
- Kim, S.O., Senthil Kumar, B.V., Bodaghi, A.: Stability and non-stability of the reciprocal-cubic and reciprocal-quartic functional equations in non-Archimedean fields. *Adv. Differ. Equ.* **77**, 12 (2017)
- Murali, R., Antony Raj, A., Thamizharasan, T.: Stability of reciprocal difference and adjoint functional equations in multi-Banach spaces: a fixed point method. *J. Emerg. Technol. Innov. Res.* **5**(8), 450–455 (2018)

21. Narasimman, P.: Solution and stability of a generalized kadditive functional equation. *J. Interdiscip. Math.* **21**(1), 171–184 (2018)
22. Park, W.G., Bae, J.H.: Solution of a vector variable bi-additive functional equation. *Commun. Korean Math. Soc.* **23**(2), 191–199 (2008)
23. Park, W.G., Bae, J.H.: Stability of a bi-additive functional equation in Banach modules over a  $C^*$ -algebra. *Discrete Dyn. Nat. Soc.* Article ID 835893 (2012)
24. Pinelas, S., Arunkumar, M., Sathya, E.: Hyers type stability of a radical reciprocal quadratic functional equation originating from 3 dimensional Pythagorean means. *Int. J. Math. Appl.* **5**(4), 45–52 (2017)
25. Prager, W., Schwaiger, J.: Stability of the multi-Jensen equation. *Bull. Korean Math. Soc.* **45**(1), 133–142 (2008)
26. Rajkumar, V.: The generalized Hyers–Ulam–Rassias stability of a quadratic functional equation. *Int. J. Res. Sci. Innov.* **5**(7), 73–76 (2018)
27. Rassias, J.M.: On approximately of approximately linear mappings by linear mappings. *J. Funct. Anal. USA* **46**, 126–130 (1982)
28. Rassias, T.M.: On the stability of the linear mapping in Banach spaces. *Proc. Am. Math. Soc.* **72**, 297–300 (1978)
29. Ravi, K., Arunkumar, M., Rassias, J.M.: On the Ulam stability for the orthogonally general Euler–Lagrange type functional equation. *Int. J. Math. Sci.* **3**(8), 36–47 (2008)
30. Ravi, K., Rassias, J.M., Pinelas, S., Narasimman, P.: The stability of a generalized radical reciprocal quadratic functional equation in Felbin’s space. *PanAm. Math. J.* **24**(1), 75–92 (2014)
31. Ravi, K., Senthil Kumar, B.V.: Ulam–Gavruta–Rassias stability of Rassias reciprocal functional equation. *Glob. J. Appl. Math. Sci.* **3**(1–2), 57–79 (2010)
32. Ravi, K., Senthil Kumar, B.V.: Generalized Hyers–Ulam–Rassias stability of a system of bi-reciprocal functional equations. *Euro. J. Pure Appl. Math.* **8**(2), 283–293 (2015)
33. Ravi, K., Suresh, S.: Generalized Hyers–Ulam stability of a cubic reciprocal functional equation. *Br. J. Math. Comput. Sci.* **20**(6), 1–9 (2017)
34. Senthil Kumar, B.V., Dutta, H.: Non-Archimedean stability of a generalized reciprocal-quadratic functional equation in several variables by direct and fixed point methods. *Filomat* **32**(9), 3199–3209 (2018)
35. Senthil Kumar, B.V., Dutta, H.: Fuzzy stability of a rational functional equation and its relevance to system design. *Int. J. Gen. Syst.* **48**(2), 157–169 (2019)
36. Senthil Kumar, B.V., Dutta, H.: Approximation of multiplicative inverse undecic and duodecic functional equations. *Math. Meth. Appl. Sci.* **42**, 1073–1081 (2019)
37. Song, A., Song, M.: The stability of quadratic-reciprocal functional equation. In: *AIP Conference Proceedings* 1955, 040171 (2018). <https://doi.org/10.1063/1.5033835>
38. Sophia, M.: Generalized Hyers–Ulam–Rassias stability of a reciprocal type functional equation in non-Archimedean fields. *Int. J. Eng. Trends Technol.* **55**(3), 153–158 (2018)
39. Ulam, S.M.: *Problems in Modern Mathematics*, Chapter VI. Wiley-Interscience, New York (1964)
40. White, M.: Incommensurables and incomparables: on the conceptual status and the philosophical use of hyperreal numbers. *Notre Dame J. Form. Log.* **40**, 420–446 (1999)

**Publisher’s Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.