

Midterm 2

Hishem Kløvnnes

November 14, 2024

1 Quantum mechanics for many-particle systems

a)

For this exercise, we will use the commutation relations:

$$[A, BC] = [A, B]C + B[A, C]$$

and

$$[AB, CD] = A[B, C]D + AC[B, D] + [A, C]DB + C[A, D]B$$

We want to show that \hat{H}_0 and \hat{V} commutes with \hat{S}_z and \hat{S}^2 .

The operators are defined as:

$$\hat{H}_0 = \xi \sum_{p\sigma} (p-1) a_{p\sigma}^\dagger a_{p\sigma}$$

$$\hat{V} = \sum_{pq} a_{p-}^\dagger a_{p+}^\dagger a_{p-} a_{p+}$$

$$\hat{S}_z = \sum_{p\sigma} \sigma a_{p\sigma}^\dagger a_{p\sigma}$$

$$\hat{S}^2 = \hat{S}_z^2 + \frac{1}{2}(\hat{S}_+ \hat{S}_- + \hat{S}_- \hat{S}_+)$$

Where $\hat{S}_\pm = \sum_p a_{p\pm}^\dagger a_{p\mp}$, $\sigma = \pm 1$ for spin and $p = 1, 2, \dots$ is the index for the single-particle states.

We start by showing that \hat{H}_0 commutes with \hat{S}_z :

$$[\hat{H}_0, \hat{S}_z] = \xi \sum_{p\sigma} (p-1) [a_{p\sigma}^\dagger a_{p\sigma}, \sum_{\tau} \tau a_{q\tau}^\dagger a_{q\tau}]$$

Lets take the operators to the side and look at those first:

$$\begin{aligned} & [a_{p\sigma}^\dagger a_{p\sigma}, a_{q\tau}^\dagger a_{q\tau}] \\ &= a_{p\sigma}^\dagger [a_{p\sigma}, a_{q\tau}^\dagger] a_{q\tau} + a_{p\sigma}^\dagger a_{q\tau}^\dagger [a_{p\sigma}, a_{q\tau}] + [a_{p\sigma}^\dagger, a_{q\tau}^\dagger] a_{p\sigma} a_{q\tau} + a_{q\tau}^\dagger [a_{p\sigma}^\dagger, a_{q\tau}] a_{p\sigma} \end{aligned}$$

Two creation operators commute and two annihilation operators commute, so two of them are directly zero, while the other two leaves croenecker deltas

$$\begin{aligned}
&= a_{p\sigma}^\dagger \delta_{pq} \delta_{\sigma\tau} a_{q\tau} - a_{q\tau}^\dagger \delta_{pq} \delta_{\sigma\tau} a_{p\sigma} \\
[\hat{H}_0, \hat{S}_z] &= \xi \sum_{p\sigma} (p-1) [a_{p\sigma}^\dagger a_{p\sigma}, \sum_{\tau} \tau a_{q\tau}^\dagger a_{q\tau}] \\
&= \xi \sum_{p\sigma} (p-1) \sum_{\tau} \tau (a_{p\sigma}^\dagger \delta_{pq} \delta_{\sigma\tau} a_{q\tau} - a_{q\tau}^\dagger \delta_{pq} \delta_{\sigma\tau} a_{p\sigma}) = 0
\end{aligned}$$

Now we want to show that \hat{H}_0 commutes with \hat{S}^2 .

$$[\hat{H}_0, \hat{S}^2] = [H_0, S_z^2] + \frac{1}{2}([H_0, S_+ S_-] + [H_0, S_- S_+])$$

We know that $[H_0, S_z] = 0$ and therefor the first term is zero.

Lets look at the second term:

$$\begin{aligned}
[H_0, S_+ S_-] &= \xi \sum_{pq\sigma} (p-1) [a_{p\sigma}^\dagger a_{p\sigma}, a_{q+}^\dagger a_{q-} a_{q-}^\dagger a_{q+}] \\
&= \xi \sum_{pq\sigma} (p-1) ([a_{p\sigma}^\dagger a_{p\sigma}, a_{q+}^\dagger a_{q-}] a_{q-}^\dagger a_{q+} + a_{q-}^\dagger a_{q+} [a_{p\sigma}^\dagger a_{p\sigma}, a_{q+}^\dagger a_{q-}]) \\
&= \xi \sum_{pq\sigma} (p-1) \left((a_{p\sigma}^\dagger [a_{p\sigma}, a_{q+}^\dagger] a_{q-} + a_{p\sigma}^\dagger a_{q-} [a_{p\sigma}, a_{q+}^\dagger]) a_{q-}^\dagger a_{q+} \right. \\
&\quad \left. + a_{q+}^\dagger a_{q-} (a_{p\sigma}^\dagger [a_{p\sigma}, a_{q-}^\dagger] a_{q+} + a_{p\sigma}^\dagger a_{q+} [a_{p\sigma}, a_{q-}^\dagger]) \right) \\
&= \xi \sum_{p\sigma} (p-1) \left((a_{p+}^\dagger a_{p-} - a_{p-}^\dagger a_{p+}) a_{p-}^\dagger a_{p+} + a_{p+}^\dagger a_{p-} (a_{p-}^\dagger a_{p+} - a_{p-}^\dagger a_{p+}) \right) = 0
\end{aligned}$$

The same goes for the last term, and therefor $[H_0, S^2] = 0$

Now we can move over to check if \hat{V} commutes with \hat{S}_z and \hat{S}^2 . First we check if \hat{V} commutes with \hat{S}_z :

$$\begin{aligned}
[\hat{V}, \hat{S}_z] &= \frac{1}{4} g [\sum_{pq} a_{p-}^\dagger a_{p+}^\dagger a_{q-} a_{q+}, \sum_{r\sigma} \sigma a_{r\sigma}^\dagger a_{r\sigma}] \\
&= \frac{1}{4} g \sum_{pqr} \sigma \left(a_{p+}^\dagger a_{p-}^\dagger [a_{q-} a_{q+}, a_{r\sigma}^\dagger a_{r\sigma}] + [a_{p+}^\dagger a_{p-}^\dagger, a_{r\sigma}^\dagger a_{r\sigma}] a_{q-} a_{q+} \right)
\end{aligned}$$

The first term:

$$\begin{aligned}
[a_{q-} a_{q+}, a_{r\sigma}^\dagger a_{r\sigma}] &= a_{q-} [a_{q+}, a_{r\sigma}^\dagger] a_{r\sigma} + a_{q-} a_{r\sigma}^\dagger [a_{q+}, a_{r\sigma}] \\
&\quad + [a_{q-}, a_{r\sigma}^\dagger] a_{r\sigma} a_{q+} + a_{r\sigma}^\dagger [a_{q-}, a_{r\sigma}] a_{q+}
\end{aligned}$$

Where the only surviving term, the ones with one creation and one annihilation operator, are:

$$\begin{aligned}
&= a_{q-} \delta_{qr} \delta_{\sigma+} a_{r\sigma} + a_{q+} \delta_{qr} \delta_{\sigma-} a_{r\sigma} \\
&= a_{q-} a_{q+} + a_{q+} a_{q-}
\end{aligned}$$

Now the second term:

$$\begin{aligned} [a_{p+}^\dagger a_{p-}^\dagger, a_{r\sigma}^\dagger a_{r\sigma}] &= a_{p+}^\dagger [a_{p-}^\dagger, a_{r\sigma}^\dagger] a_{r\sigma} + a_{p+}^\dagger a_{r\sigma}^\dagger [a_{p-}^\dagger, a_{r\sigma}] \\ &\quad + [a_{p+}^\dagger, a_{r\sigma}^\dagger] a_{r\sigma} a_{p-}^\dagger + a_{r\sigma}^\dagger [a_{p+}^\dagger, a_{r\sigma}] a_{p-}^\dagger \end{aligned}$$

Where the only surviving term, the ones with one creation and one annihilation operator, are:

$$\begin{aligned} &= -a_{p+}^\dagger \delta_{pr} \delta_{\sigma-} a_{r\sigma} - a_{p-}^\dagger \delta_{pr} \delta_{\sigma+} a_{r\sigma} \\ &= a_{p+}^\dagger a_{p-} + a_{p-}^\dagger a_{p+} \end{aligned}$$

Now we can put the two results back into the first expression, and we can see that it simply reduces to zero.

$$= \frac{1}{4} g \sum_{pq} \sigma \left(a_{p+}^\dagger a_{p-}^\dagger (a_{q-} a_{q+} + a_{q-} a_{q+}) - a_{q-} a_{q+} (a_{p+}^\dagger a_{p-}^\dagger + a_{p+}^\dagger a_{p-}^\dagger) \right) = 0$$

And now we can check if \hat{V} commutes with \hat{S}^2 :

$$[\hat{V}, \hat{S}^2] = [\hat{V}, \hat{S}_z^2] + \frac{1}{2} ([\hat{V}, \hat{S}_+ \hat{S}_-] + [\hat{V}, \hat{S}_- \hat{S}_+])$$

The first term is zero from the fact that \hat{V} commutes with \hat{S}_z . The second term is:

$$\begin{aligned} [\hat{V}, \hat{S}_+ \hat{S}_-] &= -\frac{1}{2} g \sum_{pq} [a_{p+}^\dagger a_{p-}^\dagger a_{q-} a_{q+}, \sum_r a_{r+}^\dagger a_{r-} a_{r-}^\dagger a_{r+}] \\ &= -\frac{1}{2} g \sum_{pqr} \left(a_{p+}^\dagger a_{p-}^\dagger [a_{q-} a_{q+}, a_{r+}^\dagger a_{r-}] a_{r-}^\dagger a_{r+} \right. \\ &\quad + a_{p+}^\dagger a_{p-}^\dagger a_{r+}^\dagger a_{r-} [a_{q-} a_{q+}, a_{r-}^\dagger a_{r+}] \\ &\quad + [a_{p+}^\dagger a_{p-}^\dagger, a_{r+}^\dagger a_{r-}] a_{r-}^\dagger a_{r+} a_{q-} a_{q+} \\ &\quad \left. + a_{r+}^\dagger a_{r-} [a_{p+}^\dagger a_{p-}^\dagger, a_{r-}^\dagger a_{r+}] a_{q-} a_{q+} \right) \end{aligned}$$

I will separate the commutations and look at them one by one. The first term:

$$\begin{aligned} [a_{q-} a_{q+}, a_{r+}^\dagger a_{r-}] &= a_{q-} [a_{q+}, a_{r+}^\dagger] a_{r-} + a_{q-} a_{r+}^\dagger [a_{q+}, a_{r-}] + [a_{q-}, a_{r+}^\dagger] a_{r-} a_{q+} + a_{r-} [a_{q-}, a_{r+}^\dagger] a_{q+} \\ &= a_{q-} \delta_{qr} \delta_{++} a_{r-} + \delta_{qr} \delta_{+-} a_{r-} a_{q+} = a_{q-} a_{q-} \end{aligned}$$

The second term:

$$\begin{aligned} [a_{q-} a_{q+}, a_{r-}^\dagger a_{r+}] &= a_{q-} [a_{q+}, a_{r-}^\dagger] a_{r+} + a_{q-} a_{r-}^\dagger [a_{q+}, a_{r-}] + [a_{q-}, a_{r-}^\dagger] a_{r+} a_{q+} + a_{r-}^\dagger [a_{q-}, a_{r+}] a_{q+} \\ &= a_{q-} \delta_{qr} \delta_{+-} a_{r+} + \delta_{qr} \delta_{+-} a_{r+} a_{q+} = a_{q+} a_{q+} \end{aligned}$$

The third term:

$$\begin{aligned} [a_{p+}^\dagger a_{p-}^\dagger, a_{r+}^\dagger a_{r-}] &= a_{p+}^\dagger [a_{p-}^\dagger, a_{r+}^\dagger] a_{r-} + a_{p+}^\dagger a_{r+}^\dagger [a_{p-}^\dagger, a_{r-}] + [a_{p+}^\dagger, a_{r+}^\dagger] a_{r-} a_{p-}^\dagger + a_{r+}^\dagger [a_{p+}^\dagger, a_{r-}] a_{p-}^\dagger \\ &= -a_{p+}^\dagger a_{r+}^\dagger \delta_{pr} \delta_{--} - a_{r+}^\dagger \delta_{pr} \delta_{+-} a_{p-}^\dagger = -a_{p+}^\dagger a_{p+}^\dagger \end{aligned}$$

The fourth term:

$$\begin{aligned} [a_{p+}^\dagger a_{p-}^\dagger, a_{r-}^\dagger a_{r+}] &= a_{p+}^\dagger [a_{p-}^\dagger, a_{r-}^\dagger] a_{r+} + a_{p+}^\dagger a_{r-}^\dagger [a_{p-}^\dagger, a_{r+}] + [a_{p+}^\dagger, a_{r-}^\dagger] a_{r+} a_{p-}^\dagger + a_{r-}^\dagger [a_{p+}^\dagger, a_{r+}] a_{p-}^\dagger \\ &= -a_{p+}^\dagger a_{r-}^\dagger \delta_{pr} \delta_{-+} - a_{r-}^\dagger \delta_{pr} \delta_{++} a_{p-}^\dagger = -a_{p-}^\dagger a_{p-}^\dagger \end{aligned}$$

Now we can put the results back into the expression, and since r is equal to either p or q , we can adjust according to this:

$$\begin{aligned} &= -\frac{1}{2}g \sum_{pq} \left(a_{p+}^\dagger a_{p-}^\dagger a_{q-} a_{q-} a_{q-}^\dagger a_{q+} \right. \\ &\quad + a_{p+}^\dagger a_{p-}^\dagger a_{q+} a_{q-} a_{q+} a_{q+} \\ &\quad - a_{p+}^\dagger a_{p+}^\dagger a_{p-}^\dagger a_{p+} a_{q-} a_{q+} \\ &\quad \left. - a_{p+}^\dagger a_{p-}^\dagger a_{p-}^\dagger a_{p-} a_{q-} a_{q+} \right) \end{aligned}$$

All of these terms are zero, because if we look closely, we can either see that each term is trying to annihilate the same state twice. For the second and third term this happens directly, while for the first and last term, this happens when trying to normal order the operators.

The last term is the same as the second term, because of the commutation relations between \hat{S}_+ and \hat{S}_- , and therefore the result is zero.

This means that \hat{V} commutes with \hat{S}^2 .

We can now introduce the pair-creation and pair-annihilation operators:

$$\begin{aligned} \hat{P}_p^+ &= a_{p+}^\dagger a_{p-}^\dagger \\ \hat{P}_p^- &= a_{p-} a_{p+} \end{aligned}$$

Lets re-introduce the \hat{V} operator:

$$\hat{V} = -\frac{1}{2}g \sum_{pq} a_{p+}^\dagger a_{p-}^\dagger a_{q-} a_{q+}$$

From this we can easily see that we can write \hat{V} as:

$$\hat{V} = -\frac{1}{2}g \sum_{pq} \hat{P}_p^+ \hat{P}_q^-$$

And therefore we can write the \hat{H} , with $\xi = 1$ as:

$$\hat{H} = \hat{H}_0 + \hat{V} = \sum_{p\sigma} (p-1) a_{p\sigma}^\dagger a_{p\sigma} - \frac{1}{2}g \sum_{pq} \hat{P}_p^+ \hat{P}_q^-$$

And lastly we want to show that the pair creation operators commutes among themselves:

$$[\hat{P}_p^+, \hat{P}_q^+] = [a_{p+}^\dagger a_{p-}^\dagger, a_{q+}^\dagger a_{q-}^\dagger]$$

$$a_{p+}^\dagger [a_{p-}^\dagger, a_{q+}^\dagger] a_{q-}^\dagger + a_{p+}^\dagger a_{q+}^\dagger [a_{p-}^\dagger, a_{q-}^\dagger] + [a_{p+}^\dagger, a_{q+}^\dagger] a_{q-}^\dagger a_{p-}^\dagger + a_{q+}^\dagger [a_{p+}^\dagger, a_{q-}^\dagger] a_{p-}^\dagger$$

All these terms are zero, because commuting two creation or annihilation operators will give zero.

b)

For this part we want to construct the Hamiltonian matrix for a system with no broken pairs and total spin of $S = 0$, for the case of the four lowest single-particle states.

We can start by defining a state with total spin $S = 0$:

$$|\Phi_{\alpha\beta}\rangle = P_{\alpha}^{+} P_{\beta}^{+} |0\rangle$$

We will start by finding the expectation value of the one body term:

$$\begin{aligned} \langle \Phi_{\alpha\beta} | \hat{H}_0 | \Phi_{\alpha\beta} \rangle &= \langle \Phi_{\alpha\beta} | \sum_{p\sigma} (p-1) a_{p\sigma}^{\dagger} a_{p\sigma} | \Phi_{\alpha\beta} \rangle \\ &= \sum_{p\sigma} (p-1) \langle \Phi_{\alpha\beta} | a_{p\sigma}^{\dagger} a_{p\sigma} | \Phi_{\alpha\beta} \rangle \end{aligned}$$

Here $\sigma = \pm 1$ and p can have the values $p = \alpha, \beta$, and therefor we can write the expression as:

$$\langle \Phi_{\alpha\beta} | \hat{H}_0 | \Phi_{\alpha\beta} \rangle = 2(\alpha-1) + 2(\beta-1) = 2(\alpha + \beta - 2)$$

For the interaction term \hat{V} , we will use the indexes $\alpha, \beta, \gamma, \delta$ to represent the four lowest single-particle states. We can write the expectation value as:

$$\begin{aligned} \langle \Phi_{\alpha\beta} | \hat{V} | \Phi_{\alpha\beta} \rangle &= -\frac{1}{2}g \langle \Phi_{\alpha\beta} | \sum_{pq} \hat{P}_p^{+} \hat{P}_q^{-} | \Phi_{\gamma\delta} \rangle \\ &= -\frac{1}{2}g \sum_{pq} \langle 0 | P_{\beta}^{-} P_{\alpha}^{-} P_p^{+} P_q^{-} P_{\gamma}^{+} P_{\delta}^{+} | 0 \rangle \end{aligned}$$

For this we will need to use contractions.

$$\begin{aligned} \overbrace{P_{\beta}^{-} P_{\alpha}^{-} P_p^{+} P_q^{-} P_{\gamma}^{+} P_{\delta}^{+}} &= \delta_{\beta\delta} \delta_{\alpha p} \delta_{q\gamma} \\ \overbrace{P_{\beta}^{-} P_{\alpha}^{-} P_p^{+} P_q^{-} P_{\gamma}^{+} P_{\delta}^{+}} &= \delta_{\beta\gamma} \delta_{\alpha p} \delta_{q\delta} \\ \overbrace{P_{\beta}^{-} P_{\alpha}^{-} P_p^{+} P_q^{-} P_{\gamma}^{+} P_{\delta}^{+}} &= \delta_{\beta p} \delta_{\alpha\gamma} \delta_{q\delta} \\ \overbrace{P_{\beta}^{-} P_{\alpha}^{-} P_p^{+} P_q^{-} P_{\gamma}^{+} P_{\delta}^{+}} &= \delta_{\beta p} \delta_{\alpha\delta} \delta_{q\gamma} \end{aligned}$$

Usually, when we work with contraction, we will notice how many of the lines cross to determine if the term is negative or positive. One cross, and the term is negative, while two crosses and the term is positive.

For these operators, each of them contain either two creation or two annihilation operators, and therefor the result will always be positive.

Now we can set up the matrices:

$$\hat{H}_0 = \begin{bmatrix} 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 & 0 & 0 \\ 0 & 0 & 6 & 0 & 0 & 0 \\ 0 & 0 & 0 & 6 & 0 & 0 \\ 0 & 0 & 0 & 0 & 8 & 0 \\ 0 & 0 & 0 & 0 & 0 & 10 \end{bmatrix}$$

$$\hat{V} = -\frac{1}{2}g \begin{bmatrix} 2 & 1 & 1 & 1 & 1 & 0 \\ 1 & 2 & 1 & 1 & 0 & 1 \\ 1 & 1 & 2 & 0 & 1 & 1 \\ 1 & 1 & 0 & 2 & 1 & 1 \\ 1 & 0 & 1 & 1 & 2 & 1 \\ 0 & 1 & 1 & 1 & 1 & 2 \end{bmatrix}$$

The total Hamiltonian matrix will be:

$$\hat{H} = \hat{H}_0 + \hat{V} = \begin{bmatrix} 2-g & -\frac{1}{2}g & -\frac{1}{2}g & -\frac{1}{2}g & -\frac{1}{2}g & 0 \\ -\frac{1}{2}g & 4-g & -\frac{1}{2}g & -\frac{1}{2}g & 0 & -\frac{1}{2}g \\ -\frac{1}{2}g & -\frac{1}{2}g & 6-g & 0 & -\frac{1}{2}g & -\frac{1}{2}g \\ -\frac{1}{2}g & -\frac{1}{2}g & 0 & 6-g & -\frac{1}{2}g & -\frac{1}{2}g \\ -\frac{1}{2}g & 0 & -\frac{1}{2}g & -\frac{1}{2}g & 8-g & -\frac{1}{2}g \\ 0 & -\frac{1}{2}g & -\frac{1}{2}g & -\frac{1}{2}g & -\frac{1}{2}g & 10-g \end{bmatrix}$$

The following plot and the matrix elements are calculated with the code in `energies.py`




Figure 1: The Hamiltonian matrix for a system with no broken pairs and total spin of $S = 0$, for the case of the four lowest single-particle states. Energies as a function of the strength of the interaction g . The dashed lines are to show the degeneracy of the states.

c)

In 1 the energy levels i show are from states $|\Phi_0\rangle$ to $|\Phi_5\rangle$, more explicitly:

$$|\Phi_0\rangle = P_1^+ P_2^+ |0\rangle \quad |\Phi_1\rangle = P_1^+ P_3^+ |0\rangle \quad |\Phi_2\rangle = P_1^+ P_4^+ |0\rangle$$

$$|\Phi_3\rangle = P_2^+ P_3^+ |0\rangle \quad |\Phi_4\rangle = P_2^+ P_4^+ |0\rangle \quad |\Phi_5\rangle = P_3^+ P_4^+ |0\rangle$$

For this part we want to make an approximation to the ground state energy. For this we will use at most two-particle two-holes excitations. Hence we will approximate the




Figure 2: Ground state energy as a function of the interaction strength g .

ground state energy by only using the states $|\Phi_0\rangle$ and $|\Phi_4\rangle$.

In 5 we show the energy difference between the approximation and the ground state




Figure 3: Excluding $|\Phi_5\rangle$. The Hamiltonian matrix for a system with no broken pairs and total spin of $S = 0$, for the case of the four lowest single-particle states. Energies as a function of the strength of the interaction g

energy. We can see that the approximation is very close to the ground state energy as the difference is very small. We can see that the energy difference is negative, meaning the approximated ground state has slightly higher energy than the true ground state energy. TODO: add diagrammatic representation




Figure 4: Ground state energy approximation as a function of the interaction strength g .




Figure 5: Difference between the ground state energy and the approximation as a function of the interaction strength g .

d)

The one-body operator \hat{H}_0 can be typically be expressed in terms of matrix elements $\langle p | \hat{h}_0 | q \rangle$ as:

$$\hat{H}_0 = \sum_{pq} \langle p | \hat{h}_0 | q \rangle a_p^\dagger a_q$$

When normal ordering, we can adjust the term to:

$$a_p^\dagger a_q = \{a_p^\dagger a_q\} + \delta_{pq \in i}$$

And from this we can write the one-body term as:

$$\hat{H}_0 = \sum_{pq} \langle p | \hat{h}_0 | q \rangle \{a_p^\dagger a_q\} + \sum_{pq} \langle p | \hat{h}_0 | q \rangle \delta_{pq \in i}$$

Where the second term is a constant that represents the expectation value of the one-body operator in the reference state:

$$\sum_{pq} \langle p | \hat{h}_0 | q \rangle \delta_{pq \in i} = \langle \Phi_0 | \hat{H}_0 | \Phi_0 \rangle$$

From this we get

$$\hat{H}_0 = \sum_{pq} \langle p | \hat{h}_0 | q \rangle \{a_p^\dagger a_q\} + \langle \Phi_0 | \hat{H}_0 | \Phi_0 \rangle$$

Now applying the spesific case where $\langle p | \hat{h}_0 | q \rangle = (p-1)\delta_{pq}$ and find that

$$\hat{H}_0 = \sum_{p\sigma} (p-1) \{a_p^\dagger a_{p\sigma}\} + \langle \Phi_0 | \hat{H}_0 | \Phi_0 \rangle$$

For the interaction term we will approach the normal ordering in a similar way. We can write the interaction term as:

$$\hat{H}_I = \frac{1}{4} \sum_{pqrs} \langle pq | \hat{v} | rs \rangle a_p^\dagger a_q^\dagger a_s a_r$$

where we can re-write the creation and annihilation operators as:

$$\begin{aligned} a_p^\dagger a_q^\dagger a_s a_r &= \{a_p^\dagger a_q^\dagger a_s a_r\} + a_p^\dagger \overline{a_q^\dagger a_s} a_r \\ &+ a_p^\dagger \overline{a_q^\dagger a_s} a_r + \overline{a_p^\dagger a_q^\dagger} a_s a_r + \overline{a_p^\dagger a_q^\dagger} a_s a_r + \overline{a_p^\dagger a_q^\dagger} a_s a_r + \overline{a_p^\dagger a_q^\dagger} a_s a_r \end{aligned}$$

For this spesific case we will substitute $\frac{1}{4} \langle pq | \hat{v} | rs \rangle$ with $-\frac{g}{2} \delta_{pq} \delta_{rs}$. The interaction term can now be seperated into three parts:

- 1. The two body term, which arrises from the normal ordering of the creation and annihilation operators.
- 2. The one body term, which comes from the single contractions.
- 3. The constant term, which comes from the double contractions.

From this we can construct a new normal ordered interaction term:

$$\hat{H}_I = -\frac{g}{2} \sum_{pq} \{a_p^\dagger a_q^\dagger a_q a_p\} - \frac{g}{2} \sum_{pq} a_p^\dagger a_{p\sigma} a_{p\sigma} + -\frac{g}{2} (\delta_{pr} \delta_{qs} + \delta_{ps} \delta_{qr})$$

We can combine the one-body term into the first one we found, and we get:

$$\hat{H}_0^N = \sum_{p\sigma} (p-1) \{a_p^\dagger a_{p\sigma}\} - \frac{g}{2} \sum_{pq} a_p^\dagger a_{p\sigma} a_{p\sigma}$$

The normal ordered interaction term is:

$$\hat{V}^N = -\frac{g}{2} \sum_{pq} \{a_p^\dagger a_q^\dagger a_q a_p\}$$

and the reference energy:

$$E_0^{ref} = \langle \Phi_0 | \hat{H}_0 | \Phi_0 \rangle - \frac{g}{2} \sum_{pq} (\delta_{pr} \delta_{qs} + \delta_{ps} \delta_{qr}) = 2 - g$$

where the values were obtained in the previous task.

The total normal ordered Hamiltonian is:

$$\hat{H}^N = \hat{H}_0^N + \hat{V}^N + E_0^{ref}$$

We now want to set up the standard Hartree Fock equations for the system, also known as the canonical Hartree-Fock equations. We begin by defining the single-particle operator \hat{f} , which for a general case is given by:

$$\langle p | \hat{f} | q \rangle = \langle p | \hat{H}_0 | q \rangle + \sum_j \langle pj | \hat{V} | qj \rangle_{AS}$$

where the subscript AS indicates that the matrix elements are antisymmetrized. Due to the nature of the Hamiltonian, we have $\langle p | \hat{f} | q \rangle = 0$ for $p \neq q$. In second quantization we can write the single-particle operator as:

$$\hat{F} = \sum_{pq} \langle p | \hat{f} | q \rangle a_p^\dagger a_q = \sum_p \langle p | \hat{f} | p \rangle a_p^\dagger a_p$$

In writing the operator in normal order, we have:

$$\overline{a_p^\dagger a_p} = \{a_p^\dagger a_p\} + \delta_{pp}$$

and thus:

$$\hat{F} = \sum_p \langle p | \hat{f} | p \rangle \{a_p^\dagger a_p\} + \sum_p \langle p | \hat{f} | p \rangle$$

The canonical Hartree-Fock equations are given by:

$$\hat{f} | p \rangle = \epsilon_p | p \rangle$$

where ϵ_p is the single-particle energy.

f)

g)