How to prove the arithmetic mean and geometric mean formula

hisanobu-nakamura

Title: A Proof for Arithmetic Mean and Geometric Mean Formula Date: 2020-02-10 Category: math Tags: math

1 Arithmetic Mean and Geometric Mean Formula

1.1 The Statement

Here is the arithmetic and geometric formula

$$\frac{a_1 + \dots + a_n}{n} \ge \sqrt[n]{a_1 \cdots a_n},$$

where $\forall i \ a_i > 0$.

1.2 Proof of the Statement

OK for n = 1. Equality holds. Suppose that the inequality holds for all natural numbers up to n, then

$$a_{1} + \dots + a_{n+1} = \frac{a_{2} + a_{3} \dots + a_{n+1}}{n} + \frac{a_{1} + a_{3} + \dots + a_{n+1}}{n} + \dots$$

$$+ \frac{a_{1} + a_{2} + \overset{i}{\overset{i}{\overset{i}{\smile}}} + a_{n+1}}{n} + \dots + \frac{a_{1} + \dots + a_{n}}{n}$$

$$= \sqrt[n]{a_{2} \dots a_{n+1}} + \dots + \sqrt[n]{a_{1} \dots a_{n}}$$

$$= \sqrt[n+1]{a_{1} \dots a_{n+1}} \left(\frac{\sqrt[n]{a_{2} \dots a_{n+1}}}{\sqrt[n+1]{a_{1} \dots a_{n+1}}} + \dots + \frac{\sqrt[n]{a_{1} \dots a_{n}}}{\sqrt[n+1]{a_{1} \dots a_{n+1}}} \right)$$

$$= \sqrt[n+1]{a_{1} \dots a_{n+1}} \left(\sqrt[n+1]{\frac{n+1}{a_{1}}} + \dots + \sqrt[n+1]{\frac{n+1}{a_{n+1}}} + \dots + \sqrt[n+1]{\frac{n}{a_{n+1}}} \right)$$

$$= \sqrt[n+1]{a_{1} \dots a_{n+1}} \left(\sqrt[n+1]{\frac{\alpha_{1}}{a_{1}}} + \dots + \sqrt[n+1]{\frac{\alpha_{n+1}}{a_{n+1}}} \right)$$

where $\alpha_i = \sqrt[n]{a_1 \cdot \ddot{\cdot} \cdot a_{n+1}}$ (Here, $\dot{\cdot} \dot{\cdot}$ means omitting the i-th term in the sum or the product). It is easy to see that

$$\sqrt[n+1]{\frac{\alpha_1}{a_1}} \cdots \sqrt[n+1]{\frac{\alpha_{n+1}}{a_{n+1}}} = 1.$$

So, if we can show the following proposition, we are done.

Lemma: If $b_1, \dots, b_{n+1} > 0$ and $b_1 \dots b_{n+1} = 1$, then

$$b_1 + \dots + b_{n+1} \ge n+1 \tag{1.1}$$

Now, by substituting $b_i = \sqrt[n+1]{\frac{\alpha_i}{a_i}}$, it is straightforward to see that

$$a_{1} + \dots + a_{n+1} = \sqrt[n+1]{a_{1} \cdots a_{n+1}} \left(\sqrt[n+1]{\frac{\alpha_{1}}{a_{1}}} + \dots + \sqrt[n+1]{\frac{\alpha_{n+1}}{a_{n+1}}} \right)$$

$$\geq \sqrt[n+1]{a_{1} \cdots a_{n+1}} (n+1)$$

1.3 Proof of the Lemma

Lemma: If $b_1, \dots, b_{n+1} > 0$ and $b_1 \dots b_{n+1} = 1$, then

$$b_1 + \dots + b_{n+1} \ge n+1 \tag{1.2}$$

Proof: First, if $b_1 = \cdots = b_{n+1} = 1$, then the equality holds. We notice that $\exists i$ such that $b_i > 1$ implies that $\exists j$ such that $b_j < 1$. So, let us assume the condition and proceed to prove the statement by induction. For n = 1, we can assume that $b_1 = 1 - c_1$ with $0 < c_1 < 1$. Then

$$b_1 + b_2 = b_1 + \frac{1}{b_1} = 1 - c_1 + \frac{1}{1 - c_1} \ge 1 - c_1 + 1 + c_1 = 2.$$

Now, let us suppose that (1.2) holds true for all the natural numbers up to n. Regarding the fact mentioned above, let us suppose that $b_n = 1 + c_n$ and $b_{n+1} = 1 + c_{n+1}$ with $c_n < 0$ and $c_{n+1} > 0$. Then, by the assumption,

$$(b_1 \cdots b_{n-1})(b_n b_{n+1}) = 1$$
 implies $b_1 + \cdots + b_{n-1} + b_n b_{n+1} \ge n$.

If we can say

$$b_1 + \dots + b_{n-1} + (b_n + b_{n+1}) \ge \text{or} > b_1 + \dots + b_{n-1} + (b_n b_{n+1} + 1),$$

then the inequality follows. But

$$b_n + b_{n+1} - (b_n b_{n+1} + 1) = -c_n c_{n+1} > 0.$$

Hence, it follows that

$$b_1 + \dots + b_{n-1} + b_n + b_{n+1} > b_1 + \dots + b_{n-1} + b_n b_{n+1} + 1 > n+1$$