

Name:	ID:	Section:
-------	-----	----------

Question 1 [C01] [10 Points]

Take K daily valid rainfall measurements (in millimeters). A rainfall measurement is **invalid** if the value is **below 0 or above 200**. For such inputs display: "Invalid rainfall value, rainfall measurement". Continue to take rainfall values from the user until you have K number of valid inputs. Finally, calculate a **Rainfall Score** using the following rules:

- ❖ **Add** the rainfall value if,
 - value divisible by 3
 - value is a multiple of 7
- ❖ **Otherwise, subtract** the rainfall value.

Count the number of odd digits in the Final Rainfall Score. [Before counting, check if the score is negative. If it is, multiply it by -1.]

Sample Input	Sample Output
Enter K: 5 Enter rainfall 1: 12 Enter rainfall 2: 5 Enter rainfall 3: 28 Enter rainfall 4: 25 Enter rainfall 5: 212 Enter rainfall 5: 49	Invalid rainfall value, 212. Final Rainfall Score: 59 Number of odd digits: 2
Explanation: 6 rainfall values taken as one was invalid (212). Rainfall score = $12 - 5 + 28 - 25 + 49 = 59$ The odd digit of the Temperature score is 5 & 9 so, the number of odd digits = 2.	