MDP Project

Review 2 Package

Rough draft • Pseudocode • Software justification • Execution

Built against Review 1 feedback: be relevant to the research literature, show execution and a rough draft, add pseudocode, and justify every tool. All numbers come from the committed code on the real 204-station dataset; the framing is deliberately honest and narrowed where the evidence does not support stronger claims.

Part 1 — Rough Draft (Research Paper)

Rough Draft -- Research Paper

> ***Status:*** *rough draft for Review 2. Every number is produced by the committed code on the real dataset (see* `audit/reports/`*). Framing is deliberately honest and narrowed -- see §VIII Limitations and the claim-to-evidence table. Citations are real, foundational works (see* `audit/manuscript/BIBLIOGRAPHY.md`*); DOIs are to be confirmed against publisher records before any submission.*

Title

An Open, Leakage-Safe Benchmark for Multi-Pollutant Air-Quality Machine Learning over 204 Indian Monitoring Stations

Alternative title: What Can and Cannot Be Learned from Pooled Regulatory Air-Quality Data -- A Leakage-Safe Machine-Learning Case Study over 204 Indian Monitoring Stations.

> *We moved away from the original title *"Causal Machine Learning for Multi-Pollutant Source Apportionment and Control Optimization"* because our own evaluation could not support the words causal, source apportionment, or control optimization (§VIII). We reframe the work as a **benchmark and resource contribution**: an open dataset, a leakage-safe evaluation protocol, and honest baselines others can build on. Narrowing the claim rather than overstating it is itself the contribution.*

Keywords: air quality; open benchmark; machine learning; data leakage; causal discovery; DirectLiNGAM; XGBoost; multi-objective optimization; reproducibility; India.

Abstract

We release an **open benchmark** for multi-pollutant air-quality machine learning: one year (2025) of hourly observations from 204 Indian monitoring stations (OpenAQ) merged with co-located modelled meteorology (Open-Meteo) -- about 1.18 million station-hours -- together with a **leakage-safe evaluation protocol** and honest baselines. Measurement units are **verified against the authoritative OpenAQ sensor metadata**: NO, ozone, PM2.5 and PM10 are unit-consistent, while CO, NO2 and SO2 carry duplicate ppb/µg/m3 sensors and are treated as unit-indeterminate. Under leakage-safe evaluation, a gradient-boosted model estimates contemporaneous ozone with held-out R2 ~ 0.25 (blocked-temporal) to 0.30 (unseen stations), well below the 0.41-0.43 from a leaky random split; pollutants add only ~ +0.12 R2 *beyond weather*, and a one-hour-ahead forecast beats a persistence baseline by only +0.05. Exploratory causal-structure discovery (DirectLiNGAM) recovers a single bootstrap-stable association consistent with local NO-O3 titration, but the wider graph is confounded by omitted meteorology and is not causally identified. K-means groups co-varying species only weakly (silhouette 0.26-0.39) and is not source apportionment. A linear response model admits a closed-form scenario frontier; a published metaheuristic "optimization" returned only dominated points. We conclude that the engineering is sound and reproducible, but that causal, apportionment and policy claims are unsupported by these data, and we specify the additional evidence each would require.

I. Introduction

Urban air quality in India is a major public-health problem, and open monitoring data invite machine-learning analysis. A tempting narrative runs *prediction -> causal structure -> policy optimization*. Many applied papers present each link as established. We instead **test** that narrative critically on a large but imperfect regulatory dataset and report, with uncertainty, where it holds and where it breaks.

Contributions. (1) An **open, reproducible dataset + PySpark/Dataproc pipeline** over 204 stations, with units verified against authoritative sensor metadata. (2) A **leakage-safe evaluation protocol** and honest baselines others must beat, showing how random splitting inflates apparent skill. (3) One stable, chemically-plausible association (NO-O3) reported as a hypothesis, not an identified effect. (4) A demonstration that a published metaheuristic "optimizer" was dominated by a closed-form solution. (5) An explicit account of the evidence each stronger claim would need.

Significance -- who this helps. The work is useful to the air-quality and environmental-AI community in three concrete ways: a real, open, reproducible **dataset and pipeline** others can build on; a reusable **leakage-safe evaluation recipe and diagnostic set** (bootstrap edge stability, meteorology-confounding tests, exact-vs-metaheuristic checks) that this field often omits; and an honest, evidence-backed **account of what imperfect regulatory data can and cannot support**, including a testable NO-O3 hypothesis and a requirements list for causal source apportionment. A full area-by-area breakdown is in `audit/review2/SIGNIFICANCE.md`.

II. Related Work (positioning)

Our methods are standard; our contribution is their *honest application and evaluation*. We position against five strands:

- **Tabular prediction / boosting.** XGBoost [Chen & Guestrin, 2016] is the canonical tabular regressor; we use it but subject it to leakage-safe evaluation rather than reporting a single random-split R2.
- **Data leakage.** [Kaufman et al., 2012] formalizes leakage; air-quality ML frequently random-splits autocorrelated time series. Our blocked-temporal and leave-stations-out protocol is the direct remedy.
- **Causal discovery.** DirectLiNGAM [Shimizu et al., 2006; 2011] and Pearl's identification framework [Pearl, 2009] let us *orient* edges; we add bootstrap stability and a meteorology-confounding test, which most applied uses omit.
- **Source apportionment.** Receptor models, esp. PMF [Paatero & Tapper, 1994], require chemical speciation and per-measurement uncertainties. Our data have neither, so we explicitly decline the apportionment claim.
- **Meteorological normalization.** [Grange & Carslaw, 2019] show weather must be removed before attributing air-quality changes -- motivating our weather-vs- pollutant decomposition and our refusal to read the raw graph as mechanism.

How our paper "matches" the literature. It uses the same methods as published causal-ML air-quality studies, but reports the diagnostics those studies usually skip (leakage-safe splits, bootstrap edge stability, confounding tests, an exact optimality check). The comparison table in `audit/manuscript/ORIGINAL_VS_CORRECTED.md` maps each original claim to its corrected, evidence-backed version.

III. Data

Hourly CO, NO, NO2, O3, PM10, PM2.5 and SO2 for 204 Indian stations with 2025 data (OpenAQ v3), each merged with hourly Open-Meteo reanalysis (temperature, relative humidity, wind speed, surface pressure) at the nearest ~0.1deg grid cell. Scale: **1,180,551 station-hours**, ~ 66% of a full 204 x 8,760 grid; 37 stations < 50% within-year coverage. **Units were verified against the authoritative OpenAQ per-sensor metadata:** NO (ppb), O3, PM2.5 and PM10 (µg/m3) are unit-consistent, whereas CO, NO2 and SO2 carry duplicate ppb *and* µg/m3 sensors at almost every station (e.g. CO at 183/198) with no per-row provenance, so their stored unit is **indeterminate**; we therefore restrict quantitative cross-station use to the unit-consistent pollutants (Fig. 6). Further caveats: 35 non-physical outliers and saturation caps exist; weather is modelled; the network is

urban and Delhi-NCR-heavy (not nationally representative); VOCs are structurally absent.

IV. Methodology

Full algorithmic detail is in `audit/review2/PSEUDOCODE.md`; the tool rationale is in `SOFTWARE_STACK.md`. Four analyses:

1. **Predictive baseline** -- XGBoost estimates same-hour ozone from pollutants + weather. Evaluated with leaky random (contrast only), blocked-temporal, leave-stations-out, and strict splits; preprocessing fit on train only; hyper-parameters fixed a priori; baselines = mean, climatology, persistence; uncertainty via station-block bootstrap.
2. **Structure discovery** -- DirectLiNGAM on standardized pollutants with a prior forbidding secondary->primary edges; bootstrap stability + meteorology-inclusion sensitivity.
3. **Species grouping** -- K-means on inter-species correlation profiles; silhouette model-selection + bootstrap co-assignment.
4. **Scenario analysis** -- the linear model's exact Pareto frontier derived in closed form and compared to the published metaheuristic result.

V. Experimental Setup & Execution

- **Pipeline execution.** The full three-phase pipeline runs end-to-end on Google Cloud Dataproc (custom Docker runtime) and in Cloud Shell; `main.py --phases all` completes with every phase reporting `all_ok` on the real 204-station data.
- **Audit execution.** Eight self-contained scripts (`audit/*.py`) regenerate every number offline with `SEED=42` (Track A of `RUNBOOK.md`); `pytest` guards pass.
- **Compute.** Prediction and causal-stability scripts run in ~ 2 min each on a single machine; bootstrap counts are CLI flags.

VI. Results

Figures (in `audit/reports/figures/`, regenerable via `audit/make_figures.py`): Fig. 1 data coverage; Fig. 2 R2 by evaluation protocol (leakage); Fig. 3 incremental value of pollutants beyond weather; Fig. 4 bootstrap edge stability; Fig. 5 Pareto dominance of the 19 saved policies; Fig. 6 authoritative units verification.

A. Prediction. Leakage inflates skill: random split R2 ~ 0.41, but honest evaluation gives R2 ~ 0.25 (blocked-temporal; 90% CI 0.22-0.28) and 0.30 (unseen stations; CI 0.19-0.40), down to 0.22 strict. Decomposition (Phase C2): weather alone R2 = 0.14, pollutants add **+0.12** (to 0.26), time harmonics -> 0.34; the 0.41 is only recovered by a train-only per-station mean, which is **site memorisation**, not generalisation. Persistence reaches R2 ~ 0.79-0.81 for gap-filling; a true t+1h forecast beats persistence by only **+0.05** (0.81 -> 0.87).

B. Structure. One edge is bootstrap-stable and survives meteorology: **NO -> O3, negative** (frequency 1.0, -0.138 ± 0.007; -0.108 with weather) -- local titration. PM2.5 edges collapse once weather enters (temp -> PM2.5, humidity -> O3 dominate): the pollutant-only graph was **confounded**. SO2 edges are unstable (freq <= 0.5).

C. Grouping. Silhouette weak (0.26 raw, 0.39 imputed). Core groups {NO, NO2, PM10, PM2.5} and {O3, SO2} are stable; CO's instability was an **outlier artefact** -- after flagging 35 values, CO-NO rises 0.14 -> 0.42 and CO joins the combustion group at co-assignment 1.0. Still a weak convenience partition, now physically coherent (caveat: CO and NO2 are unit-indeterminate, so their grouping may partly reflect the ppb/µg/m3 mix; the stable {O3, SO2} vs combustion split does not depend on CO).

D. Scenarios. The frontier is a short near-linear trade-off driven almost entirely by NO. The published metaheuristic's **19 "optimal" policies are all dominated** by the exact frontier (mean PM2.5 gap 0.046, max 0.298 standardized).

VII. Discussion

The honest contribution is modest and methodological: a correctly-evaluated predictive baseline that beats spatial baselines but not temporal persistence; one stable, plausible association reported as a hypothesis; a demonstration that a published "optimization" was dominated by a closed-form solution; and a cautionary account of how leakage, omitted confounders, unharmonised units and pooling inflate apparent findings. SHAP-DirectLiNGAM agreement on the NO sign is internal consistency, **not** independent validation.

VIII. Limitations

Units unharmonised (CO); weather modelled not observed; urban non-representative network; ~ 66% completeness with mixed missingness; VOC absent; a single pooled *linear* contemporaneous model cannot represent regime-dependent (VOC- vs NOx-limited) ozone chemistry; "interventions" act on standardized ambient concentrations, not emissions. **None of the analyses identify causal effects.**

What each stronger claim would require: *causal* -> an identification strategy (natural experiment / policy discontinuity) and measured confounders; *source apportionment* -> chemical speciation + per-measurement uncertainties + VOCs and a receptor model (PMF); *control optimization* -> a validated emissions->concentration model, not a fit to ambient levels.

IX. Conclusion & Future Work

The pipeline runs, is reproducible, and the dataset is real; but its original headline claims do not survive scrutiny. Reframed as an exploratory study with leakage-safe evaluation, explicit uncertainty, and honest limitations, it is a useful cautionary contribution. Future work: verify units against OpenAQ sensor metadata; obtain speciated data for apportionment; seek a genuine intervention for causal identification; and extend the predictive model to multi-step forecasting.

References

See `audit/manuscript/BIBLIOGRAPHY.md` (Shimizu et al. 2006/2011; Pearl 2009; Chen & Guestrin 2016; Lundberg & Lee 2017; Deb et al. 2002; Akiba et al. 2019; Zaharia et al. 2016; Kaufman et al. 2012; Seinfeld & Pandis 2016; Paatero & Tapper 1994; Grange & Carslaw 2019; OpenAQ; Open-Meteo). **DOIs/identifiers are drafted there** (resolve each at doi.org); confirm every one before submission.

Part 2 — Significance: How This Helps the Field

Significance -- How This Project Is Useful to the Field

This answers the Review 1 point directly: *the thing we are making should be helpful / come in use to the research area that the papers we read are on.* Below, for each research area those papers belong to, is **what our project contributes to that area** and **who it helps**. Everything here is backed by the committed code and the audit (`audit/reports/`), so each claim of usefulness is itself evidenced.

The honest headline: our strongest contributions are a **reusable dataset + pipeline**, a **leakage-safe evaluation recipe**, and a **cautionary map of the traps** in this kind of study. A negative/methodological result that stops other researchers from overclaiming is a real contribution to the area -- not a gap.

Area 1 -- Air-quality prediction & forecasting with ML

Papers in this area ask: can we predict pollutant levels from other pollutants and weather, and how well?

What we add / who it helps:

- A **reproducible benchmark** on a real 204-station Indian dataset (~1.18M station-hours) that anyone in the area can reuse, with fixed seeds and splits.
- An **honest skill ceiling**: contemporaneous ozone R2 ~ 0.25-0.30 held-out, and pollutants add only ~ +0.12 beyond weather. This tells other groups *what good actually looks like* on this data, so they don't chase inflated random-split numbers (Phase C / C2).
- A practical note for **monitoring-network operators**: persistence is a hard baseline for temporal gap-filling (R2 ~ 0.8), so model value lies in *spatial* gap-filling and multi-step forecasting, not next-hour nowcasting.

Area 2 -- Causal inference / causal discovery in environmental science

Papers in this area ask: can we recover *directional* relationships (not just correlations) among pollutants from observational data?

What we add / who it helps:

- A **worked, honest template** for applying DirectLiNGAM with the diagnostics most applied papers skip: bootstrap edge stability **and** a meteorology-confounding test. Other researchers can lift this protocol directly.
- A concrete, **testable hypothesis** for atmospheric chemists: a stable negative NO -> O3 association consistent with local titration (survives weather) -- a lead worth following with a designed study.
- A clear **requirements list** for anyone wanting a *causal* claim here: an identification strategy (natural experiment / policy discontinuity) and measured confounders. This saves the field effort by naming what's missing (Phase D).

Area 3 -- Source apportionment / receptor modelling

Papers in this area ask: which sources contribute how much to measured pollution (e.g. PMF, CMB)?

What we add / who it helps:

- A **precise statement of data requirements** -- chemical speciation, per-measurement uncertainties, and VOCs -- that regulatory ML datasets like ours lack. This is directly useful to **data providers and monitoring agencies** deciding what to measure so their data *can* support apportionment.
- Evidence that **K-means grouping is not apportionment**, with the silhouette / co-assignment numbers to prove it (Phase E) -- a caution that stops the common misread of clustering as source attribution.

Area 4 -- Multi-objective optimization for environmental policy

Papers in this area ask: how to trade off reductions across pollutants/costs (often with NSGA-II or similar metaheuristics)?

What we add / who it helps:

- A reusable **sanity check**: when the response model is linear, the exact Pareto front is closed-form -- and in our case it **dominated all 19 metaheuristic "optimal" policies** (Phase B/F). The lesson for the area: *always compare your metaheuristic against an exact or analytic baseline when one exists.*
- A caution for **policy modellers**: optimizing over *ambient concentrations* is not the same as optimizing *emissions*; the mapping must be validated first.

Area 5 -- Explainable AI (XAI) in environmental applications

Papers in this area ask: can SHAP/feature attributions explain and even "validate" model behaviour?

What we add / who it helps:

- A clear demonstration that **SHAP agreeing with a causal-discovery result is internal consistency, not independent validation** -- a precise caveat the XAI community can cite when reviewing such claims.

Area 6 -- Reproducibility & open data in environmental monitoring

Papers in this area ask: are results reproducible, and is the data/code open?

What we add / who it helps:

- A fully **open, versioned pipeline + dataset + audit** (branch `review/scientific-audit`, `RUNBOOK.md`) that runs offline with one command set -- a ready template for **students and reviewers** to reproduce and extend.
- A documented **leakage-safe evaluation recipe** (blocked-temporal, leave-stations-out, station-block bootstrap) reusable by any air-quality ML study.

One-paragraph version (for the review)

Our project is useful to the air-quality / environmental-AI research area in three concrete ways: (1) it contributes a **real, open, reproducible dataset and pipeline** others can build on; (2) it provides a **leakage-safe evaluation recipe and a set of diagnostics** (bootstrap edge stability, meteorology-confounding tests, exact-vs-metaheuristic checks) that this field often omits; and (3) it delivers an **honest, evidence-backed account of what these data can and cannot support**, including a testable NO-O3 hypothesis and a clear list of what causal source-apportionment would require. In a field prone to overclaiming from imperfect regulatory data, a careful cautionary study is a genuine service to the area.

Part 3 — Pseudocode

Pseudocode -- MDP Air-Quality ML Pipeline

Algorithm-level pseudocode for every stage. Each block mirrors the actual code (`data_collection.py`, `src/phases/`, `audit/`) but is written to be read and defended in a review, not executed. Notation: `D` = dataset (rows = station-hours, columns = pollutants + weather), `S` = set of stations, `u,sigma` = per-column train-fold mean/std.

Algorithm 0 -- Data Collection (`data_collection.py`)

```
INPUT:  country = India, year = 2025, pollutants P = {CO,NO,NO2,O3,PM10,PM25,SO2}
OUTPUT: one Parquet file per station in data/raw/telemetry/

for each station s in OpenAQ(country, has_data_in=year):          # v3 API
    sensors  <- GET /locations/{s}/sensors                        # capture units!
    meas     <- GET /measurements for each sensor, hourly, year
    weather  <- Open-Meteo reanalysis at nearest 0.1 deg grid cell of s
                for {temperature, relative_humidity, wind_speed, pressure}
    table    <- OUTER-JOIN(meas, weather) ON timestamp (hourly)
    write Parquet(table) with station_id, lat, lon, units-manifest
end
merge per-station files -> 204 files, ~1.18M station-hours
```

Design note: units are stored per sensor so that comparability can be *verified* later (the original collection discarded them -- a defect the audit fixed).

Algorithm 1 -- Phase 1: Species Grouping by K-means (`phase1_clustering.py`)

```
INPUT:  D, species columns C, k (default 3)
OUTPUT: cluster label per species + silhouette score

# features are per-species correlation PROFILES, not raw concentrations
R <- pairwise_correlation_matrix(D[C])          # |C| x |C|
X <- rows of R                                  # each species -> vector of its
                                                # correlations with all others
for k' in candidate range:                      # model-selection
    model   <- KMeans(k=k', seed=42).fit(X)
    score   <- silhouette(X, model.labels)
choose k with best silhouette
RETURN model.labels, score
```

Why correlation profiles: species that co-vary with the *same* other species are grouped, which is the closest K-means can get to "behaves like a common source".

Algorithm 2 -- Phase 2a: Predictive Baseline (`phase2_causal.py`, XGBoost)

```
INPUT:  D, target y = ozone, features F = other pollutants + weather
OUTPUT: fitted model + HONEST held-out skill

# --- leakage-safe splitting (the core correctness fix) ---
TRAIN, TEST <- blocked_temporal_split(D, train = months 1..9, test = 10..12)
          OR  leave_stations_out(D, S)          # unseen-station generalisation

mu, sigma   <- mean,std of F over TRAIN only    # fit preprocessing on train only
TRAIN, TEST <- standardize(F, mu, sigma)

model <- XGBoostRegressor(n_estimators=250, max_depth=6, lr=0.05,
                          subsample=0.8, tree_method=hist, seed=42)   # fixed a priori
model.fit(TRAIN[F], TRAIN[y])

R2 <- r2_score(TEST[y], model.predict(TEST[F]))
```

```
# baselines for honest comparison
R2_persist  <- r2(TEST[y], previous_hour_value)
R2_clim     <- r2(TEST[y], per_station_mean(TRAIN))
CI          <- station_block_bootstrap(R2)        # uncertainty
RETURN model, R2, R2_persist, R2_clim, CI
```

The *a priori* hyper-parameters and train-only preprocessing are what make the reported R2 trustworthy; tuning on TEST would re-introduce leakage.

Algorithm 3 -- Phase 2b: Causal Structure Discovery (DirectLiNGAM)

```
INPUT:  standardized pollutants Z, prior K (forbid secondary -> primary edges)
OUTPUT: adjacency matrix + per-edge bootstrap stability

model <- DirectLiNGAM(prior_knowledge = K).fit(Z)
A     <- model.adjacency_matrix                  # weighted DAG among species
persist manifest{columns, mu, sigma, A, threshold, seed, versions}

# stability + confounding check (the part that makes it defensible)
for b in 1..B:                                   # station-block bootstrap
    Z_b   <- resample stations with replacement
    A_b   <- DirectLiNGAM(K).fit(Z_b).adjacency
    tally edge frequencies and sign
re-fit WITH weather nodes {temp,RH,wind,pressure} as exogenous
RETURN edges with frequency, mean weight, and weather-sensitivity flag
```

An edge is reported only if it is bootstrap-stable **and** survives the inclusion of meteorology (otherwise it was confounded).

Algorithm 4 -- Phase 3a: Multi-Objective Policy Search (Optuna NSGA-II)

```
INPUT:  response model f (ozone, PM2.5 vs standardized concentration levers x)
OUTPUT: Pareto set of trade-off policies

study <- Optuna.create_study(directions=[minimize,minimize],
                             sampler = NSGAIISampler(seed=42))   # NOT the TPE default
for trial in 1..N:
    x       <- trial.suggest levers over supported variables only   # reject VOC etc.
    (o, pm) <- f.evaluate(x)                      # raise if x uses an unfitted var
    report (o, pm)
RETURN study.best_trials                           # non-dominated front
```

Guard (audit fix): decision variables are intersected with the *fitted* graph; asking to move a variable the model never learned raises, instead of silently returning a meaningless "optimum".

Algorithm 4' -- Exact frontier (audit check, `analytic_pareto.py`)

```
# because f is LINEAR, the true Pareto front is closed-form
front <- { x on the boundary of the feasible box minimizing lambda*o+(1-lambda)*pm
           for lambda in [0,1] }
for each saved metaheuristic policy p:
    assert exists q in front with q dominates p     # all 19 were dominated
```

Algorithm 5 -- Phase 3b: Explanation (SHAP)

```
INPUT:  fitted XGBoost model, background sample B
OUTPUT: per-feature attributions

explainer <- TreeSHAP(model)
phi       <- explainer.shap_values(B)              # exact for trees
rank features by mean(|phi|); inspect NO contribution to ozone
```

SHAP agreeing with DirectLiNGAM on the NO sign is **internal consistency** of two models on the same data -- reported as such, not as independent validation.

Part 4 — Software & Tools: What and Why

Software & Tools -- What We Use and Why It Matters

For each component: **what it is**, **why we chose it over alternatives**, and **where it is used** in the pipeline. This answers the review requirement to "explain the importance of each code and software, and why we are using it."

Infrastructure / Big-data engine

Apache Spark (PySpark)

- **What:** distributed in-memory data-processing engine with a Python API.
- **Why it matters:** the dataset is ~1.18M station-hours across 204 files; Spark lets the same code run on a laptop and scale out on a cluster without rewriting. Its DataFrame API and lazy execution plan avoid loading everything into RAM.
- **Why not pandas alone:** pandas is single-machine and would not scale to the full multi-city/multi-year target; Spark is the standard for this scale.
- **Used in:** all three pipeline phases (`src/phases/`), driven by `main.py`.

Google Cloud Dataproc

- **What:** managed Spark/Hadoop clusters on Google Cloud.
- **Why it matters:** gives a reproducible, disposable cluster so a reviewer can re-run the exact pipeline; billed per-second, so experiments are cheap and nothing persists accidentally.
- **Why not a static VM/Hadoop install:** no cluster to maintain; the custom runtime image (`Dockerfile`) pins every dependency version for reproducibility.
- **Used in:** `submit.sh`, `cloudbuild.yaml`, `Dockerfile` (deployment).

Parquet (via PyArrow)

- **What:** columnar, compressed, typed on-disk format.
- **Why it matters:** column pruning + compression make reading a few pollutants out of 204 files fast; the embedded schema prevents silent type drift.
- **Why not CSV:** CSV loses types and is far larger/slower; a real bug we hit (int-vs-float humidity) is exactly what a typed format guards against.

Machine-learning methods

K-means clustering (Spark MLlib)

- **What:** partitions items into `k` groups by minimizing within-cluster variance.
- **Why it matters:** a transparent, fast first pass at "which species behave alike"; the silhouette score gives an objective number of groups.
- **Why not DBSCAN/hierarchical here:** we cluster a handful of species on correlation profiles -- K-means with silhouette model-selection is simple, deterministic (seeded) and defensible; it is **grouping, not apportionment**.
- **Used in:** Phase 1 (`phase1_clustering.py`).

XGBoost (SparkXGBRegressor)

- **What:** gradient-boosted decision trees -- a strong tabular regressor.
- **Why it matters:** air-quality relationships are nonlinear and interacting; boosted trees capture these without manual feature engineering and are the consistent top performer on tabular benchmarks.

- **Why not linear regression / deep nets:** linear underfits the chemistry; deep nets need far more data and tuning and rarely beat boosting on tabular data.
- **Used in:** Phase 2a predictive baseline, and the SHAP explanation.

DirectLiNGAM (`lingam`)

- **What:** learns a linear, non-Gaussian acyclic causal structure (a DAG) from observational data.
- **Why it matters:** it is one of the few methods that can *orient* edges (not just correlate) under its assumptions, letting us pose directional hypotheses such as NO -> ozone.
- **Why we are cautious:** it assumes linearity, non-Gaussianity, acyclicity and **no hidden confounders** -- we explicitly test the last by adding meteorology, which is why we report only the one edge that survives.
- **Used in:** Phase 2b causal discovery.

Optuna + NSGA-II sampler

- **What:** Optuna is an optimization framework; NSGA-II is a multi-objective evolutionary algorithm that returns a Pareto front, not one answer.
- **Why it matters:** policy involves trade-offs (lower ozone vs lower PM2.5); NSGA-II surfaces the whole trade-off surface instead of a single compromise.
- **Why the sampler choice is critical:** Optuna defaults to TPE, which is single-objective; we must pass `NSGAIISampler` explicitly (a bug the audit fixed). For our *linear* model we also derived the exact front in closed form to check the metaheuristic -- and it was dominated.
- **Used in:** Phase 3a optimization + `analytic_pareto.py` cross-check.

SHAP (TreeExplainer)

- **What:** Shapley-value feature attributions; exact and fast for tree models.
- **Why it matters:** turns the black-box XGBoost into per-feature, per-prediction explanations with a sound game-theoretic basis -- needed for any "XAI" claim.
- **Why not plain feature importance:** gain/weight importances are inconsistent and global; SHAP is locally accurate and additive.
- **Used in:** Phase 3b explainability.

Data sources

OpenAQ v3 API

- **What:** open aggregator of government/reference air-quality monitors.
- **Why it matters:** real, citable regulatory measurements for India at hourly resolution -- no fabrication, and reproducible by any reviewer with a free key.
- **Caveat we surface:** per-sensor units must be captured to compare stations.

Open-Meteo reanalysis

- **What:** free historical/reanalysis weather API.
- **Why it matters:** ozone is weather-driven; we need co-located temperature, humidity, wind and pressure. Reanalysis gives complete, gap-free coverage.
- **Caveat we surface:** weather is **modelled**, not measured on-site.

Engineering & reproducibility

Tool	Why it matters
Git + feature branch	Raw data and original results preserved; all audit work isolated on `review/scientific-audit`.

Tool	Why it matters
Docker (custom image)	Pins every library version so the cluster run is byte-for-byte reproducible.
Fixed seeds (`SEED=42`**)**	Every clustering/boosting/bootstrap run is deterministic and re-checkable.
scikit-learn / SciPy / statsmodels	Standard, audited implementations of metrics, splits, bootstrap and stats used throughout the audit.
Matplotlib	Publication figures (e.g. the scenario frontier) generated from the same data.
pytest	Encodes correctness guards (e.g. "optimizer front truly dominates") as tests that must pass.

One-line summary of the stack choice: every tool is the *standard, citable, reproducible* option for its job -- Spark/Dataproc for scale, Parquet for typed storage, XGBoost for tabular prediction, DirectLiNGAM for directional hypotheses, NSGA-II for trade-offs, SHAP for explanations -- and each is paired with an honest caveat about what it can and cannot establish on this dataset.

Part 5 — Publication Plan

Publication Plan -- From Here to a Real Paper

Honest, prioritized roadmap. Status markers: [x] done - [~] in progress - [ ] to do - [YOU] needs the team (can't be automated). Based on `audit/manuscript/READINESS.md`.

Step 0 -- Decide which paper (do this first, with mam)

	Path A -- Open benchmark + honest analysis (RECOMMENDED)	Path B -- Original causal/apportionment paper
Claim	Open dataset + leakage-safe evaluation + honest baselines + cautionary findings	Causal source-apportionment + policy optimization
Feasible on our data?	**Yes**	**No** -- needs speciation, VOCs, an intervention, emissions data we don't have
Effort	~1-2 focused weeks (science mostly done)	Months + new data collection; may be infeasible as a student project
Venue fit	Workshop / conference / benchmark track	Atmospheric-science journal (very high bar)

Recommendation: Path A. It makes our honesty a strength and is achievable now. Everything below assumes Path A.

The contribution, in one sentence

> *An open, reproducible, **leakage-safe benchmark** for multi-pollutant air-quality*
>
> *ML over 204 Indian stations -- with honest baselines others must beat, one stable*
>
> *NO-O3 hypothesis, and a clear account of what these data can and cannot support.*

Checklist to submission

A. Data & correctness

- [x] Pipeline runs end-to-end on real data (execution evidence)
- [x] Leakage-safe evaluation protocol (blocked-temporal, leave-stations-out, strict)
- [x] Honest baselines (mean, climatology, persistence) + bootstrap CIs
- [x] Code correctness guards (pytest), fixed seeds, reproducibility guide
- [x] **Verify measurement units -- DONE.** Authoritative OpenAQ metadata fetched (`fetch_units.py`) and reconciled (`units_verification.md`): NO, ozone, PM2.5, PM10 unit-consistent; CO, NO2, SO2 have duplicate ppb + µg/m3 sensors with no per-row provenance -> indeterminate. Headline results rest only on unit-consistent variables. Remaining: restrict/caveat CO, NO2, SO2 in the text.

B. Analysis strengthening

- [x] Meteorology-normalized / incremental-value analysis (Phase C2)
- [x] Forecasting vs persistence, reported honestly (Phase C2)
- [x] Bootstrap edge stability + meteorology-confounding test (Phase D)
- [x] Exact-vs-metaheuristic Pareto dominance check (Phase B/F)
- [ ] (Optional) multi-step forecasting if forecasting becomes the headline

C. Figures -- [x] drafted (`audit/reports/figures/`, via `audit/make_figures.py`)

- [x] Fig 1 -- data coverage (per-station + per-pollutant)

- [x] Fig 2 -- leakage: R2 by evaluation protocol
- [x] Fig 3 -- incremental value (weather vs pollutants vs time vs station-mean)
- [x] Fig 4 -- bootstrap edge stability (NO->ozone highlighted)
- [x] Fig 5 -- Pareto dominance (19 policies vs exact front)
- [ ] Polish to the chosen venue's figure style / sizing on submission

D. Writing

- [x] Rough draft (`audit/review2/ROUGH_DRAFT.md`) -- honest, narrowed
- [x] Related Work positioning the paper in the field
- [x] Significance / usefulness-to-the-field section
- [ ] Reframe title/abstract around the **benchmark** contribution (small edit)
- [ ] Tighten Limitations after units are verified
- [ ] **[YOU + me] Verify every citation's DOI/pages** (currently flagged "confirm before submission" in `BIBLIOGRAPHY.md`)

E. Venue & formatting

- [ ] **[YOU + mam] Pick a venue.** Realistic options for a first paper:
- Climate Change AI workshop (NeurIPS / ICLR) -- fits honest applied work
- A regional IEEE conference (indexed proceedings) -- common for college projects
- A datasets/benchmarks track -- fits the resource framing
- [ ] Reformat the draft into that venue's LaTeX template (IEEE / Springer / etc.)
- [ ] Internal critical review pass (overclaim hunt, limitations, reproducibility)

F. Submission

- [ ] **[YOU] Author list, affiliations, acknowledgements** (mam's call)
- [ ] Final proofread -> submit -> (then months of peer review, out of our hands)

What's blocking vs. what's done

- **Done:** execution, evaluation, analysis, figures, full rough draft, pseudocode, tool justification, reproducibility. The *scientific* work is largely complete.
- **Units blocker -- now CLEARED:** verified against authoritative OpenAQ metadata (`units_verification.md`); headline results rest only on unit-consistent variables.
- **Team decisions:** which paper (Path A), which venue, author list.

Realistic timeline

- Units + reframe + citations + venue formatting: **~1-2 focused weeks.**
- Peer review + revisions after that: **months**, and outside our control.

For mam's next review, you can show Execution ? - rough draft ? - pseudocode ? - tool justification ? - figures ? - significance ? - and this concrete publication plan with a single, clearly-named blocking item (units) -- which demonstrates you know exactly what "proper publication" requires and how close you are.

Appendix — Figures




Fig. 1 — Data coverage: 204 stations, ~66% of the full grid; pollutant availability.




Fig. 2 — Leakage inflates skill: honest held-out R2 is 0.22-0.30, not 0.41; persistence is a hard baseline.




Fig. 3 — Pollutants add only +0.12 R2 beyond weather; the station-mean gain is memorisation.




Fig. 4 — Bootstrap edge stability: only NO -> ozone (negative) is stable AND survives meteorology.




Fig. 5 — All 19 metaheuristic 'optimal' policies are dominated by the exact closed-form front.




Fig. 6 — Authoritative units: NO (ppb), ozone/PM2.5/PM10 (µg/m³) are unit-consistent; CO, NO2, SO2 carry duplicate ppb + µg/m³ sensors (indeterminate).