# Observational Response Surfaces versus Interventional Policy: A Critical Audit of Causal Discovery and Multi-Objective Optimization Across Indian Urban Air Quality Networks

Keishav Pratyush Shreerag Jayshil Ryan

October 27, 2026

**Abstract**

The uncritical application of machine learning and multi-objective optimization to ambient air-quality telemetry can produce spurious, policy-shaped conclusions, because non-linear atmospheric chemistry and strong spatio-temporal autocorrelation violate the assumptions on which naive pipelines rest. We audit a complete prediction-to-policy pipeline built on 204 continuous ambient monitoring stations across urban India—1,180,551 station-hours of criteria pollutants ($PM_{2.5}$, $PM_{10}$, NO, $NO_2$, $SO_2$, CO, $O_3$) over calendar year 2025, merged with collocated Open-Meteo reanalysis meteorology—and report three findings. First, spatio-temporal autocorrelation inflates apparent skill under random row splitting: contemporaneous ozone estimation attains $R^2 \approx 0.41$ under an i.i.d. random split but only $R^2 \approx 0.25$ under a strict blocked temporal (Q4) holdout and $R^2 \approx 0.30$ under spatial leave-stations-out validation, while a previous-hour persistence baseline reaches $R^2 \approx 0.79$; pollutants contribute only $\Delta R^2 \approx +0.12$ beyond meteorology. Second, linear structural discovery (DirectLiNGAM) recovers a single bootstrap-stable, meteorology-robust edge consistent with *localized* nitric-oxide titration ($NO + O_3 \rightarrow NO_2 + O_2$) rather than any regional photochemical-smog mechanism, a limitation sharpened by the structural absence of volatile organic compounds (VOCs) and by an authoritative unit audit showing CO, $NO_2$ and $SO_2$ are reported in mixed ppb/$\mu g\,m^{-3}$ units with no per-record provenance. Third, we prove analytically that over the linear response surface the true Pareto frontier collapses to a one-dimensional segment, against which 100% of the heuristic NSGA-II candidate policies are dominated (generational distance = 0.084; hypervolume error = 6.6%). We reframe the artifact as an *open, leakage-safe benchmark* and propose a four-part scientific identification gate for observational environmental AI.

**Keywords:** air quality; data leakage; causal discovery; DirectLiNGAM; multi-objective optimization; Pareto dominance; source apportionment; reproducible benchmark; India.

## 1 Introduction

Urban air pollution is among the most severe environmental health burdens in South Asia. Rapid urbanization and motorization across the Indo-Gangetic Plain and the peninsular metropolitan corridors have driven chronic and widespread exceedances of the National Ambient Air Quality Standards (NAAQS), with annual-mean fine particulate matter ($PM_{2.5}$) across many Indian cities exceeding national limits by large multiples and the World Health Organization guideline by an order of magnitude [8, 17]. The expansion of the Continuous Ambient Air Quality Monitoring Station (CAAQMS) network operated under the Central Pollution Control Board, together with open aggregation platforms such as OpenAQ [15], has made hourly, multi-pollutant telemetry

available at a national scale. This abundance of observational data has, in turn, invited an increasingly ambitious class of data-driven analyses that chain together three steps: *prediction* of ambient concentrations, *discovery* of a causal structure among pollutants, and *optimization* of "policies" framed as pollutant reductions.

### 1.1 Mechanistic models versus observational learning

Two methodological traditions sit in tension. Chemical transport models (CTMs) encode the governing physics and chemistry of the atmosphere and can, in principle, answer genuinely interventional questions—what happens to concentrations if a given emission sector is curtailed. In practice their skill is bounded by bottom-up emission inventories that are frequently incomplete, out of date, or spatially mis-resolved, particularly over rapidly changing Indian source regions [8]. Observational machine learning inverts this trade-off: methods such as gradient-boosted trees [3] and random forests [2] excel at fitting the empirical covariation of pollutants and meteorology, but they estimate *associational response surfaces*, not mechanisms. The central methodological risk is that the fluency of the fit is mistaken for mechanistic or interventional content.

### 1.2 The ozone–$NO_x$ paradox

Nowhere is this risk sharper than for ozone. Tropospheric ozone is a secondary pollutant produced by the photochemical oxidation of volatile organic compounds (VOCs) and carbon monoxide in the presence of nitrogen oxides ($NO_x = NO + NO_2$) and sunlight [14, 21]. Its production rate is a famously non-linear, non-monotone function of precursor concentrations. In the *$NO_x$-limited* regime, typical of rural and background environments, ozone increases with $NO_x$; in the *VOC-limited* (or $NO_x$-saturated) regime, typical of fresh urban emissions, additional $NO_x$ *suppresses* ozone through hydroxyl-radical (OH) scavenging and direct titration, $NO + O_3 \rightarrow NO_2 + O_2$ [24]. Which regime holds at a given site and hour depends on the local VOC-to-$NO_x$ ratio. A single pooled, linear, contemporaneous model fitted across hundreds of heterogeneous urban stations cannot represent this regime dependence; at best it recovers a spatial and temporal average dominated by the urban titration signature. Any apparent "mechanism" it reports is therefore constrained—and potentially confounded—before a single parameter is estimated, especially when VOCs are not measured at all.

### 1.3 The concentration–emission conflation

A further, and more consequential, conflation underlies the "optimization" step. A response surface relates *ambient concentrations* to one another. An actual air quality policy acts on *sectoral emissions*, which map to concentrations only through transport, chemistry and the diurnally varying planetary boundary layer. An optimizer that searches over standardized ambient concentrations and returns a vector such as "set $NO_2$ to $-2\sigma$" is not recommending an emission intervention: it is selecting a point on a correlational surface that may be physically unreachable, and whose objective values carry no policy-benefit interpretation. Treating such a point as a control recommendation is a category error. The distinction between predicting concentrations and simulating interventions is the modern, data-driven instance of the long-standing gap between observation and intervention formalized in the causal-inference literature [18, 19].

### 1.4 Contributions

Rather than add another over-claimed prediction-to-policy pipeline, we audit one. We built the complete three-stage system—clustering, prediction, causal discovery, and multi-objective optimization with post-hoc explanation—ran it on real national-scale telemetry, and then subjected every claim to adversarial, reproducible scrutiny. Our contributions are:

1. **An empirical data and quality-control audit** of 204 CAAQMS stations (1,180,551 station-hours, 2025), including an authoritative per-sensor unit verification that reveals a genuine ppb/$\mu$g m$^{-3}$ unit mix for CO, $NO_2$ and $SO_2$ with no per-record provenance (Section 2).

2. **A leakage-safe predictive evaluation** that contrasts i.i.d. random splitting with blocked temporal and spatial leave-stations-out protocols and a full baseline ladder, quantifying how autocorrelation inflates apparent skill (Sections 3 and 4).

3. **A structural-equation analysis** via DirectLiNGAM with bootstrap edge stability and explicit meteorology-confounding tests, isolating the single robust, chemically interpretable association (Section 4).

4. **An analytical proof of the global Pareto frontier** for the fitted linear surrogate (Lemma 1), against which we benchmark the heuristic NSGA-II solutions and quantify their complete dominance.

5. **A scientific identification gate and an open, reproducible benchmark** for observational environmental AI (Sections 5 and 6).

# 2 Data and Quality Control

## 2.1 Network and acquisition

We assembled hourly observations for the seven criteria pollutants—$PM_{2.5}$, $PM_{10}$, NO, $NO_2$, $SO_2$, CO and $O_3$—from 204 continuous ambient monitoring stations across urban India for calendar year 2025, retrieved through the OpenAQ v3 API [15], which aggregates and standardizes the regulatory CAAQMS/CPCB feeds. Each station was merged, at hourly resolution, with collocated Open-Meteo reanalysis meteorology [28]: 2 m air temperature, relative humidity, 10 m wind speed, and surface pressure, sampled at the nearest $\sim$0.1° reanalysis grid cell and cached per coordinate to avoid redundant retrieval. All timestamps are stored in UTC; because the network spans a single nominal time zone (Indian Standard Time, UTC+5:30), diurnal features are computed after conversion to local solar-aligned hours. The assembled corpus comprises 1,180,551 station-hours.

## 2.2 Coverage and completeness

The realized corpus covers approximately 66% of a complete 204 $\times$ 8,760 station-hour grid (Figure 1a). Completeness is markedly heterogeneous across sites: the median within-year coverage is 0.77, but a left tail of 37 stations falls below 50%. Missingness is a mixture of structural (no sensor deployed for a given species), intermittent (sensor dropout), and below-detection values that are not separable in the absence of detection-limit metadata. Per-species availability within present rows (Figure 1b) is high for $PM_{2.5}$ (0.98) and the gases (0.91–0.96). Because autocorrelated gaps interact dangerously with random splitting (Section 3), all predictive evaluation uses blocked or grouped protocols rather than imputing across gaps.

## 2.3 Sanitization

Three classes of non-physical artifact were identified and flagged. (i) *Instrument saturation / round caps*: distinct spikes of mass at round ceilings (NO, $NO_2$, $O_3$ near 500; $PM_{2.5}$ near 1000). (ii) *Zero-padding and dropout*: runs of exact zeros and abrupt gaps inconsistent with ambient variability. (iii) *Non-physical outliers*: 35 values exceeding physically plausible bounds were flagged and excluded from distributional summaries. Screening these artifacts materially





Figure 1: Network coverage. (a) Distribution of within-year temporal completeness across the 204 stations (mean 66%; 37 stations below 50%, shaded). (b) Per-pollutant non-null fraction within present station-hours. Geolocation of individual monitors is not retained in the public telemetry export and is therefore not mapped; the network is urban and spatially concentrated in the National Capital Region and major metropolitan centres (a representativeness limitation, Section 5).

changes downstream covariance estimates—for example, the CO–NO correlation rises from 0.14 to 0.42—which is itself evidence that naive clustering is sensitive to data hygiene (Section 4).

### 2.4 The unit-provenance discovery

A pre-registration-style verification of measurement units against the *authoritative* OpenAQ per-sensor metadata revealed a material provenance defect (Figure 2). NO is reported in parts-per-billion (ppb) at every station; $O_3$, $PM_{2.5}$ and $PM_{10}$ are reported in mass concentration ($\mu g\,m^{-3}$) and are unit-consistent. However, CO, $NO_2$ and $SO_2$ carry *duplicate sensors reporting in both ppb and* $\mu g\,m^{-3}$ at almost every station (CO at 183/198 stations, $NO_2$ at 187/199, $SO_2$ at 178/191), while the exported telemetry retains a single value per pollutant per station-hour with *no record of which sensor produced it.* A mixing ratio (ppb) and a mass concentration ($\mu g\,m^{-3}$) are related only through molecular weight and ambient temperature and pressure; they are not comparable, and the mix cannot be reconciled post hoc from the stored data. This supersedes an earlier, value-based inference (which had suggested a uniform CO regime): the authoritative sensor metadata governs. Consequently, we restrict all quantitative cross-station inference to the unit-consistent species (NO, $O_3$, $PM_{2.5}$, $PM_{10}$); the two headline results below—ozone estimation and the NO→$O_3$ association—depend only on these.

### 2.5 Summary statistics

Table 1 reports distributional summaries of the sanitized telemetry. Values are in the stored measurement scale; for the unit-indeterminate species (CO, $NO_2$, $SO_2$) the magnitudes are not cross-station comparable and are shown for completeness only. The heavy right skew of $PM_{2.5}$ and $PM_{10}$ (large standard deviations, high 95th percentiles) is characteristic of the Indian urban winter particulate regime.





Figure 2: Authoritative unit verification per species. $PM_{2.5}$, $PM_{10}$ and $O_3$ are uniform $\mu g\,m^{-3}$; NO is uniform ppb; CO, $NO_2$ and $SO_2$ carry duplicate ppb and $\mu g\,m^{-3}$ sensors at the same station (orange), rendering their stored unit indeterminate.

Table 1: Summary statistics of the sanitized telemetry (calendar year 2025, 204 stations). Missing fraction is relative to present station-hours. CO, $NO_2$, $SO_2$ are unit-indeterminate (Section 2.4); PM in $\mu g\,m^{-3}$, $O_3$ in $\mu g\,m^{-3}$, NO in ppb.

| Pollutant | Mean | Std. dev. | 5th pct. | 95th pct. | Missing frac. |
|---|---|---|---|---|---|
| $O_3$ | 30.52 | 30.89 | 2.57 | 89.80 | 0.084 |
| NO | 14.81 | 29.13 | 0.88 | 49.20 | 0.047 |
| $NO_2$ | 27.33 | 27.37 | 2.60 | 80.10 | 0.040 |
| CO | 0.87 | 2.40 | 0.10 | 2.18 | 0.051 |
| $SO_2$ | 13.47 | 25.67 | 1.75 | 35.20 | 0.085 |
| $PM_{10}$ | 121.50 | 116.48 | 18.20 | 349.20 | 0.055 |
| $PM_{2.5}$ | 58.28 | 73.79 | 6.50 | 185.00 | 0.022 |

# 3 Methodology

## 3.1 Governing physical dynamics

The ambient concentration field $C_i(\mathbf{x}, t)$ of chemical species $i$ evolves according to the atmospheric advection–diffusion–reaction equation,

$$\frac{\partial C_i}{\partial t} = \underbrace{-\nabla\cdot(\mathbf{u}\,C_i)}_{\text{advection}} + \underbrace{\nabla\cdot(\mathbf{K}\,\nabla C_i)}_{\text{turbulent diffusion}} + \underbrace{R_i(C_1,\ldots,C_m,\,T,\,h\nu)}_{\text{chemistry}} + \underbrace{E_i(\mathbf{x},t)}_{\text{emissions}}, \tag{1}$$

where $\mathbf{u}(\mathbf{x}, t)$ is the wind velocity field, $\mathbf{K}$ the (second-order) eddy-diffusivity tensor, $R_i$ the net chemical production/loss rate for species $i$ as a function of all $m$ reacting species, temperature $T$ and the actinic photon flux $h\nu$, and $E_i$ the surface emission rate. Equation (1) makes the epistemic gap explicit: a single-site, contemporaneous regression of $C_i$ on the other $C_j$ and a few meteorological scalars discards the spatial transport operators entirely, linearizes the strongly non-linear reaction term $R_i$, and—critically— never observes the emission field $E_i$ on which policy acts. It can estimate a local, time-averaged *response surface*, not the operator in (1).

### 3.2 Predictive formulation and leakage controls

Let $y_{s,t}$ denote the standardized ozone concentration at station $s$ and hour $t$, and $\mathbf{x}_{s,t} \in \mathbb{R}^d$ the vector of contemporaneous covariates (the other six pollutants and four meteorological variables). We estimate $\hat{y} = f(\mathbf{x})$ with gradient-boosted regression trees [3], chosen for their strong tabular performance under non-linear interactions and implemented within a distributed Apache Spark pipeline [27]; hyper-parameters are fixed a priori and never tuned on any evaluation fold.

Because $y_{s,t}$ is strongly autocorrelated in both time and space, the choice of evaluation protocol is not a detail but the central inferential control [11, 20, 26]. We contrast three protocols:

1. *i.i.d. random split* (70/30 over rows): the standard but *leaky* baseline, in which near-duplicate neighbouring hours of the same station appear in both train and test.
2. *Blocked temporal holdout*: train on quarters Q1–Q3 (months 1–9), evaluate on Q4 (months 10–12). Temporal blocks are never split.
3. *Spatial leave-stations-out*: 5-fold `GroupKFold` with the station identifier as the grouping key, so that no station appears in both train and test; this probes transfer to *unmonitored* locations.

All preprocessing (standardization, imputation) is fit on the training fold only. We report against a ladder of reference baselines: (i) *lag-1 persistence*, $\hat{y}_{s,t} = y_{s,t-1}$; (ii) a *station* $\times$ *hour* $\times$ *month diurnal-seasonal mean*, the strongest purely climatological predictor; and (iii) an $L_2$-regularized *Ridge* regression [9] on the same covariates, as a linear reference for the boosted model. Uncertainty is quantified by a station-block bootstrap.

### 3.3 Covariance clustering versus bilinear receptor modelling

The pipeline's first stage groups species by $k$-means over their inter-species Pearson correlation profiles. It is essential to state precisely what this does *not* do. Quantitative source apportionment is a bilinear receptor model; positive matrix factorization (PMF) [16] factorizes a speciated concentration matrix $X = [x_{ij}]$ (sample $i$, chemical species $j$) as

$$x_{ij} = \sum_{k=1}^{p} g_{ik} f_{kj} + e_{ij}, \qquad g_{ik} \geq 0,\ f_{kj} \geq 0, \tag{2}$$

where $g_{ik}$ is the contribution of source $k$ to sample $i$, $f_{kj}$ the chemical profile of source $k$, and $e_{ij}$ a residual weighted by the measurement uncertainty $\sigma_{ij}$ through the objective $Q = \sum_{ij}(e_{ij}/\sigma_{ij})^2$. PMF requires (a) chemically speciated composition (including organic and elemental carbon and key tracers), (b) per-measurement uncertainties, and (c) enough species to identify $p$ non-negative factors. Our data provide only seven criteria pollutants, no speciation, no VOCs, and no per-measurement uncertainties. $k$-means over criteria-gas correlations is therefore an *exploratory inter-species covariance clustering*, and its clusters are mathematically distinct from the latent source factors $\{g_{ik}, f_{kj}\}$ of (2); they admit no mass-balance or source interpretation.

### 3.4 Causal discovery via DirectLiNGAM

For exploratory structure, we model the standardized pollutant vector $\mathbf{z} \in \mathbb{R}^m$ as a linear, non-Gaussian, acyclic structural equation model,

$$\mathbf{z} = \mathbf{B}\,\mathbf{z} + \mathbf{e}, \tag{3}$$

where $\mathbf{B}$ is a strictly lower-triangular (under some causal ordering) matrix of structural coefficients and $\mathbf{e}$ is a vector of mutually independent, non-Gaussian exogenous disturbances. Unlike constraint-based discovery [25], DirectLiNGAM [22, 23] estimates a causal ordering and the

coefficients $\mathbf{B}$ directly, exploiting non-Gaussianity to orient edges; non-linear generalizations exist via additive-noise models [10] but inherit the same confounding sensitivity. Its validity rests on four assumptions: (A1) *acyclicity* of the generating graph; (A2) *linearity* of the structural equations; (A3) *non-Gaussian* disturbances with non-zero variance; and (A4) *causal sufficiency*—no unobserved common causes of two or more observed variables. We encode domain knowledge as a prior-knowledge constraint matrix $K \in \{0, 1, -1\}^{m \times m}$ that forbids edges from secondary pollutants (e.g. $O_3$) into primary combustion tracers (e.g. NO, CO), consistent with emission chemistry. Edge reliability is assessed by a station-block bootstrap (reporting per-edge selection frequency and coefficient sign), and—decisively—by refitting with the meteorological variables included as exogenous nodes, which tests assumption (A4) against the dominant confounder.

### 3.5 Analytical derivation of the global Pareto frontier

The optimization stage searches, with the NSGA-II evolutionary algorithm [5] as implemented in the Optuna framework [1], for policies minimizing two standardized objectives built from the fitted linear response surface. Empirically the fitted surface is

$$z_{\mathrm{O_3}} = \alpha\, z_{\mathrm{NO}}, \qquad \alpha < 0, \tag{4}$$

$$z_{\mathrm{PM_{2.5}}} = \beta_1 z_{\mathrm{NO}} + \beta_2 z_{\mathrm{NO_2}} + \beta_3 z_{\mathrm{CO}}, \qquad \beta_1, \beta_2, \beta_3 > 0, \tag{5}$$

with fitted values $\alpha = -0.124$, $\beta_1 = 0.092$, $\beta_2 = 0.219$, $\beta_3 = 0.228$; $SO_2$ and VOC enter with zero coefficient (VOC is not a fitted variable), so the effective decision vector is $\mathbf{z} = (z_{\mathrm{NO}}, z_{\mathrm{NO_2}}, z_{\mathrm{CO}})$ over the search hypercube $z_j \in [z_{\min}, z_{\max}] = [-2, 0]$ (reductions only, bounded at $-2\sigma$).

**Lemma 1** (Collapse of the linear Pareto front)**.** *Consider the bi-objective minimization of* $\big(f_1(\mathbf{z}), f_2(\mathbf{z})\big) = \big(z_{\mathrm{O_3}}, z_{\mathrm{PM_{2.5}}}\big)$ *defined by* (4)–(5) *over* $\mathbf{z} \in [z_{\min}, z_{\max}]^3$ *with* $\alpha < 0$ *and* $\beta_1, \beta_2, \beta_3 > 0$*. Then every Pareto-optimal point satisfies* $z^\star_{\mathrm{NO_2}} = z^\star_{\mathrm{CO}} = z_{\min}$*, and the Pareto set is the one-dimensional segment*

$$\mathcal{P} = \Big\{ \big(z_{\mathrm{NO}}, z_{\min}, z_{\min}\big) \;:\; z_{\mathrm{NO}} \in [z_{\min}, z_{\max}] \Big\},$$

*whose image is the line segment* $z_{\mathrm{PM_{2.5}}} = \frac{\beta_1}{\alpha}\, z_{\mathrm{O_3}} + (\beta_2 + \beta_3)\, z_{\min}$.

*Proof.* The objective $f_1$ depends only on $z_{\mathrm{NO}}$, while $f_2$ is strictly increasing in each of $z_{\mathrm{NO_2}}$ and $z_{\mathrm{CO}}$ (as $\beta_2, \beta_3 > 0$) and does not appear in $f_1$. Fix any feasible point $\mathbf{z}$ with $z_{\mathrm{NO_2}} > z_{\min}$. The point $\mathbf{z}'$ obtained by lowering $z_{\mathrm{NO_2}}$ to $z_{\min}$ (all else equal) is feasible, leaves $f_1$ unchanged, and strictly decreases $f_2$; hence $\mathbf{z}'$ dominates $\mathbf{z}$, so $\mathbf{z}$ is not Pareto-optimal. The identical argument applies to $z_{\mathrm{CO}}$. Therefore every Pareto-optimal point has $z^\star_{\mathrm{NO_2}} = z^\star_{\mathrm{CO}} = z_{\min}$.

Restricted to this face the objectives reduce to $f_1 = \alpha z_{\mathrm{NO}}$ and $f_2 = \beta_1 z_{\mathrm{NO}} + (\beta_2 + \beta_3) z_{\min}$. Since $\alpha < 0$, $f_1$ is strictly decreasing in $z_{\mathrm{NO}}$, whereas $f_2$ is strictly increasing in $z_{\mathrm{NO}}$ (as $\beta_1 > 0$); the two objectives are therefore in strict conflict along $z_{\mathrm{NO}}$, so no interior point of the segment dominates another and the entire segment is Pareto-optimal. Eliminating $z_{\mathrm{NO}} = f_1/\alpha$ gives $f_2 = \frac{\beta_1}{\alpha} f_1 + (\beta_2 + \beta_3) z_{\min}$, a straight line. □

**Corollary 1.** *The true Pareto frontier of the fitted linear surrogate is a closed-form line segment computable without any metaheuristic search. Any evolutionary approximation can at best match it and, lacking the boundary structure of Lemma 1, will generically return interior, dominated points.*

**Benchmark metrics.** Let $\mathcal{F}$ be a dense discretization of the exact frontier $\mathcal{P}$ and $A = \{\mathbf{a}_k\}_{k=1}^n$ the archive of heuristic solutions in objective space. The *generational distance* is

$$\mathrm{GD}(A) = \Big(\frac{1}{n} \sum_{k=1}^{n} d_k^2\Big)^{1/2}, \qquad d_k = \min_{\mathbf{p} \in \mathcal{F}} \|\mathbf{a}_k - \mathbf{p}\|_2, \tag{6}$$

Table 2: Leakage-safe predictive benchmarking of contemporaneous ozone estimation. All preprocessing fit on the training fold only; hyper-parameters fixed a priori. Persistence and the diurnal-seasonal mean are defined only for the temporal holdout. Lower RMSE/MAE and higher $R^2$ are better.

| Protocol | Model | $R^2$ | RMSE | MAE |
|---|---|---|---|---|
| Random split (leaky) | XGBoost | 0.412 | 23.72 | 14.90 |
| | Ridge ($L_2$) | 0.167 | 28.21 | 18.63 |
| Blocked temporal (Q1–Q3 $\rightarrow$ Q4) | XGBoost | 0.251 | 26.49 | 16.19 |
| | Ridge ($L_2$) | 0.082 | 29.32 | 18.78 |
| | Diurnal-seasonal mean | 0.041 | 29.98 | 20.04 |
| | Persistence (lag-1) | **0.794** | **13.88** | **5.57** |
| Spatial leave-stations-out (5-fold) | XGBoost | 0.304 | 27.94 | 18.45 |
| | Ridge ($L_2$) | 0.159 | 28.33 | 18.73 |
| Temporal $\times$ spatial (strict) | XGBoost | 0.219 | 30.00 | 18.41 |

and, for a shared reference point $\mathbf{r}$ dominated by all points, the *hypervolume error* is

$$\mathrm{HVE} = \frac{\mathrm{HV}(\mathcal{F};\mathbf{r}) - \mathrm{HV}(A;\mathbf{r})}{\mathrm{HV}(\mathcal{F};\mathbf{r})}, \tag{7}$$

where $\mathrm{HV}(\cdot\,;\mathbf{r})$ is the dominated hypervolume [29]. GD $= 0$ and HVE $= 0$ iff the archive lies on the true frontier.

# 4 Results

## 4.1 Predictive benchmarking and the cost of leakage

Table 2 reports contemporaneous ozone estimation across the three evaluation protocols and the baseline ladder. The result is unambiguous: apparent skill is an artifact of the protocol. The gradient-boosted model attains $R^2 = 0.412$ under the i.i.d. random split, but this collapses to $R^2 = 0.251$ under the blocked temporal Q4 holdout and $R^2 = 0.304$ under spatial leave-stations-out, falling further to $R^2 = 0.219$ under the strict temporal$\times$spatial intersection (Figure 3). The $\sim$0.16–0.19 absolute $R^2$ gap between the random split and the blocked/grouped protocols is pure spatio-temporal leakage: neighbouring hours of the same station, near-duplicates of one another, leak across the random train/test boundary [11, 20].

Two baselines frame the honest skill. The linear Ridge reference is weak everywhere ($R^2 \leq 0.17$), confirming that the signal the boosted model captures is genuinely non-linear; but a trivial *lag-1 persistence* predictor attains $R^2 = 0.794$ on the same temporal holdout. The learned model is thus valuable only where persistence is unavailable—namely spatial interpolation to unmonitored sites—and is *beaten by doing nothing* for temporal gap-filling. The diurnal-seasonal climatology ($R^2 = 0.041$) shows that static station behaviour alone explains almost none of the hour-to-hour variance.

## 4.2 Meteorology ablation

Stripping covariate groups under the blocked temporal protocol isolates their contributions. A weather-only model attains $R^2 = 0.140$ and a pollutant-only model $R^2 = 0.132$, while the full model reaches $R^2 = 0.251$. The incremental value of the six pollutants *beyond* meteorology is therefore only $\Delta R^2 \approx +0.117$: most of the explainable ozone variance is meteorological, and the





Figure 3: Held-out $R^2$ for contemporaneous ozone by evaluation protocol. Random splitting of autocorrelated telemetry inflates skill to 0.41; honest blocked and grouped protocols give 0.22–0.30. A previous-hour persistence baseline (green) dominates for temporal gap-filling.

chemical covariates add a small, though non-trivial, increment. Any downstream "mechanistic" reading must account for the fact that meteorology, not chemistry, carries the bulk of the signal.

### 4.3 Recovered structure and its atmospheric interpretation

The bootstrap-stabilized DirectLiNGAM structure is shown in Figure 4. Of all candidate edges, exactly one is both bootstrap-stable (selection frequency 1.0) and survives the inclusion of meteorology as exogenous nodes: the negative edge NO $\rightarrow$ $O_3$, with standardized coefficient $-0.138 \pm 0.007$ in the pollutant-only model, attenuating to $-0.108$ once temperature, humidity, wind and pressure are included. This is the chemical signature of *localized* titration, $NO + O_3 \rightarrow NO_2 + O_2$, expected in fresh, NO-rich urban (traffic-dominated) air [14, 24]: ambient NO destroys ozone locally. Crucially, this is *not* regional photochemical production, which is governed by VOC oxidation that these data cannot observe. The remaining high-frequency edges (NO $\rightarrow$ $NO_2$, +0.40; $NO_2 \rightarrow PM_{2.5}$, +0.23; and the CO combustion edges, frequency 0.91) are consistent with shared combustion co-emission but involve unit-indeterminate species (Section 2.4) and, for the particulate edges, collapse once meteorology is included—evidence that the pollutant-only graph is confounded by the boundary layer (Section 5). $SO_2$ edges are unstable (frequency $\leq 0.5$) and are not retained.

### 4.4 Empirical Pareto frontier versus the NSGA-II archive

Lemma 1 predicts that the true frontier is a one-dimensional segment with $z_{NO_2} = z_{CO} = z_{min}$. We computed this exact segment in closed form and audited the 19 saved NSGA-II candidate policies against it (Table 3, Figure 5). Every one of the 19 heuristic policies is dominated by the exact frontier—a 100% dominance rate—with a mean objective-space gap of 0.046 (maximum 0.298) in standardized $PM_{2.5}$, a generational distance of GD = 0.084 (6), and a hypervolume error of HVE = 6.6% (7). The heuristic fails precisely as Corollary 1 anticipates: unaware of the boundary structure, NSGA-II scatters interior points rather than driving the reducible coordinates to their bound. For a linear surrogate, the metaheuristic is not merely unnecessary—it is strictly inferior to a closed-form solution that any analyst can write down.





Figure 4: Recovered DirectLiNGAM structure (bootstrap frequency $\geq 0.9$). Only NO $\rightarrow$ $O_3$ (red, negative) is both stable and robust to meteorological confounding. Edge width scales with |coefficient|; $SO_2$ carries no stable edge.

### 4.5 Model explainability

Figure 6 shows the global SHAP importance ranking [12, 13] for the ozone model. Relative humidity and temperature dominate, followed by NO as the leading chemical covariate—mirroring both the meteorology ablation and the single stable structural edge. We stress the epistemic status of this agreement: SHAP and DirectLiNGAM are two summaries of the *same* fitted model and data. Their concordance on the sign and prominence of NO is *internal consistency*, not independent validation; it cannot license a causal claim that the leakage-safe and confounding analyses do not already support.

## 5 Discussion: A Scientific Identification Gate

The audit in Section 4 is not a catalogue of incidental shortcomings; the three findings share a common root. A pooled, linear, observational model of ambient telemetry cannot, in principle, answer the interventional and apportionment questions the original pipeline posed. We formalize this as a four-part *identification gate*: criteria that an observational air-quality analysis must pass before causal, apportionment or policy language is warranted.

### 5.1 Gate 1: structural absence of VOCs

Ozone sensitivity is governed by the VOC-to-$NO_x$ ratio [14, 21, 24]. Without continuous hydrocarbon speciation—or an observational proxy such as the formaldehyde-to-$NO_2$ column ratio ($HCHO/NO_2$) from satellite retrievals—the VOC-limited and $NO_x$-limited regimes are empirically indistinguishable in these data. A model that never observes VOCs cannot resolve

Table 3: Audit of the heuristic NSGA-II archive against the exact analytical Pareto frontier of the fitted linear surrogate (Lemma 1). Objectives are standardized; gaps are Euclidean in objective space.

| Quantity | Value |
|---|---|
| Heuristic candidate policies audited | 19 |
| Policies dominated by the exact frontier | 19 (100%) |
| Mean gap to frontier (std. $PM_{2.5}$) | 0.046 |
| Maximum gap to frontier | 0.298 |
| Generational distance, GD | 0.084 |
| Hypervolume error, HVE | 6.6% |
| Exact frontier: $z^\star_{NO_2} = z^\star_{CO}$ | $z_{\min} = -2$ |
| Exact frontier endpoints $(z_{O_3}, z_{PM_{2.5}})$ | $(0, -0.89) \to (0.25, -1.08)$ |





Figure 5: The 19 NSGA-II "optimal" policies (orange) against the exact closed-form Pareto frontier (green) of the linear surrogate. All heuristic points are dominated.

which regime produced a given ozone value, and so cannot attribute an ozone change to a precursor intervention. Our single recovered edge, NO $\to$ $O_3$ $< 0$, is consistent with this limitation: it is the *titration* term, the one piece of ozone chemistry that does not require observing VOCs, and it must not be generalized to regional photochemical production.

### 5.2 Gate 2: unobserved planetary-boundary-layer confounding

The diurnal cycle of the planetary boundary layer (PBL) is a powerful *unobserved common cause*. At night and in winter the PBL collapses, trapping fresh emissions in a shallow volume and simultaneously raising the measured concentrations of all primary, co-emitted species; by day it deepens and dilutes them together. This single latent variable induces strong positive correlations among co-emitted pollutants that have nothing to do with any direct inter-pollutant causal mechanism, violating the causal sufficiency assumption (A4) of (3). The empirical signature is exactly what we observe: the particulate and combustion edges of Figure 4 are stable in the pollutant-only graph but collapse once meteorological proxies of mixing are included. A spuriously "discovered" CO $\to$ $PM_{2.5}$ edge is better read as two species responding to a common,





Figure 6: Global SHAP importance (mean |SHAP|) for ozone estimation. Meteorology dominates; NO is the leading chemical covariate, consistent with Figure 4. Per-sample attributions were not retained, so a beeswarm is not shown.

unmeasured mixing volume. Mitigation requires either direct PBL-height observations or explicit meteorological deconfounding [4, 6, 7].

### 5.3 Gate 3: the concentration–emission fallacy

This is the decisive gate for policy. The optimizer of Lemma 1 acts on *standardized ambient concentrations*: a recommendation such as "$z_{\mathrm{NO}_2} = z_{\min} = -2\sigma$" is a point on a correlational surface, not an emission control. Two failures follow. First, there is no validated map from this concentration vector back to a feasible set of sectoral emission reductions; the inverse problem is precisely the ill-posed source attribution that (1) would have to solve. Second, the search can select physically impossible states: for a right-skewed, non-negative pollutant whose standardized mean sits near zero, a target of $-2\sigma$ can correspond to a *negative physical mass concentration*. Formally, if $C = \mu + \sigma z$ with $\mu \lesssim 2\sigma$ (as holds for several species in Table 1), then $z = -2$ implies $C < 0$. An "optimal policy" that prescribes negative pollutant mass is a reductio: it exposes that the objective never encoded emissions, feasibility, or benefit. Optimizing a concentration response surface and reporting the result as a control strategy is a category error, independent of which optimizer is used.

### 5.4 Gate 4: leakage-aware evaluation

Finally, no skill claim survives without an autocorrelation-aware protocol. As Table 2 shows, the same model and data yield $R^2$ from 0.41 to 0.22 depending solely on the split. Random $k$-fold over rows of spatio-temporal telemetry is not a valid estimate of out-of-sample skill [11, 20]; blocked-temporal and leave-groups-out protocols, benchmarked against persistence and climatology, are the minimum standard.

### 5.5 Recommendations

For researchers and regulators deploying observational environmental AI we recommend: (i) report blocked and grouped validation with persistence and climatology baselines, never a single random-split number; (ii) treat discovered graphs as hypotheses and test every edge against meteorological confounding and bootstrap stability before any causal language; (iii)

verify measurement units and provenance against authoritative sensor metadata, not value heuristics; (iv) reserve "source apportionment" for receptor models (2) with speciation and uncertainties; and (v) never present an optimizer over ambient concentrations as an emission-control recommendation without a validated, feasibility-constrained emissions-to-concentrations model.

# 6 Conclusion

We audited a complete prediction-to-policy air-quality pipeline on 1,180,551 station-hours from 204 Indian urban monitors and found that its most consequential claims do not survive scrutiny. Empirically, spatio-temporal autocorrelation inflates apparent predictive skill from an honest $R^2 \approx 0.22$–$0.30$ to a leaky 0.41, and the learned model is beaten by trivial persistence for temporal gap-filling; pollutants add only $\Delta R^2 \approx 0.12$ beyond meteorology. Structurally, linear causal discovery recovers a single robust edge—localized NO–$O_3$ titration—rather than any regional photochemical mechanism, a ceiling imposed by the structural absence of VOCs and sharpened by a verified ppb/$\mu$g m$^{-3}$ unit-provenance defect in three species. Algorithmically, we proved (Lemma 1) that the true Pareto frontier of the fitted linear surrogate is a closed-form line segment, against which 100% of the heuristic NSGA-II policies are dominated (GD = 0.084, HVE = 6.6%).

Taken together these results motivate the four-part identification gate of Section 5 and a deliberate reframing. Rather than an over-claimed causal source-apportionment and control-optimization system, the artifact is most honestly and usefully released as an *open, leakage-safe benchmark* for multi-pollutant air-quality machine learning: a real national-scale dataset, a reproducible evaluation protocol, a baseline ladder that future methods must beat, one chemically interpretable hypothesis, and an explicit account of what each stronger claim would require. All code, data, seeds and analysis scripts are released to this end.

**Future work.** The productive path forward couples, rather than replaces, mechanism and data. We see three directions: (i) *hybrid modelling*, in which distributed observational surrogates are coupled to, and constrained by, regional three-dimensional chemical transport models (e.g. WRF-Chem, CMAQ) so that interventions are simulated through (1) rather than fit on a correlational surface; (ii) *speciation*, through low-cost VOC and carbonaceous-aerosol sensor arrays or satellite HCHO/$NO_2$ proxies, to pass Gate 1 and enable genuine ozone-regime and receptor-model analysis; and (iii) *identified causal inference*, using documented interventions or natural experiments with meteorological deconfounding and pre-registered estimands. Until an analysis passes the identification gate, the responsible contribution of observational environmental AI is a well-evaluated predictive baseline and an honest map of its own limits.


# Acknowledgements

We thank the Central Pollution Control Board and the OpenAQ and Open-Meteo projects for maintaining the open data infrastructure that made this audit possible.


# Author contributions

Keishav, Pratyush, Shreerag, Jayshil and Ryan jointly designed the study, implemented the pipeline and audit, and wrote the manuscript.

## Reproducibility statement

All code, the committed telemetry dataset, fixed random seeds, per-phase manifests, and the scripts that regenerate every number, table and figure in this manuscript are available in the project repository on the `review/scientific-audit` branch. Every result is reproducible offline without a cluster; the audit scripts fix `SEED = 42`.

## Data availability

Ambient measurements were obtained from the OpenAQ v3 API [15]; meteorological reanalysis from Open-Meteo [28]. Derived analysis artifacts are committed to the repository.

## References


[1] Takuya Akiba, Shotaro Sano, Toshihiko Yanase, Takeru Ohta, and Masanori Koyama. Optuna: A next-generation hyperparameter optimization framework. In *Proceedings of the 25th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining (KDD '19)*, pages 2623–2631, 2019. doi: 10.1145/3292500.3330701.

[2] Leo Breiman. Random forests. *Machine Learning*, 45(1):5–32, 2001. doi: 10.1023/A:1010933404324.

[3] Tianqi Chen and Carlos Guestrin. XGBoost: A scalable tree boosting system. In *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining (KDD '16)*, pages 785–794, 2016. doi: 10.1145/2939672.2939785.

[4] Victor Chernozhukov, Denis Chetverikov, Mert Demirer, Esther Duflo, Christian Hansen, Whitney Newey, and James Robins. Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal*, 21(1):C1–C68, 2018. doi: 10.1111/ectj.12097.

[5] Kalyanmoy Deb, Amrit Pratap, Sameer Agarwal, and T. Meyarivan. A fast and elitist multiobjective genetic algorithm: NSGA-II. *IEEE Transactions on Evolutionary Computation*, 6(2):182–197, 2002. doi: 10.1109/4235.996017.

[6] Stuart K. Grange and David C. Carslaw. Using meteorological normalisation to detect interventions in air quality time series. *Science of the Total Environment*, 653:578–588, 2019. doi: 10.1016/j.scitotenv.2018.10.344.

[7] Stuart K. Grange, David C. Carslaw, Alastair C. Lewis, Eirini Boleti, and Christoph Hueglin. Random forest meteorological normalisation models for Swiss PM10 trend analysis. *Atmospheric Chemistry and Physics*, 18(9):6223–6239, 2018. doi: 10.5194/acp-18-6223-2018.

[8] Sarath K. Guttikunda and Giuseppe Calori. A GIS based emissions inventory at 1 km $\times$ 1 km spatial resolution for air pollution analysis in Delhi, India. *Atmospheric Environment*, 67:101–111, 2013. doi: 10.1016/j.atmosenv.2012.10.040.

[9] Trevor Hastie, Robert Tibshirani, and Jerome Friedman. *The Elements of Statistical Learning: Data Mining, Inference, and Prediction.* Springer, 2nd edition, 2009. doi: 10.1007/978-0-387-84858-7.

[10] Patrik O. Hoyer, Dominik Janzing, Joris M. Mooij, Jonas Peters, and Bernhard Schölkopf. Nonlinear causal discovery with additive noise models. In *Advances in Neural Information Processing Systems 21 (NeurIPS)*, pages 689–696, 2009.

[11] Shachar Kaufman, Saharon Rosset, Claudia Perlich, and Ori Stitelman. Leakage in data mining: Formulation, detection, and avoidance. *ACM Transactions on Knowledge Discovery from Data*, 6(4):15:1–15:21, 2012. doi: 10.1145/2382577.2382579.

[12] Scott M. Lundberg and Su-In Lee. A unified approach to interpreting model predictions. In *Advances in Neural Information Processing Systems 30 (NeurIPS)*, pages 4765–4774, 2017. arXiv:1705.07874.

[13] Scott M. Lundberg, Gabriel Erion, Hugh Chen, Alex DeGrave, Jordan M. Prutkin, Bala Nair, Ronit Katz, Jonathan Himmelfarb, Nisha Bansal, and Su-In Lee. From local explanations to global understanding with explainable AI for trees. *Nature Machine Intelligence*, 2(1): 56–67, 2020. doi: 10.1038/s42256-019-0138-9.

[14] P. S. Monks, A. T. Archibald, A. Colette, O. Cooper, M. Coyle, R. Derwent, D. Fowler, C. Granier, K. S. Law, G. E. Mills, D. S. Stevenson, et al. Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer. *Atmospheric Chemistry and Physics*, 15(15):8889–8973, 2015. doi: 10.5194/acp-15-8889-2015.

[15] OpenAQ. OpenAQ: Open air quality data platform (api v3). https://openaq.org, 2025. Accessed 2025.

[16] Pentti Paatero and Unto Tapper. Positive matrix factorization: A non-negative factor model with optimal utilization of error estimates of data values. *Environmetrics*, 5(2):111–126, 1994. doi: 10.1002/env.3170050203.

[17] Pallavi Pant, Anuradha Shukla, Steven D. Kohl, Judith C. Chow, John G. Watson, and Roy M. Harrison. Characterization of ambient PM2.5 at a pollution hotspot in New Delhi, India and inference of sources. *Atmospheric Environment*, 109:178–189, 2015. doi: 10.1016/j.atmosenv.2015.02.074.

[18] Judea Pearl. *Causality: Models, Reasoning, and Inference*. Cambridge University Press, 2nd edition, 2009. doi: 10.1017/CBO9780511803161.

[19] Jonas Peters, Dominik Janzing, and Bernhard Schölkopf. *Elements of Causal Inference: Foundations and Learning Algorithms*. MIT Press, Cambridge, MA, 2017. ISBN 978-0-262-03731-0.

[20] David R. Roberts, Volker Bahn, Simone Ciuti, Mark S. Boyce, Jane Elith, Gurutzeta Guillera-Arroita, Severin Hauenstein, José J. Lahoz-Monfort, Boris Schröder, Wilfried Thuiller, David I. Warton, Brendan A. Wintle, Florian Hartig, and Carsten F. Dormann. Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. *Ecography*, 40(8):913–929, 2017. doi: 10.1111/ecog.02881.

[21] John H. Seinfeld and Spyros N. Pandis. *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*. John Wiley & Sons, Hoboken, NJ, 3rd edition, 2016. ISBN 978-1-118-94740-1.

[22] Shohei Shimizu, Patrik O. Hoyer, Aapo Hyvärinen, and Antti Kerminen. A linear non-Gaussian acyclic model for causal discovery. *Journal of Machine Learning Research*, 7: 2003–2030, 2006. URL https://jmlr.org/papers/v7/shimizu06a.html.

[23] Shohei Shimizu, Takanori Inazumi, Yasuhiro Sogawa, Aapo Hyvärinen, Yoshinobu Kawahara, Takashi Washio, Patrik O. Hoyer, and Kenneth Bollen. DirectLiNGAM: A direct method for learning a linear non-Gaussian structural equation model. *Journal of Machine Learning Research*, 12:1225–1248, 2011. URL https://jmlr.org/papers/v12/shimizu11a.html.

[24] Sanford Sillman. The relation between ozone, NOx and hydrocarbons in urban and polluted rural environments. *Atmospheric Environment*, 33(12):1821–1845, 1999. doi: 10.1016/S1352-2310(98)00345-8.

[25] Peter Spirtes, Clark Glymour, and Richard Scheines. *Causation, Prediction, and Search*. MIT Press, Cambridge, MA, 2nd edition, 2000.

[26] Sudhir Varma and Richard Simon. Bias in error estimation when using cross-validation for model selection. *BMC Bioinformatics*, 7:91, 2006. doi: 10.1186/1471-2105-7-91.

[27] Matei Zaharia, Reynold S. Xin, Patrick Wendell, Tathagata Das, Michael Armbrust, Ankur Dave, Xiangrui Meng, Josh Rosen, Shivaram Venkataraman, Michael J. Franklin, Ali Ghodsi, Joseph Gonzalez, Scott Shenker, and Ion Stoica. Apache Spark: A unified engine for big data processing. *Communications of the ACM*, 59(11):56–65, 2016. doi: 10.1145/2934664.

[28] Patrick Zippenfenig. Open-Meteo: Historical weather reanalysis api. https://open-meteo.com, 2025. Accessed 2025.

[29] Eckart Zitzler and Lothar Thiele. Multiobjective evolutionary algorithms: a comparative case study and the strength Pareto approach. *IEEE Transactions on Evolutionary Computation*, 3(4):257–271, 1999. doi: 10.1109/4235.797969.