## Appendix

## Lemma 4

Broadly, Lemma 4 states that for agents i and k, if k is in  $N_i^c(t)$  and is not a first-degree neighbor of i, then there exists an agent j that is a first-degree neighbor of i that is connected to both i and k. There are three properties that hold for Lemma 4 between the agents i and k. Of the 12 possible pairs for (i, k), we highlight the formulas used for the pair (1, 2) in this section. The three dL formulas representing Lemma 4 is as follows:

PrL4\_12\_1 is proven directly, and is equivalent to Inv6\_12 below. PrL4\_12\_2 is proven using Inv7\_12 and Inv8\_12. Lastly, PrL4\_12\_3 is proven using the formulas for Lemma 3, namely PrL3\_13 and PrL3\_14.

```
Inv6_12 \triangleq A1SN2 = 0 \lor A1SN2 + replan1 \neq 5 \lor (A1SN2 + replan1 = 5)
              \land ((distance(x1,y1,x3,y3) \le R \land distance(x3,y3,x2,y2) \le R \land distance(x1,y1,x4,y4) \le R
              \land distance(x4,y4,x2,y2) \leq R)
              \vee (distance(x1,y1,x3,y3) \leq R \wedge \text{distance}(x3,y3,x2,y2) \leq R) \vee (\text{distance}(x1,y1,x4,y4) \leq R
              \land distance(x4,y4,x2,y2) \leq R)))
Inv7_12 \triangleq A1SN2 = 0 \lor A1SN2 + replan1 \neq 5 \lor (A1SN2 + replan1 = 5)
             \land ((distance(x1,y1,x3,y3) \le R \land distance(x3,y3,x2,y2) \le R \land distance(x1,y1,x4,y4) \le R
                \land distance(x4,y4,x2,y2) \leq R
                \land N1C3 = 1 \land e1N3C2 = 1 \land e1N2C3 = 1 \land N1C4 = 1 \land e1N4C2 = 1 \land e1N2C4 = 1)
              \vee (distance(x1,y1,x3,y3) \leq R \wedge \text{distance}(x3,y3,x2,y2) \leq R
                \land N1C3 = 1 \land e1N3C2 = 1 \land e1N2C3 = 1 \land (N1C4 = 0 \lor e1N4C2 = 0 \lor e1N2C4 = 0))
              \vee (distance(x1,y1,x4,y4) \leq R \wedge \text{distance}(x4,y4,x2,y2) \leq R
                \land (N1C3 = 0 \lor e1N3C2 = 0 \lor e1N2C3 = 0) \land N1C4 = 1 \land e1N4C2 = 1 \land e1N2C4 = 1)))
Inv8_12 \triangleq (distance(x3,y3,x2,y2) \leq R \lor e1N3C2 = 0 \lor e1N2C3 = 0 \lor
               (distance(x3,y3,x2,y2) > R \land e1N3C2 = 1 \land e1N2C3 = 1 \land N3C2 = 1
                \land N2C3 = 1)
              \land (\mathsf{distance}(\mathsf{x4}, \mathsf{y4}, \mathsf{x2}, \mathsf{y2}) \leq R \lor \mathsf{e1N4C2} = 0 \lor \mathsf{e1N2C4} = 0 \lor
               (distance(x4,y4,x2,y2) > R \land e1N4C2 = 1 \land e1N2C4 = 1 \land N4C2 = 1
                \wedge N2C4 = 1)
```

## Lemmas 5-7

The dL formulas representing lemmas 5, 6, and 7 are weaker cases of lemma 4, and thus can be proven using the more general formulas and invariants of lemma 4.

$$\mathsf{PrL5}\_12\_1 \triangleq \mathsf{A1SN2} = 0 \lor \mathsf{A1SN2} + \mathsf{replan1} \neq 5 \lor \mathsf{N1C2} \geq \mathsf{N1C2old} \lor (\mathsf{A1SN3} + \mathsf{replan1} = 5 \lor \mathsf{N1C2old} \lor (\mathsf{A1SN3} + \mathsf{replan1} = 5 \lor \mathsf{N1C2old} \lor \mathsf{N1C2o$$

```
\land N1C2 < N1C2old \land InvDist)
\mathsf{PrL5}\_12\_2 \triangleq \mathsf{A1SN2} = 0 \lor \mathsf{A1SN2} + \mathsf{replan1} \neq 5 \lor \mathsf{N1C2} \ge \mathsf{N1C2old} \lor (\mathsf{A1SN2} + \mathsf{replan1} = 5)
                  \land N1C2 < N1C2old \land InvNiC),
PrL5_12_3 \triangleq A1SN2 = 0 \lor A1SN2 + replan1 \neq 5 \lor N1C2 \ge N1C2old
                  \lor \mathsf{slot} 13 = 0 \lor \mathsf{slot} 14 = 0 \lor (\mathsf{A1SN2} + \mathsf{replan1} = 5
                  \land N1C2 < N1C2old \land ((PrL3_13 \land PrL3_14) \lor PrL3_13 \lor PrL3_14))
PrL6_12_2 \triangleq A1SN2 = 0 \lor A1SN2 + replan1 \neq 5 \lor N1C2 \ge N1C2old \lor Rhat1old \neq 0
                  \lor (A1SN2 + replan1 = 5 \land N1C2 < N1C2old \land Rhat1old = 0 \land InvNiC)
PrL6_12_3 \triangleq A1SN2 = 0 \lor A1SN2 + replan1 \neq 5 \lor N1C2 \ge N1C2old \lor Rhat1old \neq 0
                  \lor (A1SN2 + replan1 = 5 \land N1C2 < N1C2old \land Rhat1old = 0
                  \land ((PrL3\_13 \land PrL3\_14) \lor PrL3\_13 \lor PrL3\_14))
PrL7_12_1 \triangleq A1SN2 = 0 \lor A1SN2 + replan1 \neq 5 \lor N1C2 > N1C2old \lor Rhat1old \neq 0
                  \lor (A1SN2 + replan1 = 5 \land N1C2 < N1C2old \land Rhat1old = 0 \land InvDist_2)
\mathsf{PrL7}\_\mathsf{12}\_\mathsf{2} \triangleq \mathsf{A1SN2} = 0 \lor \mathsf{A1SN2} + \mathsf{replan1} \neq 5 \lor \mathsf{N1C2} \geq \mathsf{N1C2old} \lor \mathsf{Rhat1old} \neq 0
                  \lor (A1SN2 + replan1 = 5 \land N1C2 < N1C2old \land Rhat1old = <math>0 \land InvNiC_2)
PrL7_12_3 \triangleq A2SN1 = 0 \lor A2SN1 + replan2 \neq 5 \lor N1C2 \ge N1C2old \lor Rhat1old \neq 0
                  \lor \mathsf{Rhat2old} \neq 0 \lor \lor (\mathsf{A1SN2} + \mathsf{replan1} = 5 \land \mathsf{N1C2} < \mathsf{N1C2old} \land \mathsf{Rhat1old} = 0
                  \land \mathsf{Rhat2old} = 0 \land \mathsf{InvDist\_2})
PrL7_12_4 \triangleq A2SN1 = 0 \lor A2SN1 + replan2 \neq 5 \lor N1C2 \ge N1C2old \lor Rhat1old \neq 0
                  \lor Rhat2old \neq 0 \lor \lor (A1SN2 + replan1 = 5 \land N1C2 < N1C2old \land Rhat1old = 0
                  \land \mathsf{Rhat2old} = 0 \land \mathsf{InvNiC}_2), where
   InvDist \triangleq ((distance(x1,y1,x3,y3) \leq R \land distance(x3,y3,x2,y2) \leq R \land distance(x1,y1,x4,y4) \leq R
                  \land distance(x4,y4,x2,y2) \leq R) \lor (distance(x1,y1,x3,y3) \leq R \land distance(x3,y3,x2,y2) \leq R)
                  \vee (distance(x1,y1,x4,y4) \leq R \wedge \text{distance}(x4,y4,x2,y2) \leq R)
    InvNiC \triangleq ((N1C3 = 1 \land N3C2 = 1 \land N2C3 = 1 \land N1C4 = 1 \land N4C2 = 1 \land N2C4 = 1)
                  \lor (N1C3 = 1 \land N3C2 = 1 \land N2C3 = 1 \land (N1C4 = 0 \lor N4C2 = 0 \lor N2C4 = 0))
                  \lor ((N1C3 = 0 \lor N3C2 = 0 \lor N2C3 = 0) \land N1C4 = 1 \land N4C2 = 1 \land N2C4 = 1))
 InvDist.2 \triangleq (distance(x1,y1,x3,y3) \leq R \land distance(x3,y3,x2,y2) \leq R \land distance(x1,y1,x4,y4) \leq R \land
                  \land distance(x4,y4,x2,y2) \le R) \lor (distance(x1,y1,x3,y3) \le R \land distance(x3,y3,x2,y2) \le R)
                  \vee (distance(x1,y1,x4,y4) \leq R \wedge \text{distance}(x4,y4,x2,y2) \leq R) \vee (\text{distance}(x1,y1,x3,y3) \leq R
                  \land distance(x3,y3,x4,y4) \leq R \land distance(x4,y4,x2,y2) \leq R) \lor (distance(x1,y1,x4,y4) \leq R
                  \land distance(x4,y4,x3,y3) \leq R \land distance(x3,y3,x2,y2) \leq R)
 InvNiC_2 \triangleq ((N1C3 = 1 \land N3C2 = 1 \land N2C3 = 1 \land N1C4 = 1 \land N4C2 = 1 \land N2C4 = 1)
                  \vee (N1C3 = 1 \wedge N3C2 = 1 \wedge N2C3 = 1 \wedge (N1C4 = 0 \vee N4C2 = 0 \vee N2C4 = 0))
                  \lor ((N1C3 = 0 \lor N3C2 = 0 \lor N2C3 = 0) \land N1C4 = 1 \land N4C2 = 1 \land N2C4 = 1)
                  \lor (N1C3 = 1 \land N3C4 = 1 \land N4C3 = 1 \land N4C2 = 1 \land N2C4 = 1)
```

 $\vee (N1C4 = 1 \land N4C3 = 1 \land N3C4 = 1 \land N3C2 = 1 \land N2C3 = 1))$