

Class 14 Mini Project: Pathway Analysis from RNA-Seq Results

Harshita Jha (PID: A17350910)

Table of contents

Background	1
Differential Expression Analysis	1
Data Import	1
Volcano Plot	4
Adding gene annotation	6
Pathway Analysis	8
Gene Ontology	22
Reactome Analysis	24

Background

Our data for today comes from HOX gene knock-out study.

The authors report on differential analysis of lung fibroblasts in response to loss of the developmental transcription factor HOXA1

Differential Expression Analysis

Data Import

- First step is to download the count data and associated meta data into our document.

```
library(DESeq2)
```

- Now load our data:

```
metaFile <- "GSE37704_metadata.csv"
countFile <- "GSE37704_featurecounts.csv"
```

```
colData = read.csv(metaFile, row.names=1)
head(colData)
```

```

              condition
SRR493366 control_sirna
SRR493367 control_sirna
SRR493368 control_sirna
SRR493369      hoxa1_kd
SRR493370      hoxa1_kd
SRR493371      hoxa1_kd
```

```
countData = read.csv(countFile, row.names=1)
head(countData)
```

```

              length SRR493366 SRR493367 SRR493368 SRR493369 SRR493370
ENSG00000186092    918         0         0         0         0         0
ENSG00000279928    718         0         0         0         0         0
ENSG00000279457   1982        23        28        29        29        28
ENSG00000278566    939         0         0         0         0         0
ENSG00000273547    939         0         0         0         0         0
ENSG00000187634   3214       124       123       205       207       212
              SRR493371
ENSG00000186092         0
ENSG00000279928         0
ENSG00000279457        46
ENSG00000278566         0
ENSG00000273547         0
ENSG00000187634       258
```

Q. Complete the code below to remove the troublesome first column from count-Data

```
countData <- as.matrix(countData[,-1])
head(countData)
```

```

              SRR493366 SRR493367 SRR493368 SRR493369 SRR493370 SRR493371
ENSG00000186092         0         0         0         0         0         0
```

ENSG00000279928	0	0	0	0	0	0
ENSG00000279457	23	28	29	29	28	46
ENSG00000278566	0	0	0	0	0	0
ENSG00000273547	0	0	0	0	0	0
ENSG00000187634	124	123	205	207	212	258

- Now let's get rid of the zero entries. We should remove these before any further analysis.

Q. Complete the code below to filter countData to exclude genes (i.e. rows) where we have 0 read count across all samples (i.e. columns). Tip: What will rowSums() of countData return and how could you use it in this context?

```
countData = countData[rowSums(countData) > 0, ]
head(countData)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000279457	23	28	29	29	28	46
ENSG00000187634	124	123	205	207	212	258
ENSG00000188976	1637	1831	2383	1226	1326	1504
ENSG00000187961	120	153	180	236	255	357
ENSG00000187583	24	48	65	44	48	64
ENSG00000187642	4	9	16	14	16	16

- Now, run the DESeq2:

```
dds <- DESeqDataSetFromMatrix(countData=countData,
                              colData=colData,
                              design=~condition)
```

Warning in DESeqDataSet(se, design = design, ignoreRank): some variables in design formula are characters, converting to factors

```
dds <- DESeq(dds)
```

estimating size factors

estimating dispersions

gene-wise dispersion estimates

mean-dispersion relationship

final dispersion estimates

fitting model and testing

```
dds
```

```
class: DESeqDataSet
dim: 15975 6
metadata(1): version
assays(4): counts mu H cooks
rownames(15975): ENSG00000279457 ENSG00000187634 ... ENSG00000276345
               ENSG00000271254
rowData names(22): baseMean baseVar ... deviance maxCooks
colnames(6): SRR493366 SRR493367 ... SRR493370 SRR493371
colData names(2): condition sizeFactor
```

```
res <- results(dds)
```

```
summary(res)
```

```
out of 15975 with nonzero total read count
adjusted p-value < 0.1
LFC > 0 (up)      : 4349, 27%
LFC < 0 (down)    : 4396, 28%
outliers [1]      : 0, 0%
low counts [2]    : 1237, 7.7%
(mean count < 0)
[1] see 'cooksCutoff' argument of ?results
[2] see 'independentFiltering' argument of ?results
```


Volcano Plot

Now, we will make a volcano plot of log2 fold change vs -log adjusted p-value.

```
library(ggplot2)
```

```
ggplot(res) +
  aes(log2FoldChange,
      -log(padj)) +
  geom_point()
```

Warning: Removed 1237 rows containing missing values or values outside the scale range (`geom\_point()`).



Q. Improve this plot by completing the below code, which adds color, axis labels and cutoff lines:


```
# Make a color vector for all genes
mycols <- rep("gray", nrow(res) )

# Color blue the genes with fold change above 2
mycols[ abs(res$log2FoldChange) > 2 ] <- "blue"

# Color gray those with adjusted p-value more than 0.01
mycols[ res$padj > 0.01 ] <- "gray"

ggplot(res) +
  aes(log2FoldChange,
      -log(padj)) +
  geom_point(color=mycols) +
  xlab("Log2(FoldChange)") +
  ylab("-Log(P-value)") +
  geom_vline(xintercept = c(-2,2)) +
  geom_hline(yintercept = -log(0.01))
```

Warning: Removed 1237 rows containing missing values or values outside the scale range (``geom_point()``).



Adding gene annotation

Q. Use the `mapIDs()` function multiple times to add SYMBOL, ENTREZID and GENENAME annotation to our results by completing the code below.

```
library("AnnotationDbi")
library("org.Hs.eg.db")
```

```
columns(org.Hs.eg.db)
```

[1]	"ACCNUM"	"ALIAS"	"ENSEMBL"	"ENSEMBLPROT"	"ENSEMBLTRANS"
[6]	"ENTREZID"	"ENZYME"	"EVIDENCE"	"EVIDENCEALL"	"GENENAME"
[11]	"GENETYPE"	"GO"	"GOALL"	"IPI"	"MAP"
[16]	"OMIM"	"ONTOLOGY"	"ONTOLOGYALL"	"PATH"	"PFAM"
[21]	"PMID"	"PROSITE"	"REFSEQ"	"SYMBOL"	"UCSCKG"
[26]	"UNIPROT"				

```
res$symbol = mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="SYMBOL",
                    multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
res$entrez = mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="ENTREZID",
                    multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
res$name = mapIds(org.Hs.eg.db,
                  keys=row.names(res),
                  keytype="ENSEMBL",
                  column="GENENAME",
                  multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
head(res, 10)
```

log2 fold change (MLE): condition hoxa1 kd vs control sirna

Wald test p-value: condition hoxa1 kd vs control sirna

DataFrame with 10 rows and 9 columns

	baseMean	log2FoldChange	lfcSE	stat	pvalue
	<numeric>	<numeric>	<numeric>	<numeric>	<numeric>
ENSG00000279457	29.913579	0.1792571	0.3248216	0.551863	5.81042e-01
ENSG00000187634	183.229650	0.4264571	0.1402658	3.040350	2.36304e-03
ENSG00000188976	1651.188076	-0.6927205	0.0548465	-12.630158	1.43990e-36
ENSG00000187961	209.637938	0.7297556	0.1318599	5.534326	3.12428e-08
ENSG00000187583	47.255123	0.0405765	0.2718928	0.149237	8.81366e-01
ENSG00000187642	11.979750	0.5428105	0.5215598	1.040744	2.97994e-01
ENSG00000188290	108.922128	2.0570638	0.1969053	10.446970	1.51282e-25
ENSG00000187608	350.716868	0.2573837	0.1027266	2.505522	1.22271e-02

ENSG00000188157	9128.439422	0.3899088	0.0467163	8.346304	7.04321e-17
ENSG00000237330	0.158192	0.7859552	4.0804729	0.192614	8.47261e-01
	padj	symbol	entrez		name
	<numeric>	<character>	<character>		<character>
ENSG00000279457	6.86555e-01	NA	NA		NA
ENSG00000187634	5.15718e-03	SAMD11	148398	sterile alpha motif ..	
ENSG00000188976	1.76549e-35	NOC2L	26155	NOC2 like nucleolar ..	
ENSG00000187961	1.13413e-07	KLHL17	339451	kelch like family me..	
ENSG00000187583	9.19031e-01	PLEKHN1	84069	pleckstrin homology ..	
ENSG00000187642	4.03379e-01	PERM1	84808	PPARGC1 and ESRR ind..	
ENSG00000188290	1.30538e-24	HES4	57801	hes family bHLH tran..	
ENSG00000187608	2.37452e-02	ISG15	9636	ISG15 ubiquitin like..	
ENSG00000188157	4.21963e-16	AGRN	375790		agrin
ENSG00000237330	NA	RNF223	401934	ring finger protein ..	

Q.Finally for this section let's reorder these results by adjusted p-value and save them to a CSV file in your current project directory.

```
res <- res[order(res$pvalue),]
write.csv(res, file ="deseq_results.csv")
```

Pathway Analysis

First we load up the packaged required and set up the KEGG data-sets we need:

```
library(pathview)
library(gage)
library(gageData)

data(kegg.sets.hs)
data(sigmet.idx.hs)

# Focus on signaling and metabolic pathways only
kegg.sets.hs = kegg.sets.hs[sigmet.idx.hs]

# Examine the first 3 pathways
head(kegg.sets.hs, 3)
```

```
$`hsa00232 Caffeine metabolism`
[1] "10" "1544" "1548" "1549" "1553" "7498" "9"
```


\$`hsa00983 Drug metabolism - other enzymes`

```
[1] "10"      "1066"    "10720"   "10941"   "151531"  "1548"    "1549"    "1551"
[9] "1553"    "1576"    "1577"    "1806"    "1807"    "1890"    "221223"  "2990"
[17] "3251"    "3614"    "3615"    "3704"    "51733"   "54490"   "54575"   "54576"
[25] "54577"   "54578"   "54579"   "54600"   "54657"   "54658"   "54659"   "54963"
[33] "574537"  "64816"   "7083"    "7084"    "7172"    "7363"    "7364"    "7365"
[41] "7366"    "7367"    "7371"    "7372"    "7378"    "7498"    "79799"   "83549"
[49] "8824"    "8833"    "9"       "978"
```

\$`hsa00230 Purine metabolism`

```
[1] "100"      "10201"   "10606"   "10621"   "10622"   "10623"   "107"     "10714"
[9] "108"      "10846"   "109"     "111"     "11128"   "11164"   "112"     "113"
[17] "114"      "115"     "122481"  "122622"  "124583"  "132"     "158"     "159"
[25] "1633"     "171568"  "1716"    "196883"  "203"     "204"     "205"     "221823"
[33] "2272"     "22978"   "23649"   "246721"  "25885"   "2618"    "26289"   "270"
[41] "271"      "27115"   "272"     "2766"    "2977"    "2982"    "2983"    "2984"
[49] "2986"     "2987"    "29922"   "3000"    "30833"   "30834"   "318"     "3251"
[57] "353"      "3614"    "3615"    "3704"    "377841"  "471"     "4830"    "4831"
[65] "4832"     "4833"    "4860"    "4881"    "4882"    "4907"    "50484"   "50940"
[73] "51082"    "51251"   "51292"   "5136"    "5137"    "5138"    "5139"    "5140"
[81] "5141"     "5142"    "5143"    "5144"    "5145"    "5146"    "5147"    "5148"
[89] "5149"     "5150"    "5151"    "5152"    "5153"    "5158"    "5167"    "5169"
[97] "51728"    "5198"    "5236"    "5313"    "5315"    "53343"   "54107"   "5422"
[105] "5424"     "5425"    "5426"    "5427"    "5430"    "5431"    "5432"    "5433"
[113] "5434"     "5435"    "5436"    "5437"    "5438"    "5439"    "5440"    "5441"
[121] "5471"     "548644"  "55276"   "5557"    "5558"    "55703"   "55811"   "55821"
[129] "5631"     "5634"    "56655"   "56953"   "56985"   "57804"   "58497"   "6240"
[137] "6241"     "64425"   "646625"  "654364"  "661"     "7498"    "8382"    "84172"
[145] "84265"    "84284"   "84618"   "8622"    "8654"    "87178"   "8833"    "9060"
[153] "9061"     "93034"   "953"     "9533"    "954"     "955"     "956"     "957"
[161] "9583"     "9615"
```

```
foldchanges = res$log2FoldChange
names(foldchanges) = res$entrez
head(foldchanges)
```

```
      1266      54855      1465      2034      2150      6659
-2.422719  3.201955 -2.313738 -1.888019  3.344508  2.392288
```

- Now, we can run the **gage** pathway analysis:

```
keggres = gage(foldchanges, gsets=kegg.sets.hs)
attributes(keggres)
```

```
$names
[1] "greater" "less"    "stats"
```

- Now, we can look at the first few down pathways:

```
head(keggres$less)
```

	p.geomean	stat.mean	p.val
hsa04110 Cell cycle	8.995727e-06	-4.378644	8.995727e-06
hsa03030 DNA replication	9.424076e-05	-3.951803	9.424076e-05
hsa03013 RNA transport	1.375901e-03	-3.028500	1.375901e-03
hsa03440 Homologous recombination	3.066756e-03	-2.852899	3.066756e-03
hsa04114 Oocyte meiosis	3.784520e-03	-2.698128	3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis	8.961413e-03	-2.405398	8.961413e-03

	q.val	set.size	exp1
hsa04110 Cell cycle	0.001448312	121	8.995727e-06
hsa03030 DNA replication	0.007586381	36	9.424076e-05
hsa03013 RNA transport	0.073840037	144	1.375901e-03
hsa03440 Homologous recombination	0.121861535	28	3.066756e-03
hsa04114 Oocyte meiosis	0.121861535	102	3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis	0.212222694	53	8.961413e-03

- Pathway for the cell cycle:

```
pathview(gene.data=foldchanges, pathway.id="hsa04110")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/harshitajha/BIMM143/Class14

Info: Writing image file hsa04110.pathview.png


```
pathview(gene.data=foldchanges, pathway.id=keggresids, species="hsa")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/harshitajha/BIMM143/Class14

Info: Writing image file hsa04640.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/harshitajha/BIMM143/Class14

Info: Writing image file hsa04630.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/harshitajha/BIMM143/Class14

Info: Writing image file hsa00140.pathview.png

'select()' returned 1:1 mapping between keys and columns


Info: Working in directory /Users/harshitajha/BIMM143/Class14



Info: Writing image file hsa04142.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/harshitajha/BIMM143/Class14

Info: Writing image file hsa04330.pathview.png





Q. Can you do the same procedure as above to plot the pathview figures for the top 5 down-regulated pathways?


```
## Focus on top 5 down-regulated pathways
keggrespathways_down <- rownames(keggres$less)[1:5]

# Extract the 8 character long IDs part of each string
keggresids_down <- substr(keggrespathways, start=1, stop=8)
keggresids_down
```

```
[1] "hsa04640" "hsa04630" "hsa00140" "hsa04142" "hsa04330"
```

```
pathview(gene.data=foldchanges, pathway.id=keggresids_down, species="hsa")
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/harshitajha/BIMM143/Class14
```

```
Info: Writing image file hsa04640.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/harshitajha/BIMM143/Class14
```

```
Info: Writing image file hsa04630.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/harshitajha/BIMM143/Class14
```

```
Info: Writing image file hsa00140.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```


```
Info: Working in directory /Users/harshitajha/BIMM143/Class14
```


```
Info: Writing image file hsa04142.pathview.png
```


```
'select()' returned 1:1 mapping between keys and columns
```

Info: Writing image file hsa04330.pathview.png









G0:0007156	homophilic cell adhesion	8.519724e-05	3.824205	8.519724e-05
G0:0002009	morphogenesis of an epithelium	1.396681e-04	3.653886	1.396681e-04
G0:0048729	tissue morphogenesis	1.432451e-04	3.643242	1.432451e-04
G0:0007610	behavior	1.925222e-04	3.565432	1.925222e-04
G0:0060562	epithelial tube morphogenesis	5.932837e-04	3.261376	5.932837e-04
G0:0035295	tube development	5.953254e-04	3.253665	5.953254e-04

		q.val	set.size	exp1
G0:0007156	homophilic cell adhesion	0.1951953	113	8.519724e-05
G0:0002009	morphogenesis of an epithelium	0.1951953	339	1.396681e-04
G0:0048729	tissue morphogenesis	0.1951953	424	1.432451e-04
G0:0007610	behavior	0.1967577	426	1.925222e-04
G0:0060562	epithelial tube morphogenesis	0.3565320	257	5.932837e-04
G0:0035295	tube development	0.3565320	391	5.953254e-04

\$less

		p.geomean	stat.mean	p.val
G0:0048285	organelle fission	1.536227e-15	-8.063910	1.536227e-15
G0:0000280	nuclear division	4.286961e-15	-7.939217	4.286961e-15
G0:0007067	mitosis	4.286961e-15	-7.939217	4.286961e-15
G0:0000087	M phase of mitotic cell cycle	1.169934e-14	-7.797496	1.169934e-14
G0:0007059	chromosome segregation	2.028624e-11	-6.878340	2.028624e-11
G0:0000236	mitotic prometaphase	1.729553e-10	-6.695966	1.729553e-10

		q.val	set.size	exp1
G0:0048285	organelle fission	5.841698e-12	376	1.536227e-15
G0:0000280	nuclear division	5.841698e-12	352	4.286961e-15
G0:0007067	mitosis	5.841698e-12	352	4.286961e-15
G0:0000087	M phase of mitotic cell cycle	1.195672e-11	362	1.169934e-14
G0:0007059	chromosome segregation	1.658603e-08	142	2.028624e-11
G0:0000236	mitotic prometaphase	1.178402e-07	84	1.729553e-10

\$stats

		stat.mean	exp1
G0:0007156	homophilic cell adhesion	3.824205	3.824205
G0:0002009	morphogenesis of an epithelium	3.653886	3.653886
G0:0048729	tissue morphogenesis	3.643242	3.643242
G0:0007610	behavior	3.565432	3.565432
G0:0060562	epithelial tube morphogenesis	3.261376	3.261376
G0:0035295	tube development	3.253665	3.253665

Reactome Analysis

Now we do over-representation enrichment analysis and pathway-topology analysis with **Reactome database** using the previous list of significant genes generated from our differential expression results above.

```
sig_genes <- res[res$padj <= 0.05 & !is.na(res$padj), "symbol"]
print(paste("Total number of significant genes:", length(sig_genes)))
```

```
[1] "Total number of significant genes: 8147"
```

```
write.table(sig_genes, file="significant_genes.txt", row.names=FALSE, col.names=FALSE, quote=
```

Q: What pathway has the most significant “Entities p-value”? Do the most significant pathways listed match your previous KEGG results? What factors could cause differences between the two methods?

- The pathway that has the most significant “Entities p-value” is the mitotic phase of the cell cycle. The most significant pathways in the KEGG results were cell cycle, DNA replication, homologous recombination, oocyte meiosis etc while the most significant pathways in the Reactome analysis was Mitotic phase of cell cycle, cell cycle checkpoints, M-phase of mitosis etc. Thus, the most significant pathways in the KEGG results and Reactome results both are closely related to the cell cycle/division and therefore, they closely match. Some factors that could cause differences between the two methods were that KEGG and Reactome databases put the pathways together slightly differently with Reactome database dividing the data into more detailed sub-steps while KEGG results are more broad.