Supplementary Information

Tidal potential measured with distant optical clocks

Lei Hou et al.

Reference numbers follow the main manuscript.

Supplementary Methods

SM1–SM5 describe the filtered validation analysis, also labelled the primary reduction. SM6 and SM7 describe the main analysis, also labelled the auxiliary reduction, unless a filtered-data comparison is explicitly identified. Supplementary Table 4 summarizes their methodological differences.

SM1. Physical model

We use $k_{20} = 0.29525$, $k_{22} = 0.29801$ and $k_3 = 0.093$. The IERS 2010 resonance formula, applied constituent by constituent for k_{21}, gives 0.2588 for K_1, 0.2873 for P_1 and 0.2975 for O_1. The radial displacement is computed with the IERS algorithm, including the frequency and latitude dependence of the Love number h [1].

PREM load Love numbers are used through degree $n = 696$ [32], from the gravity-toolkit distribution [33]. Above this degree, h'_n is joined smoothly to its Boussinesq limit of −6.22, and the series is summed with a Kummer transformation through $n = 40,000$. The convolution uses full grid resolution within 3° of each station and sums blocks of 8×8 cells farther away, which changes the differential M_2 and K_1 loading by at most 0.05%.

The full loading potential exceeds the vertical-displacement contribution $-g\delta h$ by factors of approximately 1.4 for the semidiurnal and 1.8 for the diurnal constituents. The radial displacement from the same convolution agrees with the FES2022 load-tide grid to within 2.2% in amplitude and 1.2° in phase for seven of the eight constituents (Supplementary Fig. 4); the attraction and induced-potential terms use the same PREM load Love numbers. For S_2, whose load-tide grid derives from a different ocean-model release, the amplitude differs by about 18% and the phase at Wuhan by 54°. The S_2 loading is 0.62×10^{-18} in Shanghai and 0.18×10^{-18} in Wuhan, a differential amplitude of 0.63×10^{-18} compared with 1.42×10^{-18} for M_2. Substituting the S_2 loading inferred from the load-tide grid changes the full-template amplitude by 0.001 and the semidiurnal amplitude by 0.000, and leaves the lunar amplitude unchanged.

The Shanghai clock is 22 km from the nearest ocean cell. Filling land cells within 0.3° or 0.6° of the coastline in a $4^\circ \times 5^\circ$ box centred on Shanghai with the nearest ocean values, strips approximately

33–66 km wide, increases the Shanghai M_2 loading potential expressed as $\delta W/g$ from 15.3 mm to 17.0 or 18.5 mm and decreases the lunar response by 0.05 or 0.08, respectively.

Setting the remainder of the eight-constituent model to zero or doubling it gives semidiurnal amplitudes of 0.95–0.96 and lunar amplitudes of 0.95–0.99.

SM2. Selection and timing

Adding the tidal model with factors of −1, 1 or 3 leaves the removed samples unchanged; six robust standard deviations is the lowest tested threshold with this invariance up to three times the model. Block lengths of 60 or 300 s and thresholds from four to eight standard deviations give lunar amplitudes from 0.93 to 1.02. Retaining the excursions gives 0.96 ± 0.14, and the full-template and semidiurnal amplitudes change by less than 0.04. Before transient removal, the procedure reproduces the summer sample set of ref. 23; after removal, the campaign diurnal amplitudes are mutually consistent ($p = 0.21$). Three of the nine identified excursions coincide with approximately equal and opposite link-monitor excursions.

Omitting endpoint trimming changes the lunar amplitude by −0.009 and the full-template amplitude by 0.008. Shifting the model by ±10 min changes A by less than 0.02. Long-period tides are not fitted: the similar site latitudes make the predicted differential semiannual tide only about 2×10^{-20}. Durations refer to retained 1-s samples; the exposure of the filtered data at 600-s steps is slightly shorter.

SM3. Uncertainty calibration and campaign consistency

For the lunar group, two independent sets of 200 realizations give standard deviations of 0.13 and 0.14, compared with 0.13 for the larger of the scaled least-squares and jackknife errors; we quote 0.14, with a resampling bias of +0.01. With the daily disturbances of Methods, which have r.m.s. values up to 1.9×10^{-18}, 5,000 realizations per scenario give standard deviations rounding to 0.14 (0.13–0.15 for the 300 realizations of Extended Data Table 3) and coverage of 0.67–0.68 for ±0.14 and 0.945–0.952 for ±1.96 × 0.14.

For the semidiurnal response, the group-bootstrap standard deviation of 0.08 is smaller than the calibrated value of 0.12, which is quoted. The segment bootstrap gives a lunar amplitude of $0.98 \pm$

0.11 and phase of $-6^\circ \pm 8^\circ$. The correlation between the lunar and neighbouring semidiurnal coefficients is −0.32.

Summer supplies 61% of the conditional information on the lunar response and gives 0.87 ± 0.18, compared with 1.05 ± 0.31 for the earlier campaigns together. The five lunar estimates are consistent ($\chi^2 = 4.1$, four degrees of freedom, $p = 0.40$; Extended Data Table 1), as are the semidiurnal estimates ($\chi^2 = 3.2$, $p = 0.52$; Extended Data Fig. 2). Resampling the operating groups gives a 95% lunar interval of [0.76, 1.18].

SM4. Prediction, environmental and daily-term checks

The whole-template prediction reduces residual sums of squares in seven of eight held-out periods, by 4.9% when pooled; the lunar prediction improves six of eight periods, by 1.7% when pooled.

Over the 98-h overlap, the blackbody shift calculated from chamber temperature and filtered like the ratio has an r.m.s. of 2.8×10^{-18}. As a free regressor it gives 0.92 ± 0.74, or 0.72 ± 0.74 after excluding the chamber-temperature transient on 30 June (day-by-day jackknife errors), consistent with zero. Without real-time correction, the shift would contain a daily component of 2.0×10^{-18} peaking at 12:13 local time, with a peak-to-peak daily range of 4×10^{-18}; the fitted 24-h ratio term in the same overlap is $(0.9 \pm 1.1) \times 10^{-18}$ with maximum at 10:54 ± 5.0 h.

On the primary sampling after segment-mean removal, the predicted solid-tide and loading r.m.s. fractional frequencies are 3.02×10^{-18} and 1.35×10^{-18}, and that of their sum is 4.09×10^{-18}. The response fitted to the solid-Earth tide alone is 1.19 ± 0.20, compared with 1.32 expected on this sampling for the combined model, and the separately fitted loading coefficient is 0.89 ± 0.55. This coefficient comes from a fit of the solid-Earth and loading templates with two free coefficients and segment offsets to the filtered primary data with equal weights; its uncertainty is the larger of the scaled least-squares and day-by-day jackknife errors.

The summer 24-h term has amplitude $(2.8 \pm 1.3) \times 10^{-18}$ and peaks at 15:48 local time with a 1.5-h uncertainty; the 12-h term is $(0.6 \pm 0.7) \times 10^{-18}$. With summer projection coefficients of 0.016 and 0.043 per 10^{-18}, they shift the lunar response by at most 0.045 and 0.024 (Extended Data Table 1). For the full record, a common-phase 12-h perturbation of 10^{-18} changes the whole-template response by at most 0.087.

The lunar responses for alternative daily models (Fig. 4a) are: campaign-specific 24-h terms with air-temperature phases, 0.99 ± 0.14; free-phase campaign-specific 24-h terms, 0.97 ± 0.15; air-temperature anomalies at both sites, 0.98 ± 0.13; common 24-h and 12-h terms, 1.00 ± 0.13; campaign-specific free-phase 12-h terms, 0.82 ± 0.17, or 0.86 ± 0.18 with 24-h terms also included. Operating-group or period offsets in place of segment offsets give 0.97 ± 0.11 and 0.99 ± 0.11, and omitting each earlier campaign in turn gives 0.89–0.99.

SM5. Alternative processing and spectra

The segment-wise estimator of ref. 23 demeans filtered data and model within each segment, fits the amplitude through the origin using non-overlapping 1,200-s windows, and combines segments with inverse-variance weights. For the summer record it gives $A = 0.78 \pm 0.19$ with the present physical model and 0.87(15) with the model of ref. 23; the joint lunar fit gives 0.87 ± 0.18.

Independent processing of the 2026 records uses non-overlapping 600-s means, corrected clock systematic shifts and fixed ten-day calendar baselines. It gives a whole-template response of 0.88 ± 0.10 without empirical daily terms and a lunar response of 0.89 ± 0.28 with daily terms in each ten-day block, and its whole-template prediction improves all six held-out 2026 periods. Simulations with white, Ornstein–Uhlenbeck (correlation times of 1 h to 3 d), Student-t and moving-average noise give biases below 0.04 and 95%-interval coverage of 94.5–99.8% (Supplementary Table 3). Simpler solid-tide approximations change its whole-template scale by less than 0.01.

At M_2, the harmonic power of the full record after projection of segment offsets decreases by 99% after subtraction of the fitted whole template (Extended Data Fig. 1c). In the three-group fit the diurnal coefficient is 0.35 ± 0.51; the two-band estimate is 0.36 ± 0.50 (Extended Data Fig. 2b).

SM6. Auxiliary response and uncertainty details

Supplementary Table 4 compares the main and filtered validation analyses; Supplementary Table 2 gives the main response estimates and their uncertainties. In the phase-relaxed fit, the lunar in-phase and quadrature estimates are 1.0007 ± 0.1320 and -0.0969 ± 0.1989; the scaled least-squares quadrature error is 0.1273.

The projected solid-Earth/loading template correlation is 0.725 and the coefficient correlation under day deletion is −0.888. For M_2 and S_2, at the measured amplitudes 4.5047×10^{-18} and 1.9214×10^{-18}, observing-period noise propagation gives standard errors of 0.6153×10^{-18} and 0.6788×10^{-18}. For the diurnal-group and loading coefficients, the period-noise standard errors are 0.3631 and 0.4337 and the UTC-day-jackknife errors 0.5381 and 0.5438.

Propagating the observing-period noise coefficients, $\Sigma_{kk} = \alpha^2_{p(k)}/\Delta t_k$, through the unchanged estimators gives S/R values of 7.32, 2.83, 1.59 and 1.12 for M_2, S_2, the diurnal group and ocean loading. In the triangular-filter reduction, the corresponding ratios are 6.93 for the lunar group, 3.10 for the neighbouring semidiurnal group and 0.72 for the two-band diurnal estimate.

SM7. Compensation, alias and correlation diagnostics

For donor D and target T, the joint fit supplies the fixed donor coefficient vector $\hat{\beta}_D$. For 61 values of λ from 0 to 1.5, we form $y_\lambda = y - \lambda X_D \hat{\beta}_D$ and fit only the target pair together with N. With $P_T = R_N X_T$ and $P_D = R_N X_D$,

$$\hat{\beta}_T(\lambda) = \hat{\beta}_{T,\text{single}} - \lambda P_T^+ P_D \hat{\beta}_D,$$

where P_T^+ is the Moore–Penrose pseudoinverse. We repeat the sweep with the donor's single-frequency estimate. A fit retaining both pairs after subtraction leaves the target unchanged by construction and is an algebraic control. Amplitude is the norm of the two coefficients and phase is $\text{atan2}(\beta_{\text{sin}}, \beta_{\text{cos}})$, corresponding to $A\cos(2\pi f(t-t_0)-\phi)$. Amplitudes are converted to equivalent height as in Methods.

Template lag correlations evaluate the fitted donor waveform at $t-\ell$ and the target at t, residualize both on the observation support and normalize their inner product, for lags within ±24 h in 5-min steps. The zero-lag correlation is 0.1955, and the maximum absolute lag correlations are 0.2413 for shifted M_2 against S_2 and 0.2113 in the reverse direction. The largest cross-component coefficient correlations between the joint M_2 and S_2 estimates, $R_{ij} = C_{ij}/\sqrt{C_{ii}C_{jj}}$, are 0.248 under scaled least squares and 0.261 under UTC-day deletion (Extended Data Fig. 6).

For alias diagnostics, we scan 20–25 μHz with spacing $1/(12T_{\mathrm{span}})$ and, for each donor, select the three strongest local maxima of its pure-injection least-squares power at least $2/T_{\mathrm{span}}$ from its own frequency. At each selected frequency, the fitted coefficient vector follows $\hat{b}_f(\lambda) = \hat{b}_f(0) - \lambda\hat{b}_{f,D}$, and the equivalent-height amplitude is $K\|\hat{b}_f(0) - \lambda\hat{b}_{f,D}\|$ with $K = (c^2/g_0)\,10^{-16}$ cm. At 22.1016 and 22.6283 μHz, subtraction of the jointly fitted lunar waveform reduces the amplitudes from 3.149 to 0.204 cm and from 3.307 to 0.266 cm, reductions of 93.51% and 91.97%, with minima at compensation factors 0.9974 and 1.0490 (Supplementary Fig. 2).

Supplementary Tables

Supplementary Table 1 | Free sinusoidal coefficients in the auxiliary reduction. Joint M_2 and S_2 cosine–sine coefficients, after controlling 54 segment constants and freely phased N_2 and K_2 terms. Coefficients and standard errors are in units of 10^{-18}. Phases refer to 6 September 2025, 16:05 UTC. Scaled least-squares errors use the residual variance; UTC-day jackknife errors delete each of the 43 observed dates and refit all coefficients.

Coefficient	Estimate	Scaled least-squares standard error	UTC-day jackknife standard error
M_2 cosine	-1.667	0.688	0.880
M_2 sine	-4.185	0.679	0.635
S_2 cosine	-0.460	0.738	0.701
S_2 sine	-1.866	0.750	1.043

Supplementary Table 2 | Template-response signal-to-noise ratios. The first four rows use the non-overlapping full-template or three-group fits; their empirical errors are the larger of scaled-least-squares and UTC-day-jackknife errors. The final two rows use the equal-weight complete solid-Earth/loading fit with UTC-day-jackknife errors. The common-white-noise calculation retains each estimator and uses $\alpha = 5.4 \times 10^{-16}\sqrt{\text{s}}$. The dimensionless factor η_i is defined in Methods.

Fit and coefficient	$\hat{A}_i \pm u_i$	ρ_i	$\rho_{i,\text{white}}$	η_i
Full tide, one template	0.955 ± 0.108	8.83	9.79	0.865
Lunar group, three-group fit	1.000 ± 0.132	7.58	7.50	0.840
Near-12-h group (S_2, K_2)	0.925 ± 0.365	2.54	3.14	0.856
Diurnal group, three-group fit	0.579 ± 0.538	1.08	1.55	0.740
Solid Earth, two-template fit	1.082 ± 0.322	3.36	5.81	0.611
Ocean loading, two-template fit	0.486 ± 0.544	0.89	1.17	0.612


Supplementary Table 3 | Noise-model simulations of the alternative processing. Bias and coverage of nominal 95% intervals based on the largest of the least-squares, delete-day and delete-period standard errors. T denotes the full template without daily terms, and L the lunar group with daily terms, using the ten-day calendar-baseline design in Extended Data Table 2 and the actual observation times. Gaussian scenarios contain 1,024 simulations; Student-t and moving-average scenarios contain 4,096. OU denotes Ornstein–Uhlenbeck noise and MA(1) a first-order moving-average process.

Noise model	Bias (T)	Coverage (T), %	Bias (L)	Coverage (L), %
White	−0.000	96.7	−0.001	98.3
White + OU, 1 h	−0.007	94.5	−0.002	96.9
White + OU, 12 h	−0.002	96.1	+0.031	98.8
White + OU, 3 d	+0.003	95.9	+0.004	99.8
White + period constants	+0.001	96.0	+0.004	98.3
Student-t_7 MA(1)	+0.002	98.3	+0.004	98.2
Student-t_5 MA(1)	+0.001	98.9	+0.003	98.3
Student-t_7 MA(1) + OU	+0.001	97.2	+0.005	98.5


Supplementary Table 4 | Main and filtered validation analyses. Both reductions use the same 54 observing segments, each with its own offset. They differ in selection, filtering, tidal-template realization and weighting, so their estimates are related but are not independent measurements.

	Filtered validation (primary)	Main analysis (auxiliary)
Data	1-s samples after transient screening; 544 h	3,299 non-overlapping 600-s means after segment-median counter-jump rejection and per-segment wall-clock binning (Methods), without the additional running-median transient screening; 545.9 h
Filtering	1,200-s triangular window at 600-s steps, applied to data and model	None
Tidal template	IERS 2010 solid tide with constituent-dependent Love numbers; ocean-loading potential by Green's-function convolution of FES2022 (SM1)	IERS radial displacement converted with h_2 and k_2; effective-potential loading convolution with 18 constituents (Methods)
Weights	Equal	Observing-period noise for the full-template and three-group fits; equal for spectral scans, joint sinusoid fits and the solid-Earth/loading fit
Uncertainties	Residual resampling for the full template, semidiurnal band and lunar group; larger of scaled least squares and day-by-day jackknife otherwise	Larger of scaled least squares and UTC-day jackknife
Used in	Fig. 1b,d, Figs. 2 and 4, Extended Data Figs. 1–3 and 7, Extended Data Tables 1–3, Supplementary Figs. 1 and 3	Fig. 3, Table 1, Extended Data Figs. 4–6, Supplementary Fig. 2, Supplementary Tables 1 and 2


Supplementary Figures



Supplementary Fig. 1 | April campaign in calendar time and frequency. a, Frequency-ratio observations after the 1,200-s triangular filter and removal of fitted segment offsets, their 3-h averages, and predictions with and without ocean loading. The horizontal axis retains actual calendar dates and gaps (local time, UTC+8). **b,** Sinusoidal power before and after subtracting the model, expressed as amplitude squared divided by two on a linear vertical axis, in units of 10^{-36}. The dashed curve is the 99% level at each frequency; the dotted line is the 99% level for the maximum over 15–50 μHz, both obtained from 1,000 resampled residual series. In the April record, K_1, P_1 and O_1 merge into a diurnal band and M_2, S_2 and N_2 into a semidiurnal band. Power below 8 μHz exceeds the displayed range.




Supplementary Fig. 2 | Sampling aliases and amplitude response to compensation. **a,b,** Equivalent-height amplitude spectra of the observations, of the observations after subtraction of the jointly fitted donor waveform, and of that donor alone, on identical timestamps and nuisance projection, with 0–6 cm zooms below. Coloured vertical lines mark three off-centre candidate frequencies selected from the pure-injection response. **c,d,** Observed amplitude at those frequencies as the compensation factor varies. Amplitude axes are linear and share scales within rows. Amplitudes at the controlled N_2 and K_2 frequencies are undefined, and responses close to them are weakly constrained.



177

178 **Supplementary Fig. 3 | Transient excursions and sample exclusion. a,** Sixty-second averages around the
179 excursion on 7 July 2026; red points mark averages consisting mainly of samples excluded by the criterion
180 in Methods. **b,** Distribution of deviations of 300-s means from a running median over ±30 min, divided by
181 the robust standard deviation within each segment. The dashed line marks the threshold of six. The count
182 axis is logarithmic. **c,** Observation time excluded in each campaign (summer in red).



Supplementary Fig. 4 | Radial-displacement validation of the loading convolution. a,b, Amplitude ratios (**a**) and phase differences (**b**) between radial displacements calculated with the loading Green's function and the FES2022 load-tide grid in the centre-of-mass frame, at Shanghai and Wuhan, for eight constituents. The S_2 difference is discussed in Supplementary Methods SM1.