


GDB-9 & Small Molecule Universes

Stefan O. Gugler


Massachusetts Institute of Technology
Department of Chemical Engineering

May 31, 2018



Reduction of the full space



Reduction of the full space



Reduction of the full space



Full Space → Scored Space

- charge $\in [-2, +2]$
- H atoms $\in [0, 4]$
- element $\in \{\text{C, N, O, P, S}\}$

Full Space:

M:125, D:5625, B:5625



} ① charge, sterics, octet, shell, bond order

Scored Space:

50, 1500, 201

These rules ① are demonstrated in the following for the di-heavy-atoms.

Rules ① to reduce Full Space → Scored Space

Constraints:

- Charge $c = c_1 + c_2 \leq 1$
- Sterics: H atoms < 4 on connecting atoms
- Closed shell (even number of electrons)

→ Reduces 5625 ligands to 1171.

Rules ① to reduce Full Space → Scored Space

$$u_{\text{octet},i} = \begin{cases} 10 + 2 \cdot (8 - VE_i) & \text{if } 8 - VE_i < 0 \\ 10 - 1 \cdot (8 - VE_i) & \text{if } 8 - VE_i \geq 0 \end{cases} \quad (1)$$

$$u_{\text{charge}} = \begin{cases} 0 & \text{if } c_1 + c_2 > 0 \\ 3 & \text{if } 0 \geq c_1 + c_2 \geq -2 \\ 1 & \text{if } c_1 + c_2 = -3 \\ 0 & \text{if } c_1 + c_2 = -4 \end{cases} \quad (2)$$

$$u_{\text{VSEPR}} = 5 - | \underbrace{(VE_1 - 2 \cdot LP_1 + c_1 - 2 \cdot h_1)}_{\text{ready electrons}} - \underbrace{(VE_2 - 2 \cdot LP_2 + c_2 - 2 \cdot h_2)}_{\text{ready electrons}} | \quad (3)$$

Rules ① to reduce Full Space → Scored Space



$$u_{\text{CA}} = \begin{cases} 1 & \text{if } h_1 = 4 \\ 2 & \text{if } h_1 = 3 \\ 3 & \text{if else} \end{cases} \quad (4)$$

$$u_{\text{shell}} = \begin{cases} 1 & \text{if } \neg(VE_1 + VE_2) \% 2 \\ 0 & \text{if else} \end{cases} \quad (5)$$

$$u_{\text{total}} = u_{\text{shell}} \cdot \left(\frac{1}{2}(u_{\text{octet},1} + u_{\text{octet},2}) + u_{\text{charge}} + u_{\text{VSEPR}} + u_{\text{CA}} \right) \quad (6)$$

where VE_i denotes the number of valence electrons, LP_i the number of lone pairs.

Distribution and Examples



Reduction of the full small ligand universe: Mo, Di, Bi

Scored Space:
50, 1500, 201



} ② score > s

Desired Space:
29, 374, 60

Reduction of the full small ligand universe: Mo, Di, Bi

Desired Space:
29, 374, 60



} ③ $p\%$ of each isoelectronic

Enumerated Space:
20, 80, 50

Reduction of the full small ligand universe: Mo, Di, Bi

Enumerated Space:

20, 80, 50



④

random whim?

Sample & Build



Subsets of octahedral space

The sizes of the selected subsets of octahedral space.

Set	description	size
Homoleptics	$\text{eq} = \text{ax}$	553
"5+1" symmetric	$\text{eq} = \text{ax}_1 \neq \text{ax}_2$	163,620
"4+2" symmetric	$\text{eq}_1 \neq \text{eq}_2 = \text{ax}$	185,376
Strongly symmetric	$\text{eq} \neq \text{ax}$	245,316
Equatorially asymmetric	$\text{eq}_1 \neq \text{eq}_2 \neq \text{ax}$	15,924,796
Weakly symmetric	$\text{eq} \neq \text{ax}_1 \neq \text{ax}_2$	45,077,310
Complete Heteroleptics	$L_i \neq L_j$	$\approx 5.9 \cdot 10^{12}$
Octahedral Space	all	$> 1.8 \cdot 10^{14}$


Properties of the sets

- Reduce space to facilitate sampling from non-homoleptics
- Example: strongly symmetric, monodentate ligand fields (163,620)
- Exclude all with charge smaller than -4, which results in 87,150 ligand fields (53 %).



Principal Component Analysis

The homoleptics (ho) span the strong symmetry (ss) and "5+1" (fo) set.




Footprint and Entropy calculation

We use five properties to characterize the ligand field and generate a five dimensional distribution:

- total charge
- total valence electrons
- electronegativity of the connecting atom
- $\chi_{\text{ax,eq}}^{\text{lc}} = \sum EN_{\text{CA}} \cdot EN_i$
- $\chi'_{\text{ax,eq}}^{\text{lc}} = \sum EN_{\text{CA}} - EN_i$



We then calculate the entropy, H_{KDE} , of the Kernel Density Estimated distribution.

Correlation analysis for strongly symmetric monodentates



Example of KDE slice

Dimensions $\frac{lc}{ax, eq} \chi_1$ vs. charge in H_{KDE} for strongly symmetric monodentates.




Monodentate Footprints





Table : Entropic footprint

Set	$H_{\text{KDE}}^{\text{monodent}}$	$H_{\text{KDE}}^{\text{bident}}$
Homoleptics	19.7	15.63
"5+1" symmetric	13.7	-
Strongly symmetric AC	-	9.47
Strongly symmetric ADC	12.70	5.53
"4+2" symmetric	12.70	9.47
Weakly symmetric	8.1	7.7
Equatorially asymmetric AC	-	10.04
Equatorially asymmetric ADC		

Entropy histogram



Actual calculations



- 0: $[NH_3]$, $[N]\#[N]$, $[C+]\#[O^-]$, $[C+]\#[NH^-]$, $[N]\#[CH]$
- 1: $[CH_+] = [CH_3^-]$, $[NH] = [O]$, $[CH_+] = [OH^-]$, $[CH_2^+] - [OH_2^-]$
- 8: $[OH_2^-] - [PH^+]$, $[P^+] = [OH^-]$, $[PH^+] - [OH_2^-]$, $[NH_2^-] = [CH^+]$