

# SU(2) NJL model: vacuum sector and meson properties

Hui-Jae Lee,<sup>1,\*</sup> Ho-Yeon Won,<sup>1</sup> and Yu-Son Jun<sup>1</sup>

<sup>1</sup>*Hadron Theory Group, Department of Physics,  
Inha University, Incheon 22212, Republic of Korea*

## I. MESONIC PROPERTIES

The action  $S$  in this process is

$$S = -N_c \text{Tr} [\log D] + \int d^4x \frac{1}{2G} (\sigma^2 + \vec{\pi}^2) \quad (1)$$

$$= -\frac{1}{2} N_c \text{Tr} [\log D^\dagger D] + \int d^4x \frac{1}{2G} (\sigma^2 + \vec{\pi}^2) \quad (2)$$

where  $D^\dagger D$  is

$$D^\dagger D = -\partial^2 + \sigma^2 + \vec{\pi}^2 - \{\not{\partial}(\sigma + i\vec{\pi} \cdot \vec{\tau}\gamma_5)\} = V^{-1} + A \quad (3)$$

and  $V^{-1} = -\partial^2 + M^2$ ,  $A = \sigma^2 + \vec{\pi}^2 - M^2 - \{\not{\partial}(\sigma + i\vec{\pi} \cdot \vec{\tau}\gamma_5)\}$ . By the expansion for log,

$$\begin{aligned} \text{Tr} [\log (V^{-1} + A)] &\approx \text{Tr} [\log (V^{-1})] - \frac{1}{2} \text{Tr} [V A V A] + 8I_1(M) \int d^4x (\sigma^2 + \vec{\pi}^2 - M^2) \\ \implies S &= -\frac{1}{2} N_c \left[ \text{Tr} [\log (V^{-1})] - \frac{1}{2} \text{Tr} [V A V A] + 8I_1(M) \int d^4x (\sigma^2 + \vec{\pi}^2 - M^2) \right] + \int d^4x \frac{1}{2G} ((\sigma - \hat{m})^2 + \vec{\pi}^2) \end{aligned} \quad (4)$$

We have to calculate  $\text{Tr} [V A V A]$  hand by hand.

### A. Calculation of $\text{Tr} [V A V A]$

$$\begin{aligned} \text{Tr} [V A V A] &= \text{Tr} [V (\sigma^2 + \vec{\pi}^2 - M^2 - \{\not{\partial}(\sigma + i\vec{\pi} \cdot \vec{\tau}\gamma_5)\}) V (\sigma^2 + \vec{\pi}^2 - M^2 - \{\not{\partial}(\sigma + i\vec{\pi} \cdot \vec{\tau}\gamma_5)\})] \\ &= \text{Tr} \left[ V \left( 2\sigma_0 \tilde{\sigma} + 2\vec{\pi}_0 \tilde{\vec{\pi}} + \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 - \{\not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5)\} \right) V \left( 2\sigma_0 \tilde{\sigma} + 2\vec{\pi}_0 \tilde{\vec{\pi}} + \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 - \{\not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5)\} \right) \right] \\ &= \text{Tr} \left[ V \left( 2\sigma_0 \tilde{\sigma} + 2\vec{\pi}_0 \tilde{\vec{\pi}} + \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 - \{\not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5)\} \right) V \left( 2\sigma_0 \tilde{\sigma} + 2\vec{\pi}_0 \tilde{\vec{\pi}} + \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 - \{\not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5)\} \right) \right] \\ &= \text{Tr} \left[ 4\sigma_0^2 V \tilde{\sigma} V \tilde{\sigma} + 2i\sigma_0 V \tilde{\sigma} V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} + 2i\sigma_0 V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} V \tilde{\sigma} - V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} \right. \\ &\quad \times V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} + 2\sigma_0 V \tilde{\sigma} V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) + 2\sigma_0 V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) V \tilde{\sigma} + iV \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} \\ &\quad \left. + iV \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) + V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) \right] \end{aligned} \quad (5)$$

We can erase terms using relations  $\text{tr}[\gamma^\mu] = 0$  and  $\text{tr}[\gamma^\mu \gamma_5] = 0$  in the spin space.

$$\begin{aligned} \text{Tr} [V A V A] &= \text{Tr} \left[ 4\sigma_0^2 V \tilde{\sigma} V \tilde{\sigma} - V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} + 2\sigma_0 \left\{ V \tilde{\sigma}, V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) \right\} \right. \\ &\quad \left. + V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) \right]. \end{aligned} \quad (6)$$

---

\* hjlee6674@inha.edu

To treat second term, following relation is needed.

$$\begin{aligned}\gamma_5 \not{\tilde{\pi}}(\tilde{\pi} \cdot \tilde{\tau}) \gamma_5 \not{\tilde{\pi}}(\tilde{\pi} \cdot \tilde{\tau}) &= -\not{\tilde{\pi}}(\tilde{\pi} \cdot \tilde{\tau}) \not{\tilde{\pi}}(\tilde{\pi} \cdot \tilde{\tau}) = -\partial_\mu \gamma^\mu \tilde{\pi}_i \tau_i \partial_\nu \gamma^\nu \tilde{\pi}_j \tau_j = -\partial_\mu \gamma^\mu \tilde{\pi}_i \partial_\nu \gamma^\nu \tilde{\pi}_j (\delta_{ij} + i\epsilon_{ijk} \tau_k) \\ &= -\partial_\mu \gamma^\mu \tilde{\pi}_i \partial_\nu \gamma^\nu \tilde{\pi}_i - \epsilon_{ijk} \partial_\mu \gamma^\mu \tilde{\pi}_i \partial_\nu \gamma^\nu \tilde{\pi}_j \tau_k\end{aligned}\quad (7)$$

Because this is in the trace and  $\text{tr}[\tau_i]=0$ , last term is a zero. Therefore

$$\gamma_5 \not{\tilde{\pi}}(\tilde{\pi} \cdot \tilde{\tau}) \gamma_5 \not{\tilde{\pi}}(\tilde{\pi} \cdot \tilde{\tau}) = -\not{\tilde{\pi}} \cdot \not{\tilde{\pi}} \quad (8)$$

$$\begin{aligned}\text{Tr}[VAV A] &= \text{Tr} \left[ 4\sigma_0^2 V \tilde{\sigma} V \tilde{\sigma} - V \not{\tilde{\sigma}} V \not{\tilde{\sigma}} - V \not{\tilde{\pi}} \cdot V \not{\tilde{\pi}} - 2i\gamma_5 V \not{\tilde{\sigma}} \cdot \not{\tilde{\pi}}(\tilde{\pi} \cdot \tilde{\tau}) + 2\sigma_0 \left\{ V \tilde{\sigma}, V(\tilde{\sigma}^2 + \tilde{\pi}^2) \right\} \right. \\ &\quad \left. + V(\tilde{\sigma}^2 + \tilde{\pi}^2) V(\tilde{\sigma}^2 + \tilde{\pi}^2) \right].\end{aligned}\quad (9)$$

Since  $\text{tr}[\gamma_5] = 0$ ,

$$\begin{aligned}\text{Tr}[VAV A] &= \text{Tr} \left[ 4\sigma_0^2 V \tilde{\sigma} V \tilde{\sigma} - V \not{\tilde{\sigma}} V \not{\tilde{\sigma}} - V \not{\tilde{\pi}} \cdot V \not{\tilde{\pi}} + 2\sigma_0 \left\{ V \tilde{\sigma}, V(\tilde{\sigma}^2 + \tilde{\pi}^2) \right\} \right. \\ &\quad \left. + V(\tilde{\sigma}^2 + \tilde{\pi}^2) V(\tilde{\sigma}^2 + \tilde{\pi}^2) \right].\end{aligned}\quad (10)$$

From  $\phi = (\sigma, \vec{\pi})$ ,

$$\begin{aligned}\phi &= \phi_0 + \tilde{\phi} = (\sigma_c = M, 0) + (\tilde{\sigma}, \tilde{\pi}) \\ \implies \phi_0 \tilde{\phi} &= \sigma_0 \tilde{\sigma}, \quad \tilde{\phi} \tilde{\phi} = \tilde{\sigma}^2 + \tilde{\pi}^2\end{aligned}\quad (11)$$

substituting Eq (10),

$$\text{Tr}[VAV A] = \text{Tr} \left[ 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \not{\tilde{\phi}} V \not{\tilde{\phi}} + 2 \left\{ V \phi_0 \tilde{\phi}, V \tilde{\phi} \tilde{\phi} \right\} + V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} \right]. \quad (12)$$

## B. Action and Propagator

Therefore the action Eq (4) is

$$\begin{aligned}S &= -\frac{1}{2} N_c \left[ \text{Tr} [\log(V^{-1})] - \frac{1}{2} \text{Tr} \left[ 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \not{\tilde{\phi}} V \not{\tilde{\phi}} + 2 \left\{ V \phi_0 \tilde{\phi}, V \tilde{\phi} \tilde{\phi} \right\} + V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} \right] \right. \\ &\quad \left. + 8I_1(M) \int d^4x (\sigma^2 + \vec{\pi}^2 - M^2) \right] + \int d^4x \frac{1}{2G} \left( (\sigma - \hat{m})^2 + \vec{\pi}^2 \right)\end{aligned}\quad (13)$$

We already know from the previous calculation in the paper that

$$8\sigma_c G N_c I_1(\sigma_c) = \sigma_c - \hat{m}. \quad (14)$$

Then last two terms are

$$\begin{aligned}&-4N_c I_1(M) \int d^4x (\sigma^2 + \vec{\pi}^2 - M^2) + \int d^4x \frac{1}{2G} \left( (\sigma - \hat{m})^2 + \vec{\pi}^2 \right) \\ &= -\frac{M - \hat{m}}{2MG} \int d^4x (\phi^2 - M^2) + \int d^4x \frac{1}{2G} (\phi^2 - 2\sigma\hat{m} + \hat{m}^2) \\ &= \left( \frac{\hat{m}}{2MG} - \frac{1}{2G} \right) \int d^4x (\phi^2 + 2\phi\tilde{\phi} + \tilde{\phi}^2 - M^2) + \frac{1}{2G} \int d^4x (\phi^2 - 2\sigma\hat{m} + \hat{m}^2) \\ &= \left( \frac{\hat{m}}{2MG} - \frac{1}{2G} \right) \int d^4x (2M\tilde{\sigma} + \tilde{\phi}^2) + \frac{1}{2G} \int d^4x (M^2 + 2M\tilde{\sigma} + \tilde{\phi}^2 - 2\sigma\hat{m} + \hat{m}^2) \\ &= \int d^4x \left( \frac{\hat{m}\tilde{\sigma}}{G} + \frac{\hat{m}}{2MG} \tilde{\phi}^2 \right) - \int d^4x \left( \frac{M\tilde{\sigma}}{G} + \frac{1}{2G} \tilde{\phi}^2 \right) + \int d^4x \left( \frac{M^2}{2G} + \frac{M\tilde{\sigma}}{G} + \frac{1}{2G} \tilde{\phi}^2 - \frac{M\hat{m}}{G} + \frac{\tilde{\sigma}\hat{m}}{G} + \frac{\hat{m}^2}{2G} \right) \\ &= \int d^4x \left( \frac{2\hat{m}\tilde{\sigma} - M\hat{m}}{G} + \frac{M^2 + \hat{m}^2}{2G} \right) + \int d^4x \frac{\hat{m}}{2MG} \tilde{\phi}^2.\end{aligned}\quad (15)$$

We have to keep in mind the last term which will be included the second derivative of the action. Let us write the action one more time:

$$S = -\frac{1}{2}N_c \left[ \text{Tr} [\log (V^{-1})] - \frac{1}{2} \text{Tr} \left[ 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} + 2 \left\{ V \phi_0 \tilde{\phi}, V \tilde{\phi} \tilde{\phi} \right\} + V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} \right] \right] \\ + \int d^4x \left( \frac{2\hat{m}\tilde{\sigma} - M\hat{m}}{G} + \frac{M^2 + \hat{m}^2}{2G} \right) + \int d^4x \frac{\hat{m}}{2MG} \tilde{\phi}^2, \quad (16)$$

$$S = -\frac{1}{2}N_c \left[ \text{Tr} [\log (V^{-1})] - \frac{1}{2} \text{Tr} \left[ 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} + 2 \left\{ V \phi_0 \tilde{\phi}, V \tilde{\phi} \tilde{\phi} \right\} + V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} \right] \right] + R_1 + R_2 \tilde{\phi}^2. \quad (17)$$

Since we use saddle-point approximation, the action can be expressed the expansion for  $\phi$  with a zero first derivative:

$$S[\phi] \approx S[\phi_0] + \frac{\delta}{\delta\phi} S[\phi_0] \tilde{\phi} + \frac{1}{2} \frac{\delta^2}{\delta\phi^2} S[\phi_0] \tilde{\phi}^2 \\ = S[\phi_0] + \frac{1}{2} \frac{\delta^2}{\delta\phi^2} S[\phi_0] \tilde{\phi}^2. \quad (18)$$

Comparing with Eq (17), we can find the second derivative term

$$\frac{1}{2} \frac{\delta^2}{\delta\phi^2} S[\phi_0] \tilde{\phi}^2 = -\frac{1}{2} N_c \left[ -\frac{1}{2} \text{Tr} \left[ 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} \right] \right] + R_2 \tilde{\phi}^2 \\ = \frac{N_c}{4} \int \frac{d^4p}{(2\pi)^4} \text{tr} \left\langle p \left| 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} \right| p \right\rangle + R_2 \tilde{\phi}^2 \\ = 2N_c \int \frac{d^4p}{(2\pi)^4} \left\langle p \left| 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} \right| p \right\rangle + R_2 \tilde{\phi}^2. \quad (19)$$

Now we will use the completeness relation:

$$\mathbb{I} = \frac{d^4p'}{(2\pi)^4} |p'\rangle \langle p'| \quad (20)$$

$$\implies \int \frac{d^4p}{(2\pi)^4} \left\langle p \left| 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} \right| p \right\rangle = \int \frac{d^4p}{(2\pi)^4} \int \frac{d^4p'}{(2\pi)^4} \left( \left\langle p \left| 4\phi_0^2 V \tilde{\phi} \right| p' \right\rangle \left\langle p' \left| V \tilde{\phi} \right| p \right\rangle \right. \\ \left. - \left\langle p \left| V \tilde{\phi} \tilde{\phi} \right| p' \right\rangle \left\langle p' \left| V \tilde{\phi} \tilde{\phi} \right| p \right\rangle \right) \quad (21)$$

In the momentum space representation,  $V = 1/(p^2 + M^2)$ . So

$$\frac{1}{2} \frac{\delta^2}{\delta\phi^2} S[\phi_0] \tilde{\phi}^2 = 2N_c \int \frac{d^4p}{(2\pi)^4} \int \frac{d^4p'}{(2\pi)^4} \frac{1}{(p^2 + M^2)(p'^2 + M^2)} \left( \left\langle p \left| 4\phi_0^2 \tilde{\phi} \right| p' \right\rangle \left\langle p' \left| \tilde{\phi} \right| p \right\rangle - \left\langle p \left| \tilde{\phi} \tilde{\phi} \right| p' \right\rangle \left\langle p' \left| \tilde{\phi} \tilde{\phi} \right| p \right\rangle \right) + R_2 \tilde{\phi}^2 \quad (22)$$

Let us calculate term by term.

$$\left\langle p \left| 4\phi_0^2 \tilde{\phi} \right| p' \right\rangle = 4M^2 \int d^4x \left\langle p \left| \tilde{\phi} \right| x \right\rangle \langle x | p' \rangle = 4M^2 \int d^4x \tilde{\phi}(x) \langle p | x \rangle \langle x | p' \rangle = 4M^2 \int d^4x \tilde{\phi}(x) e^{ix(p-p')} \\ = 4M^2 \tilde{\phi}(p - p'), \\ \left\langle p' \left| \tilde{\phi} \right| p \right\rangle = \tilde{\phi}(p' - p), \\ \left\langle p \left| \tilde{\phi} \tilde{\phi} \right| p' \right\rangle = \int d^4x \left\langle p \left| \tilde{\phi} \tilde{\phi} \right| x \right\rangle \langle x | p' \rangle = \int d^4x \tilde{\phi} \tilde{\phi}(x) \langle p | x \rangle \langle x | p' \rangle = \int d^4x \tilde{\phi} \tilde{\phi}(x) e^{ix(p-p')} \\ = \int d^4x \tilde{\phi} \left( \tilde{\phi}(x) e^{ix(p-p')} \right) - \int d^4x \tilde{\phi}(x) \tilde{\phi} e^{ix(p-p')} = - \int d^4x \tilde{\phi}(x) i(p - p') e^{ix(p-p')} \\ = -i(p - p') \tilde{\phi}(p - p'), \\ \left\langle p' \left| \tilde{\phi} \tilde{\phi} \right| p \right\rangle = -i(p' - p) \tilde{\phi}(p' - p). \quad (23)$$

Then Eq (22) is

$$\frac{1}{2} \frac{\delta^2}{\delta\phi^2} S[\phi_0] \tilde{\phi}^2 = 2N_c \int \frac{d^4 p}{(2\pi)^4} \int \frac{d^4 p'}{(2\pi)^4} \frac{4M^2 \tilde{\phi}(p-p') \tilde{\phi}(p'-p) - (p-p')(p'-p) \tilde{\phi}(p-p') \tilde{\phi}(p'-p)}{(p^2 + M^2)(p'^2 + M^2)} + R_2 \tilde{\phi}^2. \quad (24)$$

For convenience, let us introduce changing variable:

$$p' + p = 2k, \quad p' - p = q \quad (25)$$

and

$$p' = k + \frac{q}{2}, \quad p = k - \frac{q}{2}. \quad (26)$$

Since we change two variables  $(p', p)$  to  $(k, q)$ , 4-integral variables becomes  $d^4 k d^4 q$ . Hence

$$\begin{aligned} \frac{1}{2} \frac{\delta^2}{\delta\phi^2} S[\phi_0] \tilde{\phi}^2 &= 2N_c \int \frac{d^4 k}{(2\pi)^4} \int \frac{d^4 q}{(2\pi)^4} \frac{4M^2 \tilde{\phi}(-q) \tilde{\phi}(q) + q^2 \tilde{\phi}(-q) \tilde{\phi}(q)}{\left\{ \left(k - \frac{q}{2}\right)^2 + M^2 \right\} \left\{ \left(k + \frac{q}{2}\right)^2 + M^2 \right\}} + R_2 \tilde{\phi}^2 \\ &= 2N_c \int \frac{d^4 k}{(2\pi)^4} \int \frac{d^4 q}{(2\pi)^4} \frac{4M^2 + q^2}{\left\{ \left(k - \frac{q}{2}\right)^2 + M^2 \right\} \left\{ \left(k + \frac{q}{2}\right)^2 + M^2 \right\}} \tilde{\phi}(-q) \tilde{\phi}(q) + R_2 \tilde{\phi}^2 \\ &= 2N_c \int \frac{d^4 q}{(2\pi)^4} (4M^2 + q^2) f(q) \tilde{\phi}(-q) \tilde{\phi}(q) + R_2 \tilde{\phi}^2, \quad f(q) = \int \frac{d^4 k}{(2\pi)^4} \frac{1}{\left\{ \left(k - \frac{q}{2}\right)^2 + M^2 \right\} \left\{ \left(k + \frac{q}{2}\right)^2 + M^2 \right\}}. \end{aligned} \quad (27)$$

$\tilde{\phi}(q)$  can be expressed into the Fourier transformation

$$\begin{aligned} \tilde{\phi}(-q) \tilde{\phi}(q) &= \int d^4 x \int d^4 y \tilde{\phi}(y) e^{iq(x-y)} \tilde{\phi}(x) \\ \implies \frac{1}{2} \frac{\delta^2}{\delta\phi^2} S[\phi_0] \tilde{\phi}^2 &= 2N_c \int d^4 x \int d^4 y \int \frac{d^4 q}{(2\pi)^4} (4M^2 + q^2) f(q) \tilde{\phi}(y) e^{iq(x-y)} \tilde{\phi}(x) \\ &\quad + \int d^4 x \int d^4 y \frac{\hat{m}}{2MG} \delta^{(4)}(x-y) \tilde{\phi}(y) \tilde{\phi}(x). \end{aligned} \quad (28)$$

Actually, the exact expression of the second derivative is

$$\frac{1}{2} \frac{\delta^2}{\delta\phi^2} S[\phi_0] \tilde{\phi}^2 = \frac{1}{2} \int d^4 x \int d^4 y \left. \frac{\delta^2 S}{\delta\phi(x) \delta\phi(y)} \right|_{\phi_0} \tilde{\phi}(x) \tilde{\phi}(y) \quad (29)$$

and this form is seem to be comparable with the LHS of Eq (28). Therefore

$$\begin{aligned} \left. \frac{\delta^2 S}{\delta\phi(x) \delta\phi(y)} \right|_{\phi_0} &= 2N_c \int \frac{d^4 q}{(2\pi)^4} (4M^2 + q^2) f(q) e^{iq(x-y)} + \frac{\hat{m}}{2MG} \delta^{(4)}(x-y) \\ &= \frac{1}{2} \int \frac{d^4 q}{(2\pi)^4} \left[ 4N_c (4M^2 + q^2) f(q) + \frac{\hat{m}}{MG} \right] e^{iq(x-y)}. \end{aligned} \quad (30)$$

Finally the meson propagator  $K_\phi(q^2)$  is

$$\begin{aligned} \left. \frac{\delta^2 S}{\delta\phi(x) \delta\phi(y)} \right|_{\phi_0} &= \int \frac{d^4 q}{(2\pi)^4} K^{-1}(q^2) e^{iq(x-y)} = \int \frac{d^4 q}{(2\pi)^4} \left[ 4N_c (4M^2 + q^2) f(q) + \frac{\hat{m}}{MG} \right] e^{iq(x-y)} \\ &= K(q^2) = \frac{1}{4N_c (4M^2 + q^2) f(q) + \frac{\hat{m}}{MG}} = \frac{1}{Z_q(q^2)} \frac{1}{4M^2 + q^2 + \frac{\hat{m}}{MG Z_q(q^2)}}, \quad Z_q(q^2) = 4N_c f(q). \end{aligned} \quad (31)$$

### C. $Z_q(q^2)$ Calculation

Now only one process is remain: calculating  $Z_q(q^2)$ . From Eq (27),

$$\begin{aligned} Z_q(q^2) &= 4N_c \int \frac{d^4 k}{(2\pi)^4} \frac{1}{\left\{ \left( k - \frac{q}{2} \right)^2 + M^2 \right\} \left\{ \left( k + \frac{q}{2} \right)^2 + M^2 \right\}} \\ &= 4N_c \int \frac{d^4 k}{(2\pi)^4} \frac{1}{\left( k^2 + M^2 + \frac{q^2}{4} - qk \right) \left( k^2 + M^2 + \frac{q^2}{4} + qk \right)}. \end{aligned} \quad (32)$$

There are useful integral rules:

$$\frac{1}{AB} = \int_0^1 dx \frac{1}{[xA + (1-x)B]^2}, \quad \frac{1}{Q^2} = \int_0^\infty du u e^{-uQ}. \quad (33)$$

Let us  $A = \left( k^2 + M^2 + \frac{q^2}{4} - qk \right)$  and  $B = \left( k^2 + M^2 + \frac{q^2}{4} + qk \right)$ . substituting to the first rule,

$$\begin{aligned} Z_q(q^2) &= 4N_c \int \frac{d^4 k}{(2\pi)^4} \int_0^1 dx \left[ x \left( k^2 + M^2 + \frac{q^2}{4} - qk \right) + (1-x) \left( k^2 + M^2 + \frac{q^2}{4} + qk \right) \right]^{-2} \\ &= 4N_c \int \frac{d^4 k}{(2\pi)^4} \int_0^1 dx \left[ -2qkx + \left( k^2 + M^2 + \frac{q^2}{4} + qk \right) \right]^{-2} \\ &= 4N_c \int \frac{d^4 k}{(2\pi)^4} \int_0^1 dx \frac{1}{(Kx + B)^2} = 4N_c \int \frac{d^4 k}{(2\pi)^4} \int_0^1 dx \frac{1}{(Kx + B)^2} \end{aligned} \quad (34)$$

where  $K = -2qk$  and  $B = k^2 + M^2 + \frac{q^2}{4} + qk$ . By the second rule,

$$\begin{aligned} Z_q(q^2) &= 4N_c \int \frac{d^4 k}{(2\pi)^4} \int_0^1 dx \int_0^\infty du u \exp \left[ -u \left( -2qkx + k^2 + M^2 + \frac{q^2}{4} + qk \right) \right] \\ &= 4N_c \int_0^1 dx \int_0^\infty du u \exp \left[ -\frac{u}{4} \{ 1 - (1-2x)^2 \} q^2 - uM^2 \right] \int \frac{d^4 k}{(2\pi)^4} \exp \left[ -u \left( k + \frac{1}{2}q(1-2x) \right)^2 \right]. \end{aligned} \quad (35)$$

For convenience, let  $1 - 2x = y$ , and  $-2dx = dy$

$$\begin{aligned} Z_q(q^2) &= 4N_c \int_0^1 dx \int_0^\infty du u \exp \left[ -\frac{u}{4} \{ 1 - (1-2x)^2 \} q^2 - uM^2 \right] \int \frac{d^4 k}{(2\pi)^4} \exp \left[ -u \left( k + \frac{1}{2}q(1-2x) \right)^2 \right] \\ &= -2N_c \int_1^{-1} dy \int_0^\infty du u \exp \left[ -\frac{u}{4} (1-y^2) q^2 - uM^2 \right] \int \frac{d^4 k}{(2\pi)^4} \exp \left[ -u \left( k + \frac{1}{2}qy \right)^2 \right] \\ &= 2N_c \int_{-1}^1 dy \int_0^\infty du u \exp \left[ -\frac{u}{4} (1-y^2) q^2 - uM^2 \right] \int \frac{d^4 k}{(2\pi)^4} \exp \left[ -u \left( k + \frac{1}{2}qy \right)^2 \right]. \end{aligned} \quad (36)$$

The last integral is the 4-dimensional spherical integral.

$$\begin{aligned} \int \frac{d^4 k}{(2\pi)^4} \exp \left[ -u \left( k + \frac{1}{2}qy \right)^2 \right] &= \int_0^\infty \frac{dk}{(4\pi)^2} k^3 \exp \left[ -u \left( k + \frac{1}{2}qy \right)^2 \right] \\ &= \int_0^\infty \frac{dk}{(4\pi)^2} \left( k - \frac{1}{2}qy \right)^3 e^{-uk^2} \end{aligned} \quad (37)$$

Expanding quadratic, there are only gaussian integrals. But the terms which have the odd degree of  $y$  are erase since of the integral for  $y$  is symmetric.

$$\begin{aligned}
\int_0^\infty \frac{dk}{(4\pi)^2} \left(k - \frac{1}{2}qy\right)^3 e^{-uk^2} &= \int_0^\infty \frac{dk}{(4\pi)^2} \left(k^3 - \frac{3}{2}qyk^2 + \frac{3}{4}q^2y^2k - \frac{1}{8}q^3y^3\right) e^{-uk^2} \\
&= \int_0^\infty \frac{dk}{(4\pi)^2} \left(k^3 + \frac{3}{4}q^2y^2k\right) e^{-uk^2} \\
&= \frac{1}{(4\pi)^2} \left(\frac{1}{2u^2} + \frac{3q^2y^2}{8u}\right).
\end{aligned} \tag{38}$$

Hence

$$\begin{aligned}
Z_q(q^2) &= \frac{2N_c}{(4\pi)^2} \int_{-1}^1 dy \int_0^\infty du u \exp \left[ -\frac{u}{4} (1-y^2) q^2 - uM^2 \right] \left( \frac{1}{2u^2} + \frac{3q^2y^2}{8u} \right) \\
&= \frac{2N_c}{(4\pi)^2} \int_{-1}^1 dy \int_0^\infty du \left( \frac{1}{2u} e^{-u\left(\frac{q^2}{4}(1-y^2)+M^2\right)} + \frac{3q^2y^2}{8} e^{-u\left(\frac{q^2}{4}(1-y^2)+M^2\right)} \right).
\end{aligned} \tag{39}$$

The last term diverges at  $y^2 = \frac{M^2}{q^2} + 1$ . So we remain only first term and cut the integral interval for  $u$ (regularization). Finally

$$\begin{aligned}
Z_q(q^2) &= \frac{2N_c}{(4\pi)^2} \int_{-1}^1 dy \int_0^\infty du u \exp \left[ -\frac{u}{4} (1-y^2) q^2 - uM^2 \right] \left( \frac{1}{2u^2} + \frac{3q^2y^2}{8u} \right) \\
&= \frac{2N_c}{(4\pi)^2} \int_{-1}^0 dy \int_{\Lambda^{-2}}^\infty du \frac{1}{u} e^{-u\left(M^2 + \frac{q^2}{4}(1-y^2)\right)}.
\end{aligned} \tag{40}$$

This is the Pauli-Vilars regularized expression.