

# SU(2) NJL model: vacuum sector and meson properties

Hui-Jae Lee,<sup>1,\*</sup> Ho-Yeon Won,<sup>1</sup> and Yu-Son Jun<sup>1</sup>

<sup>1</sup>*Hadron Theory Group, Department of Physics,  
Inha University, Incheon 22212, Republic of Korea*

## I. MESONIC PROPERTIES

The action  $S$  in this process is

$$S = -N_c \text{Tr} [\log D] + \int d^4x \frac{1}{2G} (\sigma^2 + \vec{\pi}^2) \quad (1)$$

$$= -\frac{1}{2} N_c \text{Tr} [\log D^\dagger D] + \int d^4x \frac{1}{2G} (\sigma^2 + \vec{\pi}^2) \quad (2)$$

where  $D^\dagger D$  is

$$D^\dagger D = -\partial^2 + \sigma^2 + \vec{\pi}^2 - \{\not{\partial}(\sigma + i\vec{\pi} \cdot \vec{\tau}\gamma_5)\} = V^{-1} + A \quad (3)$$

and  $V^{-1} = -\partial^2 + M^2$ ,  $A = \sigma^2 + \vec{\pi}^2 - M^2 - \{\not{\partial}(\sigma + i\vec{\pi} \cdot \vec{\tau}\gamma_5)\}$ . By the expansion for log,

$$\begin{aligned} \text{Tr} [\log (V^{-1} + A)] &\approx \text{Tr} [\log (V^{-1})] - \frac{1}{2} \text{Tr} [V A V A] + 8I_1(M) \int d^4x (\sigma^2 + \vec{\pi}^2 - M^2) \\ \implies S &= -\frac{1}{2} N_c \left[ \text{Tr} [\log (V^{-1})] - \frac{1}{2} \text{Tr} [V A V A] + 8I_1(M) \int d^4x (\sigma^2 + \vec{\pi}^2 - M^2) \right] + \int d^4x \frac{1}{2G} ((\sigma - \hat{m})^2 + \vec{\pi}^2) \end{aligned} \quad (4)$$

We have to calculate  $\text{Tr} [V A V A]$  hand by hand.

### A. Calculation of $\text{Tr} [V A V A]$

$$\begin{aligned} \text{Tr} [V A V A] &= \text{Tr} [V (\sigma^2 + \vec{\pi}^2 - M^2 - \{\not{\partial}(\sigma + i\vec{\pi} \cdot \vec{\tau}\gamma_5)\}) V (\sigma^2 + \vec{\pi}^2 - M^2 - \{\not{\partial}(\sigma + i\vec{\pi} \cdot \vec{\tau}\gamma_5)\})] \\ &= \text{Tr} \left[ V \left( 2\sigma_0 \tilde{\sigma} + 2\vec{\pi}_0 \tilde{\vec{\pi}} + \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 - \{\not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5)\} \right) V \left( 2\sigma_0 \tilde{\sigma} + 2\vec{\pi}_0 \tilde{\vec{\pi}} + \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 - \{\not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5)\} \right) \right] \\ &= \text{Tr} \left[ V \left( 2\sigma_0 \tilde{\sigma} + 2\vec{\pi}_0 \tilde{\vec{\pi}} + \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 - \{\not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5)\} \right) V \left( 2\sigma_0 \tilde{\sigma} + 2\vec{\pi}_0 \tilde{\vec{\pi}} + \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 - \{\not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5)\} \right) \right] \\ &= \text{Tr} \left[ 4\sigma_0^2 V \tilde{\sigma} V \tilde{\sigma} + 2i\sigma_0 V \tilde{\sigma} V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} + 2i\sigma_0 V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} V \tilde{\sigma} - V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} \right. \\ &\quad \times V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} + 2\sigma_0 V \tilde{\sigma} V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) + 2\sigma_0 V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) V \tilde{\sigma} + iV \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} \\ &\quad \left. + iV \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) + V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) \right] \end{aligned} \quad (5)$$

We can erase terms using relations  $\text{tr}[\gamma^\mu] = 0$  and  $\text{tr}[\gamma^\mu \gamma_5] = 0$  in the spin space.

$$\begin{aligned} \text{Tr} [V A V A] &= \text{Tr} \left[ 4\sigma_0^2 V \tilde{\sigma} V \tilde{\sigma} - V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} V \left\{ \not{\partial}(\tilde{\sigma} + i\tilde{\vec{\pi}} \cdot \vec{\tau}\gamma_5) \right\} + 2\sigma_0 \left\{ V \tilde{\sigma}, V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) \right\} \right. \\ &\quad \left. + V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) V \left( \tilde{\sigma}^2 + \tilde{\vec{\pi}}^2 \right) \right]. \end{aligned} \quad (6)$$

---

\* hjlee6674@inha.edu

To treat second term, following relation is needed.

$$\begin{aligned}\gamma_5 \not{\partial}(\tilde{\pi} \cdot \tilde{\tau}) \gamma_5 \not{\partial}(\tilde{\pi} \cdot \tilde{\tau}) &= -\not{\partial}(\tilde{\pi} \cdot \tilde{\tau}) \not{\partial}(\tilde{\pi} \cdot \tilde{\tau}) = -\partial_\mu \gamma^\mu \tilde{\pi}_i \tau_i \partial_\nu \gamma^\nu \tilde{\pi}_j \tau_j = -\partial_\mu \gamma^\mu \tilde{\pi}_i \partial_\nu \gamma^\nu \tilde{\pi}_j (\delta_{ij} + i\epsilon_{ijk} \tau_k) \\ &= -\partial_\mu \gamma^\mu \tilde{\pi}_i \partial_\nu \gamma^\nu \tilde{\pi}_i - \epsilon_{ijk} \partial_\mu \gamma^\mu \tilde{\pi}_i \partial_\nu \gamma^\nu \tilde{\pi}_j \tau_k\end{aligned}\quad (7)$$

Because this is in the trace and  $\text{tr}[\tau_i]=0$ , last term is a zero. Therefore

$$\gamma_5 \not{\partial}(\tilde{\pi} \cdot \tilde{\tau}) \gamma_5 \not{\partial}(\tilde{\pi} \cdot \tilde{\tau}) = -\not{\partial} \tilde{\pi} \cdot \not{\partial} \tilde{\pi} \quad (8)$$

$$\begin{aligned}\text{Tr}[VAV A] &= \text{Tr} \left[ 4\sigma_0^2 V \tilde{\sigma} V \tilde{\sigma} - V \not{\partial} \tilde{\sigma} V \not{\partial} \tilde{\sigma} - V \not{\partial} \tilde{\pi} \cdot V \not{\partial} \tilde{\pi} - 2i\gamma_5 V \not{\partial} \tilde{\sigma} \cdot \not{\partial} (\tilde{\pi} \cdot \tilde{\tau}) + 2\sigma_0 \left\{ V \tilde{\sigma}, V (\tilde{\sigma}^2 + \tilde{\pi}^2) \right\} \right. \\ &\quad \left. + V (\tilde{\sigma}^2 + \tilde{\pi}^2) V (\tilde{\sigma}^2 + \tilde{\pi}^2) \right].\end{aligned}\quad (9)$$

Since  $\text{tr}[\gamma_5] = 0$ ,

$$\begin{aligned}\text{Tr}[VAV A] &= \text{Tr} \left[ 4\sigma_0^2 V \tilde{\sigma} V \tilde{\sigma} - V \not{\partial} \tilde{\sigma} V \not{\partial} \tilde{\sigma} - V \not{\partial} \tilde{\pi} \cdot V \not{\partial} \tilde{\pi} + 2\sigma_0 \left\{ V \tilde{\sigma}, V (\tilde{\sigma}^2 + \tilde{\pi}^2) \right\} \right. \\ &\quad \left. + V (\tilde{\sigma}^2 + \tilde{\pi}^2) V (\tilde{\sigma}^2 + \tilde{\pi}^2) \right].\end{aligned}\quad (10)$$

From  $\phi = (\sigma, \tilde{\pi})$ ,

$$\begin{aligned}\phi &= \phi_0 + \tilde{\phi} = (\sigma_c = M, 0) + (\tilde{\sigma}, \tilde{\pi}) \\ \implies \phi_0 \tilde{\phi} &= \sigma_0 \tilde{\sigma}, \quad \tilde{\phi} \tilde{\phi} = \tilde{\sigma}^2 + \tilde{\pi}^2\end{aligned}\quad (11)$$

substituting Eq (10),

$$\text{Tr}[VAV A] = \text{Tr} \left[ 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \not{\partial} \tilde{\phi} V \not{\partial} \tilde{\phi} + 2 \left\{ V \phi_0 \tilde{\phi}, V \tilde{\phi} \tilde{\phi} \right\} + V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} \right]. \quad (12)$$

Therefore the action Eq (4) is

$$\begin{aligned}S &= -\frac{1}{2} N_c \left[ \text{Tr} [\log (V^{-1})] - \frac{1}{2} \text{Tr} \left[ 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \not{\partial} \tilde{\phi} V \not{\partial} \tilde{\phi} + 2 \left\{ V \phi_0 \tilde{\phi}, V \tilde{\phi} \tilde{\phi} \right\} + V \tilde{\phi} \tilde{\phi} V \tilde{\phi} \tilde{\phi} \right] \right. \\ &\quad \left. + 8I_1(M) \int d^4x (\sigma^2 + \tilde{\pi}^2 - M^2) \right] + \int d^4x \frac{1}{2G} \left( (\sigma - \hat{m})^2 + \tilde{\pi}^2 \right)\end{aligned}\quad (13)$$

Since we use saddle-point approximation, the action can be expressed the expansion for  $\phi$  with a zero first derivative:

$$\begin{aligned}S[\phi] &\approx S[\phi_0] + \frac{\delta}{\delta \phi} S[\phi_0] \tilde{\phi} + \frac{1}{2} \frac{\delta^2}{\delta \phi^2} S[\phi_0] \tilde{\phi}^2 \\ &= S[\phi_0] + \frac{1}{2} \frac{\delta^2}{\delta \phi^2} S[\phi_0] \tilde{\phi}^2.\end{aligned}\quad (14)$$

Comparing with Eq (13), we can find the second derivative term

$$\begin{aligned}\frac{1}{2} \frac{\delta^2}{\delta \phi^2} S[\phi_0] \tilde{\phi}^2 &= -\frac{1}{2} N_c \left[ -\frac{1}{2} \text{Tr} \left[ 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \not{\partial} \tilde{\phi} V \not{\partial} \tilde{\phi} \right] \right] = \frac{N_c}{4} \int \frac{d^4p}{(2\pi)^4} \text{tr} \left\langle p \left| 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \not{\partial} \tilde{\phi} V \not{\partial} \tilde{\phi} \right| p \right\rangle \\ &= 2N_c \int \frac{d^4p}{(2\pi)^4} \left\langle p \left| 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \not{\partial} \tilde{\phi} V \not{\partial} \tilde{\phi} \right| p \right\rangle.\end{aligned}\quad (15)$$

Now we will use the completeness relation:

$$\mathbb{I} = \frac{d^4p'}{(2\pi)^4} |p'\rangle \langle p'| \quad (16)$$

$$\begin{aligned}\implies \int \frac{d^4p}{(2\pi)^4} \left\langle p \left| 4\phi_0^2 V \tilde{\phi} V \tilde{\phi} - V \not{\partial} \tilde{\phi} V \not{\partial} \tilde{\phi} \right| p \right\rangle &= \int \frac{d^4p}{(2\pi)^4} \int \frac{d^4p'}{(2\pi)^4} \left( \left\langle p \left| 4\phi_0^2 V \tilde{\phi} \right| p' \right\rangle \left\langle p' \left| V \tilde{\phi} \right| p \right\rangle \right. \\ &\quad \left. - \left\langle p \left| V \not{\partial} \tilde{\phi} \right| p' \right\rangle \left\langle p' \left| V \not{\partial} \tilde{\phi} \right| p \right\rangle \right)\end{aligned}\quad (17)$$

In the momentum space representation,  $V = 1/(p^2 + M^2)$ . So

$$\frac{1}{2} \frac{\delta^2}{\delta\phi^2} S[\phi_0] \tilde{\phi}^2 = 2N_c \int \frac{d^4 p}{(2\pi)^4} \int \frac{d^4 p'}{(2\pi)^4} \frac{1}{(p^2 + M^2)(p'^2 + M^2)} \left( \langle p | 4\phi_0^2 \tilde{\phi} | p' \rangle \langle p' | \tilde{\phi} | p \rangle - \langle p | \tilde{\phi} \tilde{\phi} | p' \rangle \langle p' | \tilde{\phi} \tilde{\phi} | p \rangle \right) \quad (18)$$

Let us calculate term by term.

$$\begin{aligned} \langle p | 4\phi_0^2 \tilde{\phi} | p' \rangle &= 4M^2 \int d^4 x \langle p | \tilde{\phi} | x \rangle \langle x | p' \rangle = 4M^2 \int d^4 x \tilde{\phi}(x) \langle p | x \rangle \langle x | p' \rangle = 4M^2 \int d^4 x \tilde{\phi}(x) e^{ix(p-p')} \\ &= 4M^2 \tilde{\phi}(p-p'), \\ \langle p' | \tilde{\phi} | p \rangle &= \tilde{\phi}(p'-p), \\ \langle p | \tilde{\phi} \tilde{\phi} | p' \rangle &= \int d^4 x \langle p | \tilde{\phi} \tilde{\phi} | x \rangle \langle x | p' \rangle = \int d^4 x \tilde{\phi} \tilde{\phi}(x) \langle p | x \rangle \langle x | p' \rangle = \int d^4 x \tilde{\phi} \tilde{\phi}(x) e^{ix(p-p')} \\ &= \int d^4 x \tilde{\phi} \left( \tilde{\phi}(x) e^{ix(p-p')} \right) - \int d^4 x \tilde{\phi}(x) \tilde{\phi} e^{ix(p-p')} = - \int d^4 x \tilde{\phi}(x) i(p-p') e^{ix(p-p')} \\ &= -i(p-p') \tilde{\phi}(p-p'), \\ \langle p' | \tilde{\phi} \tilde{\phi} | p \rangle &= -i(p'-p) \tilde{\phi}(p'-p). \end{aligned} \quad (19)$$

Then Eq (18) is

$$\frac{1}{2} \frac{\delta^2}{\delta\phi^2} S[\phi_0] \tilde{\phi}^2 = 2N_c \int \frac{d^4 p}{(2\pi)^4} \int \frac{d^4 p'}{(2\pi)^4} \frac{4M^2 \tilde{\phi}(p-p') \tilde{\phi}(p'-p) + (p-p')(p'-p) \tilde{\phi}(p-p') \tilde{\phi}(p'-p)}{(p^2 + M^2)(p'^2 + M^2)}. \quad (20)$$

For convenience, let us introduce changing variable:

$$p' + p = 2k, \quad p' - p = q \quad (21)$$

and

$$p' = k + \frac{q}{2}, \quad p = k - \frac{q}{2}. \quad (22)$$

Since we change two variables  $(p', p)$  to  $(k, q)$ , 4-integral variables becomes  $d^4 k d^4 q$ . Hence

$$\begin{aligned} \frac{1}{2} \frac{\delta^2}{\delta\phi^2} S[\phi_0] \tilde{\phi}^2 &= 2N_c \int \frac{d^4 k}{(2\pi)^4} \int \frac{d^4 q}{(2\pi)^4} \frac{4M^2 \tilde{\phi}(-q) \tilde{\phi}(q) - q^2 \tilde{\phi}(-q) \tilde{\phi}(q)}{\left\{ \left( k - \frac{q}{2} \right)^2 + M^2 \right\} \left\{ \left( k + \frac{q}{2} \right)^2 + M^2 \right\}} \\ &= 2N_c \int \frac{d^4 k}{(2\pi)^4} \int \frac{d^4 q}{(2\pi)^4} \frac{4M^2 - q^2}{\left\{ \left( k - \frac{q}{2} \right)^2 + M^2 \right\} \left\{ \left( k + \frac{q}{2} \right)^2 + M^2 \right\}} \tilde{\phi}(-q) \tilde{\phi}(q) \end{aligned} \quad (23)$$