

Permütasyonda sıralama veya diziliş söz konusudur. Seçilmiş nesnelerin sıralanışı veya dizilişi önemlidir.

Kombinasyonda ise, seçim veya seçme söz konusudur.

Discrete Mathematics

Counting

Ayşegül Gençata Yayımlı H. Turgut Uyar

2013-2016

sıra önemli değilse
daha az elemanlı küme

Permutations are for lists (order matters) and combinations are for groups (order doesn't matter).

1 / 39

License



© 2013-2016 A. Yayımlı, T. Uyar

You are free to:

- Share – copy and redistribute the material in any medium or format
- Adapt – remix, transform, and build upon the material

Under the following terms:

- Attribution – You must give appropriate credit, provide a link to the license, and indicate if changes were made.
- NonCommercial – You may not use the material for commercial purposes.
- ShareAlike – If you remix, transform, or build upon the material, you must distribute your contributions under the same license as the original.

For more information:

<https://creativecommons.org/licenses/by-nc-sa/4.0/>

Read the full license:

<https://creativecommons.org/licenses/by-nc-sa/4.0/legalcode>

2 / 39

Topics

Combinatorics

Introduction
Basic Principles

sıra önemsiz

çiğköfte yapma 5 kişilik takımdan 3 kişi seçme

Permutations

Introduction
Circular Arrangements

eldeki malzemelerle pasta yapma
5 kişilik takımda 3 kişi sıralama

Combinations

Introduction
With Repetition

3 / 39

Combinatorics

farklı nesne düzenleme (seçme) işidir ve bu süreçte
sıranın bu düzenleme de önemi yoktur.

seçme, düzenleme

- **combinatorics**: study of arrangement of objects
- **enumeration**: counting of objects with certain properties
- to solve a complicated problem:
- break it down into smaller problems
- piece together solutions to these smaller problems

sayma

divide and conquer

4 / 39

Sum Rule

- ▶ $task_1$ can be performed in n_1 distinct ways
- ▶ $task_2$ can be performed in n_2 distinct ways
- ▶ $task_1$ and $task_2$ cannot be performed simultaneously
- ▶ performing either $task_1$ or $task_2$ can be accomplished in $n_1 + n_2$ ways

5 / 39

Sum Rule Example

- ▶ a college library has 40 textbooks on sociology, and 50 textbooks on anthropology
- ▶ to learn about sociology or anthropology a student can choose from $40 + 50 = 90$ textbooks

seçmeli ders: 10 bilg, 5 elektronikten var. 1 dersi 15 farklı yolla

elbise 3 üst, 5 alt var. 1 tane elbise seçeceksin üzerindeki takımdan farklı olabilmesi için yani ya alt ya üst seçeceksen eğer kaç farklı seçimin olur
 $3+5$

6 / 39

Sum Rule Example

- ▶ a computer science instructor has two colleagues
- ▶ one colleague has 3 textbooks on "Introduction to Programming"
- ▶ the other colleague has 5 textbooks on the same subject
- ▶ n : maximum number of different books that can be borrowed
- ▶ $5 \leq n \leq 8$
- ▶ both colleagues may own copies of the same book

7 / 39

Product Rule

- ▶ a procedure can be broken down into $stage_1$ and $stage_2$
- ▶ n_1 possible outcomes for $stage_1$
- ▶ for each of these, n_2 possible outcomes for $stage_2$
- ▶ procedure can be carried out in $n_1 \cdot n_2$ ways

elbise 3 üst, 5 alt . kaç farklı üst-alt takımı oluşturabilirsin
 $3 \cdot 5$

8 / 39

Product Rule Example

- ▶ drama club is holding tryouts for a play
- ▶ 6 men and 8 women auditioning for the leading roles
- ▶ director can cast leading couple in $6 \cdot 8 = 48$ ways

çift seçecek yani 1 kadın 1 erkek

Product Rule Example

- ▶ license plates with 2 letters, followed by 4 digits
- ▶ how many possible plates?
- ▶ no letter or digit can be repeated:
 $26 \cdot 25 \cdot 10 \cdot 9 \cdot 8 \cdot 7 = 3,276,000$
- ▶ repetitions allowed:
 $26 \cdot 26 \cdot 10 \cdot 10 \cdot 10 \cdot 10 = 6,760,000$
- ▶ repetitions allowed, only vowels and even digits:
 $5 \cdot 5 \cdot 5 \cdot 5 \cdot 5 \cdot 5 = 15,625$

Product Rule Example

- ▶ a byte consists of 8 bits
- ▶ a bit has two possible values: 0 or 1
- ▶ number of possible values for a byte:
 $2 \cdot 2 \cdots 2 = 2^8 = 256$

Counting Example

- ▶ pastry shop menu:
6 kinds of muffins, 8 kinds of sandwiches
hot coffee, hot tea, iced tea, cola, orange juice
- ▶ buy either a muffin and a hot beverage,
or a sandwich and a cold beverage
- ▶ how many possible purchases?
- ▶ muffin and hot beverage: $6 \cdot 2 = 12$
- ▶ sandwich and cold beverage: $8 \cdot 3 = 24$
- ▶ total: $12 + 24 = 36$

Permutation

sıralama önemli

- ▶ **permutation**: a linear arrangement of distinct objects
- ▶ **order important** sıralama, dizilenmesi

13 / 39

Permutation Example

- ▶ a class has 10 students: A, B, C, \dots, I, J
- ▶ 4 students are to be seated in a row for a picture:
 $BCEF, CEFI, ABCF, \dots$
- ▶ how many such arrangements?
- ▶ filling of a position: a stage of the counting procedure
 $10 \cdot 9 \cdot 8 \cdot 7 = 5,040$

14 / 39

Permutation Example

$$\begin{aligned} 10 \cdot 9 \cdot 8 \cdot 7 &= 10 \cdot 9 \cdot 8 \cdot 7 \cdot \frac{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1} \\ &= \frac{10!}{6!} \end{aligned}$$

15 / 39

Permutations

n : total obje sayısı $r=10, r=4$
 r : sıralanacak obje

- ▶ n distinct objects
- ▶ number of permutations of size r (where $1 \leq r \leq n$):

$$\begin{aligned} P(n, r) &= n \cdot (n-1) \cdot (n-2) \cdots (n-r+1) \\ &= \frac{n!}{(n-r)!} \end{aligned}$$

- ▶ if repetitions are allowed: n^r

16 / 39

Permutations Example

- ▶ if size equals number of objects: $r = n$

$$P(n, n) = \frac{n!}{(n - n)!} = \frac{n!}{0!} = n!$$

example

- ▶ number of permutations of the letters in "COMPUTER":
8!

17 / 39

Arrangements Example

- ▶ number of arrangements of the letters in "BALL"
- ▶ two L's are indistinguishable

A	B	L	L	L	A	B	L
A	L	B	L	L	A	L	B
A	L	L	B	L	B	A	L
B	A	L	L	L	L	B	L
B	L	A	L	L	L	L	A
B	L	L	A	L	L	L	B

- ▶ number of arrangements: $\frac{4!}{2!} = 12$

18 / 39

Arrangements Example

- ▶ arrangements of all letters in "DATABASES" ayirt edilemeyen
- ▶ for each arrangement where A's are **indistinguishable**,
3! = 6 arrangements where A's are **distinguishable**:
 $DA_1TA_2BA_3SES, DA_1TA_3BA_2SES, DA_2TA_1BA_3SES,$
 $DA_2TA_3BA_1SES, DA_3TA_1BA_2SES, DA_3TA_2BA_1SES$
- ▶ for each of these, 2 arrangements where S's are distinguishable:
 $DA_1TA_2BA_3S_1ES_2, DA_1TA_2BA_3S_2ES_1$
- ▶ number of arrangements: $\frac{9!}{2! \cdot 3!} = 30,240$

19 / 39

Generalized Rule


genel kural

- ▶ n objects ayirt edilemeyen
- ▶ n_1 indistinguishable objects of $type_1$
 n_2 indistinguishable objects of $type_2$
...
- ▶ n_r indistinguishable objects of $type_r$
- ▶ $n_1 + n_2 + \dots + n_r = n$
- ▶ number of linear arrangements:

$$\frac{n!}{n_1! \cdot n_2! \cdot \dots \cdot n_r!}$$

20 / 39

Arrangements Example



- ▶ go from (2,1) to (7,4)
- ▶ each step one unit to the right (*R*) or one unit upwards (*U*)
- ▶ *RURRURRU*, *URRRUURR*
- ▶ how many such paths?

- ▶ each path consists of 5 *R*'s and 3 *U*'s
- ▶ number of paths: $\frac{8!}{5!3!} = 56$

asıl

21 / 39

Circular Arrangements Example

- ▶ 6 people seated around a round table: *A, B, C, D, E, F*
- ▶ arrangements considered to be the same when one can be obtained from the other by rotation:
ABEFCD, *DABEFC*, *CDABEF*, *FCDABE*, *EFCDAB*, *BEFCDA*
- ▶ how many different circular arrangements?
- ▶ each circular arrangement corresponds to 6 linear arrangements
- ▶ number of circular arrangements: $\frac{6!}{6} = 120$

(n-1)!

22 / 39

Combination Kombinasyon

sıra önemsiz seçme

- ▶ **combination**: choosing from distinct objects
- ▶ order not important

sıra önemsiz

elbise kombinasyonu

kot- tişört veya tişört-kot aynı kombinasyondur

23 / 39

Combination Example

permütasyon örneği

- ▶ a deck of 52 playing cards
- ▶ 4 suits: clubs, diamonds, hearts, spades
- ▶ 13 ranks in each suit: Ace, 2, 3, ..., 10, Jack, Queen, King
- ▶ draw 3 cards in succession, without replacement
- ▶ how many possible draws?

$$52 \cdot 51 \cdot 50 = \frac{52!}{49!} = P(52, 3) = 132,600$$

P(52,3)

24 / 39

Combination Example

kombinasyon örneği

kupa, karo, sinek, maça

örnek

- ▶ one such draw: **kupa as** **sinek** **karo papaz**
AH (ace of hearts), 9C (9 of clubs), KD (king of diamonds)
- ▶ if order doesn't matter
- ▶ 6 permutations of (AH, 9C, KD) correspond to just one selection

$$C(52, 3) = \frac{52!}{3! \cdot 49!} = 22,100$$

sıralama önemli değil,

52 kartlık bir desteden 3 kart çekiliyor:
Örneğin AH (kupa ası), 9C (sinek dokuz), KD (karo papazı).

Eğer kartların çekiliş sırası önemli değilse,
bu 3 kartın 6 farklı dizilişi (permütasyonu) aslında aynı seçime karşılık gelir.

Buna göre, 52 kartlık bir desteden sırası önemli olmadan 3 kart seçmenin kaç farklı yolu vardır?

25 / 39

Number of Combinations

- ▶ n distinct objects
- ▶ each combination of r objects: $r!$ permutations of size r
- ▶ number of combinations of size r (where $0 \leq r \leq n$):

$$C(n, r) = \binom{n}{r} = \frac{P(n, r)}{r!} = \frac{n!}{r! \cdot (n - r)!}$$

26 / 39

Number of Combinations

- ▶ number of combinations:

$$C(n, r) = \frac{n!}{r! \cdot (n - r)!}$$

- ▶ note that:

$$\begin{aligned} C(n, 0) &= 1 = C(n, n) \\ C(n, 1) &= n = C(n, n - 1) \end{aligned}$$

27 / 39

Number of Combinations Example

- ▶ Lynn and Patti buy a powerball ticket
- ▶ match five numbers selected from 1 to 49
- ▶ and then match powerball, 1 to 42
- ▶ how many possible tickets?
- ▶ Lynn selects five numbers from 1 to 49: $C(49, 5)$
- ▶ Patti selects the powerball from 1 to 42: $C(42, 1)$
- ▶ possible tickets: $\binom{49}{5} \binom{42}{1} = 80,089,128$

5 ana sayı: 1 ile 69 arasında seçilir. 1 Powerball (kırmızı top): 1 ile 26 arasında seçilir.

28 / 39

Number of Combinations Examples

C(53,9)

- ▶ for a volleyball team, gym teacher must select nine girls from junior and senior classes
- ▶ 28 junior and 25 senior candidates
- ▶ how many different ways?
- ▶ if no restrictions: $\binom{53}{9} = 4,431,613,550$
- ▶ if two juniors and one senior are best spikers and must be on the team: $\binom{50}{6} = 15,890,700$
- ▶ if there has to be four juniors and five seniors: $\binom{28}{4}\binom{25}{5} = 1,087,836,750$

$$C(28,4) * C(25,5)$$

$$C(53,9) = 53! / (44! * 9!)$$

$$(50 * 49 * 48 * 47 * 46 * 45) / 6!$$

$$C(50,6)$$

29 / 39

Binomial Theorem

Theorem

if x and y are variables and n is a positive integer, then:

$$\begin{aligned} (x + y)^n &= \binom{n}{0} x^0 y^n + \binom{n}{1} x^1 y^{n-1} + \binom{n}{2} x^2 y^{n-2} + \dots \\ &\quad + \binom{n}{n-1} x^{n-1} y^1 + \binom{n}{n} x^n y^0 \\ &= \sum_{k=0}^n \binom{n}{k} x^k y^{n-k} \end{aligned}$$

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

- ▶ $\binom{n}{k}$: binomial coefficient

30 / 39

Binomial Theorem Examples

- ▶ in the expansion of $(x + y)^7$, coefficient of $x^5 y^2$:
 $\binom{7}{5} = \binom{7}{2} = 21$

$$C(7,2)$$

31 / 39

Multinomial Theorem

Theorem

For positive integers n, t , the coefficient of $x_1^{n_1} x_2^{n_2} x_3^{n_3} \dots x_t^{n_t}$ in the expansion of $(x_1 + x_2 + x_3 + \dots + x_t)^n$ is

$$\frac{n!}{n_1! \cdot n_2! \cdot n_3! \cdot \dots \cdot n_t!}$$

where each n_i is an integer with $0 \leq n_i \leq n$, for all $1 \leq i \leq t$, and $n_1 + n_2 + n_3 + \dots + n_t = n$.

32 / 39

Multinomial Theorem Examples

- ▶ in the expansion of $(x + y + z)^7$, coefficient of $x^2y^2z^3$:

$$\binom{7}{2, 2, 3} = \frac{7!}{2! \cdot 2! \cdot 3!} = 210$$

33 / 39

Combinations with Repetition Example

Tekrarlı Kombinasyon

- ▶ 7 students visit a restaurant
- ▶ each of them orders one of the following:
cheeseburger (c), hot dog (h), taco (t), fish sandwich (f)
- ▶ how many different purchases are possible?

Kaç farklı satın alma işlemi yapılabilir?

34 / 39

Combinations with Repetition Example

$n=4, r=7$ r arrangement'lı yapılacak nesne

c	c	h	h	t	t	f	x	x		x	x		x	x		x
c	c	c	c	h	t	f	x	x	x	x		x		x		x
c	c	c	c	c	c	f	x	x	x	x	x	x				x
h	t	t	f	f	f	f		x		x	x		x	x	x	x
t	t	t	t	t	t	t			x	x	x	x	x	x	x	
f	f	f	f	f	f	f				x	x	x	x	x	x	x

- ▶ enumerate all arrangements of 10 symbols consisting of seven x's and three |'s
- ▶ number of different purchases: $\frac{10!}{7! \cdot 3!} = \binom{10}{7} = 120$

$C(7+4-1, 7)$

ayıraç sayısı mı?

3 tanesini söyleyince son kalana gerek var mı?

35 / 39

Number of Combinations with Repetition

4 yemek içinden 7 kişiye sipariş

- ▶ select, with repetition, r of n distinct objects
- ▶ considering all arrangements of r x's and $n - 1$ |'s

$$\frac{(n + r - 1)!}{r! \cdot (n - 1)!} = \binom{n + r - 1}{r}$$

7 tane yemek dağıtılıyor
 $r=7$

7 kişiye farklı şekilde 4 yemek çeşidini dağıt
 $r=7$
 $n=4$

36 / 39

Number of Combinations with Repetition Example

3 tane muz dağıtılıyor, $r=3$

- ▶ distribute 7 bananas and 6 oranges among 4 children
- ▶ each child receives at least one banana
- ▶ how many ways?

- ▶ step 1: give each child a banana
- ▶ step 2: distribute 3 bananas to 4 children

1	1	1	0	b		b		b	
1	0	2	0	b			b	b	
0	0	1	2			b		b	b
0	0	0	3				b	b	b

4 çocuk

- ▶ $C(6, 3) = 20$ ways

$n=4, r=3$

3 muzu farklı şekilde

$r=3, n=4$

3 muz, 4 çocuğa. distinct object muz

37 / 39

Number of Combinations with Repetition Example

- ▶ step 3: distribute 6 oranges to 4 children

1	2	2	1	o		o	o		o	o		o
1	2	0	3	o		o	o			o	o	o
0	3	3	0		o	o	o		o	o	o	
0	0	0	6				o	o	o	o	o	o

- ▶ $C(9, 6) = 84$ ways

- ▶ step 4: by the rule of product: $20 \cdot 84 = 1,680$ ways

38 / 39

References

Required Reading: Grimaldi

- ▶ Chapter 1: Fundamental Principles of Counting
 - ▶ 1.1. The Rules of Sum and Product
 - ▶ 1.2. Permutations
 - ▶ 1.3. Combinations
 - ▶ 1.4. Combinations with Repetition

39 / 39