

Cosmology Tutorial 1

Howard Kinsman

February 7, 2017

1

$$f = \frac{L}{4\pi r^2} \quad (1)$$

and differentiating

$$\frac{df}{dr} = -\frac{L}{2\pi r^3} \quad (2)$$

Number density = $\frac{1}{r}$ so

$$N = \frac{4}{3}\pi r^3 \times \frac{1}{r} \quad (3)$$

$$N = \frac{4\pi r^2}{3} \quad (4)$$

and differentiating

$$\frac{dN}{dr} = \frac{8\pi r}{L} \quad (5)$$

dividing equation 5 by equation 2 gives

$$\frac{dN}{df} = \frac{-48\pi^2 r^4}{L} \quad (6)$$

so that

$$\frac{dN}{df} \propto r^4 \quad (7)$$

$r^2 \propto f^{-1}$ so that

$$\frac{dN}{df} \propto f^{-2} \quad (8)$$

Therefore $\alpha = -2$. $d \ln f = \frac{1}{f} df$ so that

$$\mathcal{N} = \frac{dN}{d \ln f} = f \frac{dN}{df} \quad (9)$$

which gives $\mathcal{N} \propto f^{-1}$ and so

$$\log \mathcal{N} = -\log f + \text{const} \quad (10)$$

This is linear so measuring stellar fluxes would allow an observer to determine if the distribution was homogeneous.

2

The NRAO VLA Sky Survey (NVSS) is a radio survey covering the sky north of -40 declination at 1.4GHz. It contains over a million rows so I extracted a subset of sources where the flux density $> 100mJy$ giving approximately 62,000 sources.

The Two Micron All Sky Survey (2MASS) Redshift Survey (2MRS) was a ten-year project to map the full three-dimensional distribution of galaxies in the nearby universe. It provides a good census of baryonic matter in the nearby universe (to about 300 Mpc). It contains about 400,000 rows so I only extracted sources with a magnitude < 13 in the K band giving about 44,000 rows.

The 2MASS Extended Source Catalog (2MASX) contains over 1.6 millions rows and suveys the entire sky in the J, H and K bands. I restricted the output to sources with a magnitude of < 13 in the J band giving about 77,000 rows.

3

Figure 1 shows a log of number counts per log flux for the NVSS. The slope was calculated:

$$\frac{\log(2000/1000)}{\log(100 - 200)} = -1 \quad (11)$$

This figure matches quite well with the figure of -1.5 calculated for a homogeneous universe. The slope is constant apart from very small flux densities (probably due to sampling errors) and large flux densities (the universe is not homogeneous on smaller scales).




Figure 1: NVSS

Figure 2 shows a log of number counts per magnitude in the K band for the 2MRS survey.




Figure 2: 2MRS

The slope was calculated as:

$$\frac{\log(1000/100)}{10/7} = .8 \quad (12)$$

This also reasonably matches the expected slope of .6 for a homogeneous universe. Finally figure 3 shows a log of number counts per magnitude in the J band for the 2MASX survey.




Figure 3: 2MASX

The slope was calculated as:

$$\frac{\log(10000/1000)}{12/11} = .9 \quad (13)$$

This is a similar figure to the 2MRS survey.

4

Figure 4 shows the 2MRS data in galactic coordinates restricted to the the closest half of the galaxies i.e. $cz < 8000$ km/s. It clearly shows evidence of structure indicating that the universe is not homogeneous on smaller scales.




Figure 4: 2MRS