

MSc Project - Modelling a Double-Barred Galaxy as a Double Binary

Howard Kinsman

August 14, 2017

Abstract

TODO

1 Introduction

1.1 Double-Barred Galaxies

Double-barred galaxies, where a smaller inner bar is nested inside a larger outer bar, were first observed in 1975 by de Vaucouleurs (de Vaucouleurs, 1975), however it was only in the 1990s that double-barred galaxies were recognised as being a distinct category of galaxies. The current estimates of double-barred galaxy frequencies are $\approx 30\%$ of barred galaxies and $\approx 20\%$ of all galaxies (Erwin, 2003). Erwin (2011) surveyed 38 barred galaxies and found a total of 10 double-barred galaxies with no preference for Hubble type. They found that inner bars were about 12% the size of the outer bars with semi-major axes from $\approx 250\text{pc}$ to 1 kpc. Secondary bars probably rotate independently of, but in the same direction as their associated primary bars Erwin (2011). Erwin and Sparke (2002) believe that inner bars probably rotate independently of their outer bars and suggest that they are relatively long-lived structures. Inner bar size has been found to correlate slightly with outer bar size (Erwin, 2003). The median size ratio is ≈ 0.12 with an upper limit of ≈ 0.25 which is consistent with the theory that inner bars cannot be too large without disrupting the orbit of the outer bar.




Figure 1: This is a schematic drawing of the various components of a double-barred galaxy. Credit: ESO

It has been suggested that nested bars may provide a mechanism for fueling active galactic nuclei (Shlosman et. al., 1989) however more recently (Erwin, 2003) believe that inner bars only play a minor role in this activity. (Erwin and Sparke, 2002) compared single and double-barred galaxies and found that there was no significant difference in nuclear activity. To investigate this theory further Lorenzo-Caceres et. al. (2002) carried out a study of NGC 357. This study aimed to present the first detailed morphological, kinematical and stellar population analysis of the bulge, inner and outer bars of a double-barred galaxy. They concluded that the bulge and inner bar show similar stellar population properties whilst the outer bar was less metal-rich. This result implied that, for the case of NGC 357, the outer bar was formed in a shorter timescale than the inner structures and so disagrees with traditional idea that gas flown along the outer structure triggers star formation causing the formation of the inner structure. This was strengthened by a further study of four double-barred galaxies by Lorenzo-Caceres et. al. (2013)

where they concluded that the stellar populations of inner bars are younger and more metal-rich than the outer bars. This suggests that at present inner bars play a moderate or even minor role in the morphological evolution of double-barred galaxies (Lorenzo-Caceres et. al., 2013).

Through the use of computer modeeling Saha and Maciejewski (2013) showed that double bars can form without gas in a dark matter dominated halo. They produced a model where the inner bar was rotating at almost as slowly as the outer bar. The route to the formation of double bars maybe very different to that of a single bar.

1.2 N-Body Simulations and the Four-Body Problem

Studies of three or more bodies interacting gravitationally dates back to the time of Newton. However it is only with the widespread use of computers that orbits of more than three bodies have been extensively studied. Today N-body simulations are an essential tool in the study of solar system dynamics and galactic dynamics (Binney and Tremaine, 2008). Celestial mechanics during the preceding centuries has mainly focussed on the three body problem due to the complexities involved in adding more bodies. It is estimated that within the Milky Way approximately two thirds of all stars exist in binary systems and a further one fifth of these are in triple systems whilst a further fifth still exist in systems with more than three bodies (Steves and Roy, 1998). Steves and Roy (1998) estimate that $\approx 2 \times 10^9$ quadruple systems exist in the Milky Way galaxy. They found that quadruple systems exist in two forms: a double binary or linear.

The classical equations of motion for an N-body problem are:

$$m_i \ddot{\mathbf{r}} = \sum_{i \neq j} \frac{m_i m_j}{r_{ij}^3} \mathbf{r}_{ij} \quad i = 1, 2, 3, \dots \quad (1)$$

where $\mathbf{r}_i = (x_i, y_i)$ and $\mathbf{r}_{ij} = \mathbf{r}_j - \mathbf{r}_i$. The units have been selected so that the gravitational constant (G) is equal to one. In simulations it is natural to choose units where $G = 1$ and the units of length and mass are conventionally chosen so that the total mass of the system is $M = 1$ (Heggie and Hut, 2003).

The four body problem, specifically, has been studied extensively by Steves and Roy (1998). Here they model four equal mass bodies orbiting in circular coplanar orbits. This model they term the Caledonian problem, because their research is being carried out at Glasgow Caledonian University. Approximately a century ago the Copenhagen problem was studied extensively in Copenhagen which consisted of a three body problem: two equal mass bodies and an infinitesimally small test particle. Szell et. al. (2004) also

studied the Caledonian problem and found that the stability of the system is dependent upon a constant termed the Szebehely constant. The Caledonian Symmetric Double Binary Problem (CSDBP) is relevant in studying the stability and evolution of symmetric quadruple stellar clusters and exoplanetary systems of two planets orbiting a binary pair of stars (AlvarezRamirez and Medina, 2014).

Methods for the study of N-body simulations have changed considerably since the advent of computers. In a study of N-body simulations Trenti and Hut (2008) mentions a pioneering attempt by Holmberg in 1941 which used light bulbs and galvanometers to track the evolution of 37 particles. However it wasn't until the early seventies and the advent of computers that this field really got started. Nowadays simulations can up to a million particles (Heggie and Hut, 2003) which is of particular relevance to the study of globular clusters. I refer the reader to Aarseth (2003) for much more detailed explanation of the field of N-body simulations.

1.3 Orbits, Loops, Resonances and Double-Barred Galaxy Simulations

Double-barred galaxy simulations and models have taken many forms. Du et. al. (2015) explored the formation and evolution of double-barred galaxies (which is not very well understood) by showing that a dynamically cool inner disk embedded in a hotter outer disk can generate a secondary bar while the outer disk forms a large-scale primary bar. Maciejewski et. al. (2001) used hydrodynamical simulations to study gas inflow in barred galaxies and the effect this has on secondary bars. They concluded that the secondary bar can prevent, rather than enhance, the inflow. More recently, Debattista and Shen (2007) used collisionless N-body simulations to show that inner bars can form spontaneously without requiring gas. They also found that secondary bars rotate faster than primary bars and that the secondary bars pulsate. This is considered further in the Discussion.

In an attempt to explain the dynamics of the orbits of nested bars Maciejewski and Sparke (1999) introduced the concept of ‘loops’. They wanted to investigate how two independently rotating bars can maintain themselves as gravitating systems. A gravitational potential consisting of two concentric bisymmetric bars that are independently rotating is not stationary in any reference frame Maciejewski and Sparke (1999). In the potential of a single bar particles would stay on a stable closed periodic orbit but with nested bars orbits will not be closed because the particles would be experience two separate forces. The ‘loop’ therefore was introduced as an extension of the

concept of orbit. Particles in a ‘loop’ would return to their original positions every time the bars come back to the same relative orientation (Maciejewski and Sparke, 1999). This lead to the conclusion that loops supporting the inner bar are thicker with parallel bars and thinner with perpendicular bars (Maciejewski, 2008). This also suggests that the two bars do not rotate through each other as rigid bodies but the inner bar should pulsate and that the bars should spend more time nearly orthogonal (Maciejewski, 2008). These predictions were confirmed by (Debattista and Shen, 2007) and (Shen and Debattista, 2008). This was further confirmed by Maciejewski and Small (2010) who found that the inner bar should end well within its corotation radius and that the inner bar extends further out for lower pattern speeds than higher ones; the inner bar pulsates; and that the angular velocity of the inner bar is not uniform and that faster inner bars rotate more coherently.

Wozniak (2015) also carried out double-barred galaxy simulations and achieved long-lived inner bars lasting for up to 7 Gyr. These were N-body/hydrodynamical simulations. They found that the two bar lengths, ratio of pattern speeds, as well as the age of the inner stellar bar populations in the simulations matched well with observations. They concluded that because, unlike previous simulations, there was no overlap between primary bar inner Lindblad resonance (ILR) and the inner bar corotation (CR) or any kind of resonance overlap, then some other kind of mechanism must feed the central waves (Wozniak, 2015). It is suggested that star formation may be responsible for bringing energy into the inner bar.

Garzon (2014) studied the dynamical evolution of two isolated bars, within the same galaxy, under their mutual gravitational interaction. They considered two cases purely analytically: rotation of rigid bodies and rotation of deformable rotation. They found that the bars oscillate in size according to the relative angle between the bars. This is broadly in agreement with (Debattista and Shen, 2007). What follows is an attempt to verify the calculations in (Garzon, 2014) but using a simpler model of a double-barred galaxy using a double binary.

2 Methods

Fortran 90 was selected as the language of choice in order to avoid the fixed line limitations of Fortran 77. The model was created using OdeInt from Numerical Recipes. I first converted OdeInt to Fortran 90 for this purpose. Please see the Compilers subsection for a few notes on compilers - may or may not be relevant to the project. So for a four body problem this equation

results in the following:

$$\ddot{\mathbf{r}}_1 = \frac{m_2 \mathbf{r}_{12}}{r_{12}^3} + \frac{m_3 \mathbf{r}_{13}}{r_{13}^3} + \frac{m_4 \mathbf{r}_{14}}{r_{14}^3} \quad (2)$$

$$\ddot{\mathbf{r}}_2 = \frac{m_1 \mathbf{r}_{21}}{r_{21}^3} + \frac{m_3 \mathbf{r}_{23}}{r_{23}^3} + \frac{m_4 \mathbf{r}_{24}}{r_{24}^3} \quad (3)$$

$$\ddot{\mathbf{r}}_3 = \frac{m_1 \mathbf{r}_{31}}{r_{31}^3} + \frac{m_2 \mathbf{r}_{32}}{r_{32}^3} + \frac{m_4 \mathbf{r}_{34}}{r_{34}^3} \quad (4)$$

$$\ddot{\mathbf{r}}_4 = \frac{m_1 \mathbf{r}_{41}}{r_{41}^3} + \frac{m_2 \mathbf{r}_{42}}{r_{42}^3} + \frac{m_3 \mathbf{r}_{43}}{r_{43}^3} \quad (5)$$

The above equations were encoded into Fortran. A full code listing is supplied in a separate file.

2.1 Initial Conditions

As a starting point both the outer binary and inner binary were placed in circular orbits and tested separately and then combined to ensure a stable starting point. The inner binary had initially a negligible mass. The outer binary initial conditions were:

$m1=.5$, $m2=.5$, $x1=-.5$, $x2=.5$, $y1=0$, $y2=0$,
 $vx1=0$, $vx2=0$, $vy1=-.5$, $vy2=.5$

and for the inner binary:

$m3=.001$, $m4=.001$, $x3=-.001$, $x4=.001$, $y3=0$, $y4=0$,
 $vx3=0$, $vx4=0$, $vy3=-.5$, $vy4=.5$

Plots of the two initial binaries are shown in Fig. 2 and 3. The scale of the inner binary is so small that it is difficult to see in GnuPlot.




Figure 2: Outer Binary




Figure 3: Inner Binary

I then proceeded to perturb the system by gradually increasing the mass of the inner binary. As the system became unstable I compensated for the

increasing mass of the inner binary by increasing the velocity of the outer binary and also decreasing the velocity and increasing the binary separation of the inner binary. In many of the configurations the systems were inherently unstable and produced collisions with some or all of the bodies being ejected from the system. These were rejected and only 'stable' configurations were considered. Table 1 shows the perturbations made to the system - only the variables in this table changed and all the other initial conditions remained the same.

2.2 Compilers


As I had converted OdeInt into Fortran 90 I wanted to make sure I hadn't introduced any errors. I therefore compared the output of the Numerical Recipes version of OdeInt with mine. I was compiling with GFortran and the Fortran 77 version was compiled with g77. They were different! So naturally concerned I then compiled the Fortran 77 version with GFortran. The results were identical to mine, which satisfied me that I hadn't introduced any errors.

3 Results

As can be seen from Fig. 4 even a negligible mass inner binary causes the outer binary to begin to separate and become less tightly bound. Figures Fig. 4 to 39 show plots of the double binary using the parameters given in Table 1. In many cases the inner binary is only visible as a point at the centre of the outer binary. As the mass of the inner binary increased the system became increasingly unstable and the effect on the outer binary became more pronounced. Even the 'stable' configurations listed in Table 1 hold for only a few orbits. I was unable to recreate a stable system for a mass $> .08$ for the inner bar. The plots below reveal that increasing the mass of the inner binary causes the outer binary to separate and form wider orbits.

Table 1: Configurations

Config.	m3	m4	vy1	vy2	x3	x4	vy3	vy4
1	.001	.001	-.5	.5	-.001	.001	-.5	.5
2	.002	.002	-.5	.5	-.001	.001	-.5	.5
3	.003	.003	-.5	.5	-.001	.001	-.5	.5
4	.004	.004	-.5	.5	-.004	.004	-.5	.5
5	.004	.004	-.55	.55	-.004	.004	-.5	.5
6	.004	.004	-.57	.57	-.004	.004	-.5	.5
7	.005	.005	-.58	.58	-.005	.005	-.3	.3
8	.005	.005	-.58	.58	-.005	.005	-.4	.4
9	.005	.005	-.58	.58	-.005	.005	-.35	.35
10	.006	.006	-.6	.6	-.006	.006	-.3	.3
11	.025	.025	-.65	.65	-.02	.02	-.5	.5
12	.025	.025	-.7	.7	-.025	.025	-.5	.5
13	.03	.03	-.7	.7	-.03	.03	-.5	.5
14	.035	.035	-.75	.75	-.04	.04	-.4	.4
15	.04	.04	-.75	.75	-.045	.045	-.4	.4
16	.05	.05	-.75	.75	-.045	.045	-.4	.4
17	.05	.05	-.75	.75	-.05	.05	-.35	.35
18	.05	.05	-.75	.75	-.05	.05	-.3	.3
19	.06	.06	-.8	.8	-.06	.06	-.25	.25
20	.08	.08	-.9	.9	-.08	.08	-.15	.15
21	.09	.09	-.95	.95	-.09	.09	-.1	.1
22	.1	.1	-1	1	-.1	.1	-.05	.05
23	.1	.1	-1	1	-.1	.1	-.04	.04
24	.1	.1	-1	1	-.1	.1	-.03	.03
25	.1	.1	-1	1	-.1	.1	-.02	.02
26	.12	.12	-1.05	1.05	-.11	.11	-.015	.015
27	.12	.12	-1.1	1.1	-.11	.11	-.015	.015
28	.12	.12	-1.1	1.1	-.115	.115	-.015	.015
29	.12	.12	-1.1	1.1	-.12	.12	-.015	.015
30	.1	.1	-1	1	-.1	.1	-.1	.1
31	.1	.1	-1	1	-.102	.102	-.1	.1
31	.1	.1	-1	1	-.1021	.1021	-.1	.1






Figure 5: Configuration 1 - Inner Bar




Figure 6: Configuration 2




Figure 7: Configuration 2 - Inner Bar




Figure 8: Configuration 3




Figure 9: Configuration 3 - Inner Bar




Figure 10: Configuration 4




Figure 11: Configuration 4 - Inner Bar

1




Figure 12: Configuration 5




Figure 13: Configuration 5 - Inner Bar




Figure 14: Configuration 6




Figure 15: Configuration 6 - Inner Bar




Figure 16: Configuration 7




Figure 17: Configuration 7 - Inner Bar




Figure 18: Configuration 8




Figure 19: Configuration 8 - Inner Bar




Figure 20: Configuration 9




Figure 21: Configuration 9 - Inner Bar




Figure 22: Configuration 10




Figure 23: Configuration 10 - Inner Bar




Figure 24: Configuration 11




Figure 25: Configuration 11 - Inner Bar




Figure 26: Configuration 12




Figure 27: Configuration 12 - Inner Bar




Figure 28: Configuration 13




Figure 29: Configuration 13 - Inner Bar




Figure 30: Configuration 14




Figure 31: Configuration 14 - Inner Bar




Figure 32: Configuration 15




Figure 33: Configuration 15 - Inner Bar




Figure 34: Configuration 16




Figure 35: Configuration 16 - Inner Bar




Figure 36: Configuration 17




Figure 37: Configuration 17 - Inner Bar




Figure 38: Configuration 18




Figure 39: Configuration 18 - Inner Bar




Figure 40: Configuration 19




Figure 41: Configuration 19 - Inner Bar




Figure 42: Configuration 20




Figure 43: Configuration 20 - Inner Bar




Figure 44: Configuration 21




Figure 45: Configuration 21 - Inner Bar




Figure 46: Configuration 22




Figure 47: Configuration 22 - Inner Bar




Figure 48: Configuration 23




Figure 49: Configuration 23 - Inner Bar




Figure 50: Configuration 24




Figure 51: Configuration 24 - Inner Bar




Figure 52: Configuration 25




Figure 53: Configuration 25 - Inner Bar




Figure 54: Configuration 26




Figure 55: Configuration 26 - Inner Bar




Figure 56: Configuration 27




Figure 57: Configuration 27 - Inner Bar




Figure 58: Configuration 28




Figure 59: Configuration 28 - Inner Bar




Figure 60: Configuration 29




Figure 61: Configuration 29 - Inner Bar




Figure 62: Configuration 30




Figure 63: Configuration 30 - Inner Bar




Figure 64: Configuration 31




Figure 65: Configuration 31 - Inner Bar




Figure 66: Configuration 32




Figure 67: Configuration 32 - Inner Bar

4 Discussion

The final three runs (configurations 30, 31, and 32) were by far the most stable. The orbital periods vary slightly with time so it is difficult to give accurate figures, so I am using the periods of the first orbit. The following table shows the periods of the two binaries together with the ratios of the corresponding periods. The data shows that as the periodic ratios tends

Table 2: Periods

Config.	Inner	Outer	Ratio	Length of simulation
30	.51	11.56	1:22.7	25
31	.54	11.5	1:21.3	25
32	.55	11.5	1:20.9	35

towards 1:21 the system considerably stabilises. Run 32 was by far the most stable run and the orbits persisted until the integrator errored with a 'Step size underflow'. 1:21 could possibly be an orbital resonance. Fourier analysis? Rewrite in Python, automate, introduce machine learning algorithm

(Garzon, 2014), (Erwin and Sparke, 2002), (?), (?), (Maciejewski, 2008), (?), (Maciejewski and Small, 2010) (Maciejewski and Sparke, 1998), (Maciejewski and Sparke, 1999), (Manos and Athanassoula, 2011), (Shen and Debattista, 2008), (Shen and Debattista, 2009), (Debattista and Shen, 2007), (Malhotra, 1998)

References

- Aarseth, S., Gravitational N-Body Simulations, 2003, Cambridge University Press
- AlvarezRamirez, M., Medina, M., A Review of the Planar Caledonian Four-Body Problem, 2014
- Binney, J., Tremaine, S., Galactic Dynamics, 2008, Princeton University Press
- de Vaucouleurs, G., 1975, ApJS, 29, 193
- Debattista, V., Shen, J., Long-Lived Double-Barred Galaxies from Pseudobulges, 2007, The Astrophysical Journal, 654: L127L130, 2007 January 10
- Du, M., Shen, J., Debattista, V., Forming Double-Barred Galaxies from Dynamically Cool Inner Disks, 2015, The Astrophysical Journal, 804:139 (10pp), 2015 May 10
- Erwin, P., Double-Barred Galaxies I. A Catalog of Barred Galaxies with Stellar Secondary Bars and Inner Disks, 2003, arXiv:astro-ph/0310806v2
- Erwin, P., Double-Barred Galaxies, 2011, Mem. S.A.It. Suppl. Vol. 18, 145
- Erwin, P., Sparke, L., Double Bars, Inner Disks, and Nuclear Rings in Early-Type Disk Galaxies, 2002, The Astronomical Journal, 124:6577, 2002 July
- Garzon, F., Lopez-Corredoira, M., Dynamical evolution of two associated galactic bars, 2014, arXiv:1409.1916.v1
- Heggie D., Hut, P., The Gravitational Million-Body Problem, 2003, Cambridge University Press
- de Lorenzo-Caceres, A., Vazdekis, A., Aguerri, K., A., L., Corsini, E., M., Debattista, V., P., Constraining the formation of inner bars. Photometry, kinematics and stellar populations in NGC357, 2002, arXiv:1111.1718v1
- de Lorenzo-Caceres, A., Falcon-Barroso, J., Vazdekis, A., Distinct stellar populations in the inner bars of double-barred galaxies, 2013, arXiv:1302.5701v1
- Maciejewski, W., Teuben, P., J., Sparke, L., S., Stone, J., M., Gas Inflow in Barred Galaxies - Effects of Secondary Bars, arXiv:astro-ph/0109431v1

- Maciejewski, W., Orbits in corotating and counterrotating double bars, 2008, arXiv:0801.1471v1
- Maciejewski, W., Small, E., Orbital Support of Fast and Slow Inner Bars in Double Barred Galaxies, 2010, arXiv:1006.4574v1
- Maciejewski, W., Sparke, L., Bars within Bars in Galaxies, 1998, arXiv:astro-ph/9812228v1
- Maciejewski, W., Sparke, L., Orbits Supporting Bars within Bars, 1999, arXiv:astro-ph/9911281v1
- Malhotra, R., Orbital Resonances and Chaos in the Solar System, 1998, Solar system Formation and Evolution ASP Conference Series, Vol. 149, 1998
- Manos, T., Athanassoula, E., Regular and chaotic orbits in barred galaxies - I. Applying the SALI/GALI method to explore their distribution in several models, 2011, arXiv:1102.1157v2
- Moiseev, A., V., Velocity dispersion of stars and gas motion in double-barred galaxies, 2001, arXiv:astro-ph/0111220v1
- Press, W., H., Teukolsky, S., A., Vetterling, W., T., Flannery, B., P., Numerical Recipes in Fortran 77 The Art of Scientific Computing, 2nd edition
- Saha, K., Maciejewski, W., Spontaneous formation of double bars in dark matter dominated galaxies, 2013, arXiv:1304.7108v1
- Sellwood, J., Wilkinson, A., Dynamics of Barred Galaxies, 2006, arXiv:astro-ph/0608665v1
- Shen, J., Debattista, V., Long-lived double-barred galaxies in N-body simulations, 2008, Mem. S.A.It. Vol. 00, 169
- Shen, J., Debattista, V., Observable Properties of Double-Barred Galaxies in N-Body Simulations, 2009, The Astrophysical Journal, 690:758772, 2009 January 1
- Shlosman, I., Frank, J., Begeman, M.,C., 1989, Nature, 338, 45
- Steves, B., Roy, A., Some special restricted four-body problems-I. Modelling the Caledonian problem, 1998, Phrt. Spucx, Si., Vol. 46. No. I I /) 12, pp. 1465-1474, 1998

Szell, A., Erdi, B., Sandor, Zs., Steves, B., Chaotic and stable behaviour in the Caledonian Symmetric Four-Body Problem, 2004, Mon. Not. R. Astron. Soc. 347, 380388 (2004)

Trenti, M., Hut, P., Gravitational N-body Simulations, 2008, arXiv:0806.3950v1

Wozniak, H., How can double-barred galaxies be long-lived?, 2015, A and A 575, A7 (2015)