FLU EXPRESS



Flu Express is a weekly report produced by the Respiratory Disease Office of the Centre for Health Protection. It monitors and summarizes the latest local and global influenza activities.

Local Situation of Influenza Activity (as of Oct 2, 2013)

Reporting period: Sep 22 - 28, 2013 (Week 39)

- The latest surveillance data showed that the local influenza activity remained elevated.
- The Centre for Health Protection has collaborated with the Hospital Authority and private hospitals to monitor influenza associated intensive care unit (ICU) admissions or deaths (aged 18 years or above) since Jul 26, 2013. As of Oct 2, there were 46 cases of influenza associated ICU admission or death, in which 29 of them were fatal.
- Influenza can cause serious illnesses in high-risk individuals and even healthy persons. Except for those with contraindications, influenza vaccination is suitable for all members of the public.
- Children (aged between six months and less than 6 years, or attending a kindergarten or child care centre in Hong Kong) and elderly (aged 65 years or above), who are eligible, will be subsidised for seasonal influenza vaccination from enrolled private doctors participating in the Government's vaccination subsidy schemes starting from Oct 2, 2013.

Influenza-like-illness surveillance among sentinel general outpatient clinics, 2011-13

In week 39, the average consultation rate for influenza-like illness (ILI) among sentinel general outpatient clinics (GOPCs) was 3.0 ILI cases per 1,000 consultations (Figure 1).




Figure 1 ILI consultation rate at sentinel GOPC, 2011-13

Influenza-like illness surveillance among sentinel private doctors, 2011-13

In week 39, the average consultation rate for ILI among sentinel private doctors was 29.7 ILI cases per 1,000 consultations (Figure 2).




Figure 2 ILI consultation rate at sentinel GP, 2011-13

Influenza virus detections (Laboratory surveillance), 2012-13

In week 39, the number of influenza viruses detected in the Public Health Laboratory Centre was 345, including 297 influenza A (H3) viruses, 28 influenza B viruses and 20 influenza A (H1N1)pdm09 viruses (Figure 3).




Figure 3 Influenza virus detections, 2012-13

Influenza-like illness outbreak monitoring, 2012-13

In week 39, 5 ILI outbreaks occurring in schools/institutions were recorded. In the first 4 days of week 40 (Sep 29 to Oct 2, 2013), 3 ILI outbreaks occurring in schools/institutions were recorded (Figure 4).




Figure 4 ILI outbreaks, 2012-13

Rate of influenza-like illness syndrome group in accident and emergency departments, 2011-13#

In week 39, the rate of the influenza-like illness syndrome group in the accident and emergency departments (AED) was 162.5 (per 1,000 coded cases) (Figure 5).

#Note: The influenza-like illness syndrome group includes codes such as influenza, upper respiratory tract infection, fever, cough, throat pain, and nneumonia




Figure 5 Rate of ILI syndrome group in AED, 2011-13

Influenza associated hospital admission rates and deaths based on discharge coding, 2011-13

In week 39, hospital admission rates with principal diagnosis of influenza for persons aged 0-4 years, 5-64 years and 65 years or above were 1.35, 0.05 and 0.27 cases respectively (per 10,000 people in the age group) (Figure 6). Weekly number of deaths with any diagnosis of influenza in public hospitals is also shown in the figure on the right hand side.




Figure 6 Influenza associated hospital admission rates and deaths, 2011-13

Fever surveillance at sentinel child care centres/kindergartens, 2011-13

In week 39, 0.83% of children in the sentinel child care centres / kindergartens (CCC/ KG) had fever (Figure 7).




Figure 7 Percentage of children with fever at sentinel CCC/KG, 2011-13

Fever surveillance at sentinel residential care homes for the elderly, 2011-13

In week 39, 0.12% of residents in the sentinel residential care homes for the elderly (RCHEs) had fever (Figure 8).




Figure 8 Percentage of residents with fever at sentinel RCHE, 2011-13

Influenza-like illness surveillance among sentinel Chinese medicine practitioners, 2011-13

In week 39, the average consultation rate for ILI among Chinese medicine practitioners (CMPs) was 2.10 ILI cases per 1,000 consultations (Figure 9).




Figure 9 ILI consultation rate at sentinel CMP, 2011-13

Enhanced Surveillance for Severe Seasonal Influenza (Aged 18 years or above)

• From Sep 25, 2013, 12:00 noon to Oct 2, 2013, 12:00 noon, 3 cases of influenza associated ICU admission were recorded, in which 2 of them were fatal.

Note: The data reported are provisional figures and subject to further revision.

Surveillance of severe paediatric influenza-associated complication/death (Aged above one month and below 18 years)

• In week 39 and the first 4 days of week 40 (Sep 29 to Oct 2, 2013), there were no new reports of severe paediatric influenza-associated complication/death.

Note: The data reported are provisional figures and subject to further revision.

Surveillance of Tamiflu resistant influenza A(H1N1)2009 virus infection

 In week 39 and the first 4 days of week 40 (Sep 29 to Oct 2, 2013), there were no new reports of oseltamivir (Tamiflu) resistant influenza A(H1N1)pdm09 virus infection. There are totally 36 reports of Tamiflu resistant influenza A(H1N1)pdm09 virus detected in Hong Kong since 2009.

Global Situation of Influenza Activity

- In the United States (week ending Sep 21, 2013), the influenza activity remained low.
- In Canada (week ending Sep 21, 2013), the influenza activity remained low.
- In the United Kingdom (week ending Sep 22, 2013), the ILI consultation rate remained low.
- In Europe (week ending Sep 22, 2013), the ILI consultation rate were at low levels in all countries in the Region.
- In New Zealand (week ending Sep 22, 2013), the ILI consultation rate was below the baseline.
- In Australia (week ending Sep 13, 2013), nationally the 2013 influenza season appeared to have peaked. Overall influenza activity had been relatively low compared to 2011 and 2012.

Sources:

Information have been extracted from the following sources when updates are available: <u>United States Centers for Disease Control and Prevention</u>, <u>Public Health Agency of Canada</u>, <u>United Kingdom Health Protection Agency</u>, <u>World Health Organization/Europe Euroflu</u>, New Zealand Ministry of Health and Australia Department of Health and Ageing.