

FLU EXPRESS

Flu Express is a weekly report produced by the Respiratory Disease Office of the Centre for Health Protection. It monitors and summarizes the latest local and global influenza activities.

Local Situation of Influenza Activity (as of May 8, 2013)

Reporting period: Apr 28 – May 4, 2013 (Week 18)

- The latest surveillance data showed that the local influenza activity is decreasing as compared with past few weeks.
- The Centre for Health Protection has collaborated with the Hospital Authority and private hospitals to monitor influenza associated intensive care unit (ICU) admissions or deaths (aged 18 years or above) since Jan 18, 2013. As of May 8, there were 68 cases of influenza associated ICU admission or death, of which 28 of them were fatal.
- Influenza can cause serious illnesses in high-risk individuals and even healthy persons. Except for those with contraindications, influenza vaccination is suitable for all members of the public.
- Children between the age of six months and less than 6 years old as well as elderly people aged 65 years and above are eligible to receive subsidised vaccination from enrolled private doctors participating in the Government's vaccination subsidy schemes starting from Sep 24, 2012.

Influenza-like-illness surveillance among sentinel general outpatient clinics, 2011-13

In week 18, the average consultation rate for influenza-like illness (ILI) among sentinel general outpatient clinics (GOPCs) was 3.3 ILI cases per 1,000 consultations (Figure 1).




Figure 1 ILI consultation rate at sentinel GOPC, 2011-13

Influenza-like illness surveillance among sentinel private doctors, 2011-13

In week 18, the average consultation rate for ILI among sentinel private doctors was 32.8 ILI cases per 1,000 consultations (Figure 2).




Figure 2 ILI consultation rate at sentinel GP, 2011-13

Influenza virus detections (Laboratory surveillance), 2012-13

In week 18, the number of influenza viruses detected in the Public Health Laboratory Centre was 108, including 84 influenza A (H1N1)2009 viruses, 19 influenza A (H3) viruses and 5 influenza B viruses (Figure 3).




Figure 3 Influenza virus detections, 2012-13

Influenza-like illness outbreak monitoring, 2012-13

In week 18, 4 ILI outbreaks occurring in schools/institutions were recorded (Figure 4). In the first 4 days of week 19 (May 5 to 8, 2013), 4 ILI outbreaks occurring in schools/institutions were recorded.




Figure 4 ILI outbreaks, 2012-13

Rate of influenza-like illness syndrome group in accident and emergency departments, 2011-13[#]

In week 18, the rate of the influenza-like illness syndrome group in the accident and emergency departments (AED) was 178.3 (per 1,000 coded cases) (Figure 5).

[#]Note: The influenza-like illness syndrome group includes codes such as influenza, upper respiratory tract infection, fever, cough, throat pain, and pneumonia.




Figure 5 Rate of ILI syndrome group in AED, 2011-13

Influenza associated hospital admission rates and deaths based on discharge coding, 2011-13

In week 18, hospital admission rates with principal diagnosis of influenza for persons aged 0-4 years, 5-64 years and 65 years or above were 0.42, 0.02 and 0.10 cases respectively (per 10,000 people in the age group) (Figure 6). Weekly number of deaths with any diagnosis of influenza in public hospitals is also shown in the figure on the right hand side.




Figure 6 Influenza associated hospital admission rates and deaths, 2011-13

Fever surveillance at sentinel child care centres/ kindergartens, 2011-13

In week 18, 0.74% of children in the sentinel child care centres / kindergartens (CCC/ KG) had fever (Figure 7).




Figure 7 Percentage of children with fever at sentinel CCC/ KG, 2011-13

Fever surveillance at sentinel residential care homes for the elderly, 2011-13

In week 18, 0.08% of residents in the sentinel residential care homes for the elderly (RCHEs) had fever (Figure 8).




Figure 8 Percentage of residents with fever at sentinel RCHE, 2011-13

Influenza-like illness surveillance among sentinel Chinese medicine practitioners, 2011-13

In week 18 the average consultation rate for ILI among Chinese medicine practitioners (CMPs) was 2.18 ILI cases per 1,000 consultations (Figure 9).




Figure 9 ILI consultation rate at sentinel CMP, 2011-13

Enhanced Surveillance for Severe Seasonal Influenza (Aged 18 years or above)

- From May 1, 2013, 12:00 noon to May 8, 2013, 12:00 noon, one case of influenza associated ICU admission was recorded. In addition, one previously reported cases who required ICU admission passed away during this reporting period.

Note: The data reported are provisional figures and subject to further revision.

Surveillance of severe paediatric influenza-associated complication/death (Aged above one month and below 18 years)

- In week 18 and the first 4 days of week 19 (May 5 to 8, 2013), there were no new reports of severe paediatric influenza-associated complication/death.

Note: The data reported are provisional figures and subject to further revision.

Surveillance of Tamiflu resistant influenza A(H1N1)2009 virus infection

- In week 18 and the first 4 days of week 19 (May 5 to 8, 2013), there were no new reports of oseltamivir (Tamiflu) resistant influenza A(H1N1)2009 virus infection. There are totally 30 reports of Tamiflu resistant influenza A(H1N1)2009 virus detected in Hong Kong since 2009.

Global Situation of Influenza Activity

- In the United States (week ending Apr 27, 2013), the influenza activity decreased.
- In Canada (week ending Apr 27, 2013), the influenza activity continued to decline.
- In the United Kingdom (week ending Apr 28, 2013), the weekly influenza consultation rate remained stable in England and Wales and decreased in Scotland and Northern Ireland.
- In Europe (week ending Apr 28, 2013), the ILI consultation rate was low throughout the Region.
- In Singapore (week ending Apr 27, 2013), the number of consultations for acute respiratory infections was below the warning level.

Sources:

Information have been extracted from the following sources when updates are available: [United States Centers for Disease Control and Prevention](#), [Public Health Agency of Canada](#), [Public Health England](#), [World Health Organization/Europe Euroflu](#) and [Singapore Ministry of Health](#).