

Quick Review of Linearity of Expectation

COMP 3711H - HKUST
Version of 24/11/2014
M. J. Golin

Linearity of Expectation is one of the simplest and most useful tools used in the analysis of randomized algorithms.

In its easiest form it just says that,
if X, Y are *any* two random variables (not necessarily independent) then

$$E(X + Y) = E(X) + E(Y).$$

The iterated version is that
if X_1, X_2, \dots, X_n are *any* random variables, then

$$E\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n E(X_i).$$

Example: Let Z be the value seen when rolling two dice.
 $Z = X_1 + X_2$ where X_i is the value seen when rolling single die $i = 1, 2$. It's easy to calculate that

$$E(X_i) = \sum_{j=1}^6 j \Pr(X_i = j) = \sum_{j=1}^6 j \frac{1}{6} = \frac{7}{2}.$$

Then

$$E(Z) = E(X_1 + X_2) = E(X_1) + E(X_2) = \frac{7}{2} + \frac{7}{2} = 7.$$

When flipping n coins, what is the expected number of heads?

$Z = \sum_{i=1}^n X_i$, where

$X_i = 1$ if coin i is a head and 0 if it is a tail.

Set $\Pr(X_i = 1) = p_i$ and $\Pr(X_i = 0) = 1 - p_i$.

Then X_i is a *Bernoulli Random Variable* with probability p_i .

Note that $E(X_i) = 1 \cdot \Pr(X_i = 1) = p_i$ so

$$E(Z) = \sum_{i=1}^n E(X_i) = \sum_{i=1}^n \Pr(X_i = 1) = \sum_{i=1}^n p_i.$$

Examples:

$$p_i = p \text{ (all coins the same)} \Rightarrow E(Z) = pn$$

$$p_i = \frac{1}{i} \Rightarrow E(Z) = \sum_{i=1}^n p_i = \sum_{i=1}^n \frac{1}{i} = H_n \sim \ln n$$

Suppose you are flipping n coins, each with $p_i = \frac{1}{2}$, i.e., fair coins. How many times does the pattern HHH appear?

Let x_1, x_2, \dots, x_n be the list of coin tosses, i.e., $x_i \in \{\text{H(ead)}, \text{T(ail)}\}$.

$$Z = \sum_{i=3}^n X_i \text{ where } X_i = 1 \text{ iff } x_{i-2}x_{i-1}x_i = HHH$$

$$\Pr(X_i = 1) = \frac{1}{8}, \text{ so}$$

$$E(Z) = \frac{n-2}{8}.$$

Suppose an algorithm's input is a permutation of n numbers.

Let x_1, x_2, \dots, x_n be the input in its given order.

x_i is a *left to right maxima* if it's bigger than x_1, x_2, \dots, x_{i-1} .

For example, the red items in these two permutations are the l.t.r. maxima:

5 4 7 8 1 6 3 2 1 3 5 7 2 4 6 8

Some algorithms' run times depend upon Z , the number of l.t.r. maxima. Assuming all $n!$ permutations are equally likely, how can we find $E(Z)$?

$Z = \sum_{i=1}^n X_i$ where $X_i = 1$ iff x_i is a l.t.r. maxima and 0 otherwise

One way of generating a random permutation is to first randomly choose the first i items equally likely among all possible $\binom{n}{i}$ subsets. Then choose a random permutation among the i possible permutations to order them as x_1, \dots, x_i . Then randomly order the remaining items as x_{i+1}, \dots, x_n .

Probability x_i is l.t.r. maxima is prob it's largest in first i items which is now $\frac{1}{i}$. So X_i is Bernouli Random Variable with $p_i = 1/i$ and $E(Z_i) = H_n$.

Suppose we flip n coins.

i th coin is Heads with probability $p_i = 1/i$.

If i th coin is Heads you get i dollars; Tails, you get nothing.

What is expected amount you receive?

Let Z be total amount. $Z = \sum_{i=1}^n X_i$ where $X_i = i$ if i th coin is heads and is otherwise 0. Then

$$E(X_i) = ip_i = 1$$

So,
$$E(Z) = E\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n E(X_i) = \sum_{i=1}^n 1 = n$$

Make a minor change from the previous page.

Suppose we flip n coins. i th coin is Heads with probability $p_i = 1/i$.

If i th coin is Heads you run another random process Y_i to tell you how much money you receive. All you know is that $E(Y_i) = i$.

If i th coin is Tails, you get nothing. What is expected amount you receive?

Let Z be total amount. $Z = \sum_{i=1}^n X_i Y_i$ where $X_i = 1$ if i th coin is Heads and is otherwise 0, so $E(X_i) = p_i$. $E(Y_i) = i$. Then, *because X_i and Y_i are independent*

$$E(X_i Y_i) = E(X_i) E(Y_i) = \frac{1}{i} i = 1$$

So,

$$E(Z) = E\left(\sum_{i=1}^n X_i Y_i\right) = \sum_{i=1}^n E(X_i Y_i) = \sum_{i=1}^n 1 = n$$