

# COMP170

## Discrete Mathematical Tools for Computer Science

### More on “*time until first success*”

*Version 1: Last updated, Nov 30, 2005*

# Example 1

## Example 1

Throw a fair die until you see a 1.

## Example 1

Throw a fair die until you see a 1.

Then throw it until you see a 2.

## Example 1

Throw a fair die until you see a 1.

Then throw it until you see a 2.

Continue until you see all of 3, 4, 5, 6, in that order.

## Example 1

Throw a fair die until you see a 1.

Then throw it until you see a 2.

Continue until you see all of 3, 4, 5, 6, in that order.

How many times, on average, do you throw the die?

## Example 1

Throw a fair die until you see a 1.

Then throw it until you see a 2.

Continue until you see all of 3, 4, 5, 6, in that order.

How many times, on average, do you throw the die?

Set  $X_1 = \#$  of throws until you see 1.

For  $i > 1$ :  $X_i = \#$  of throws, starting from when you see  $i - 1$  for the first time, until you see  $i$  for the first time.

## Example 1

Throw a fair die until you see a 1.

Then throw it until you see a 2.

Continue until you see all of 3, 4, 5, 6, in that order.

How many times, on average, do you throw the die?

Set  $X_1 = \#$  of throws until you see 1.

For  $i > 1$ :  $X_i = \#$  of throws, starting from when you see  $i - 1$  for the first time, until you see  $i$  for the first time.

$$\begin{array}{cccccc} \underbrace{2 \ 5 \ 1}_{X_1 = 3} & \underbrace{3 \ 1 \ 4 \ 2}_{X_2 = 4} & \underbrace{6 \ 5 \ 4 \ 4 \ 3}_{X_3 = 5} & \underbrace{1 \ 3 \ 4}_{X_4 = 3} & \underbrace{6 \ 4 \ 1 \ 2 \ 5}_{X_5 = 5} & \underbrace{1 \ 4 \ 3 \ 6}_{X_6 = 4} \end{array}$$



$$\begin{array}{cccccc}
 \underbrace{2 \ 5 \ 1}_{X_1 = 3} & \underbrace{3 \ 1 \ 4 \ 2}_{X_2 = 4} & \underbrace{6 \ 5 \ 4 \ 4 \ 3}_{X_3 = 5} & \underbrace{1 \ 3 \ 4}_{X_4 = 3} & \underbrace{6 \ 4 \ 1 \ 2 \ 5}_{X_5 = 5} & \underbrace{1 \ 4 \ 3 \ 6}_{X_6 = 4}
 \end{array}$$

Total number of throws is

$$X = X_1 + X_2 + \dots + X_6.$$

$$\begin{array}{cccccc}
 \underbrace{2 \ 5 \ 1}_{X_1 = 3} & \underbrace{3 \ 1 \ 4 \ 2}_{X_2 = 4} & \underbrace{6 \ 5 \ 4 \ 4 \ 3}_{X_3 = 5} & \underbrace{1 \ 3 \ 4}_{X_4 = 3} & \underbrace{6 \ 4 \ 1 \ 2 \ 5}_{X_5 = 5} & \underbrace{1 \ 4 \ 3 \ 6}_{X_6 = 4}
 \end{array}$$

Total number of throws is

$$X = X_1 + X_2 + \dots + X_6.$$

$$X = 19$$

$$\begin{array}{cccccc}
 \underbrace{2 \ 5 \ 1}_{X_1=3} & \underbrace{3 \ 1 \ 4 \ 2}_{X_2=4} & \underbrace{6 \ 5 \ 4 \ 4 \ 3}_{X_3=5} & \underbrace{1 \ 3 \ 4}_{X_4=3} & \underbrace{6 \ 4 \ 1 \ 2 \ 5}_{X_5=5} & \underbrace{1 \ 4 \ 3 \ 6}_{X_6=4}
 \end{array}$$

Total number of throws is

$$X = X_1 + X_2 + \dots + X_6. \qquad X = 19$$

$X_i$  is a geometric random variable with  $p = 1/6$ ,  
 $\Rightarrow E(X_i) = \frac{1}{p} = 6$  .

$$\begin{array}{cccccc}
 \underbrace{2 \ 5 \ 1}_{X_1=3} & \underbrace{3 \ 1 \ 4 \ 2}_{X_2=4} & \underbrace{6 \ 5 \ 4 \ 4 \ 3}_{X_3=5} & \underbrace{1 \ 3 \ 4}_{X_4=3} & \underbrace{6 \ 4 \ 1 \ 2 \ 5}_{X_5=5} & \underbrace{1 \ 4 \ 3 \ 6}_{X_6=4}
 \end{array}$$

Total number of throws is

$$X = X_1 + X_2 + \dots + X_6. \qquad X = 19$$

$X_i$  is a geometric random variable with  $p = 1/6$ ,  
 $\Rightarrow E(X_i) = \frac{1}{p} = 6$ .

Then, by linearity of expectation,

$$E(X) = E(X_1) + E(X_2) + \dots + E(X_6) = 6 \cdot 6 = 36.$$

## Example 2

## Example 2

Throw a fair die until you have seen all 6 numbers.

## Example 2

Throw a fair die until you have seen all 6 numbers.

How many times do you throw the die on average?

## Example 2

Throw a fair die until you have seen all 6 numbers.

How many times do you throw the die on average?

Let  $N_i$  be the  $i$ th new number we see.



## Example 2

Throw a fair die until you have seen all 6 numbers.

How many times do you throw the die on average?

Let  $N_i$  be the  $i$ th new number we see.

Let  $X_1 = 1$ :

For  $i > 1$ :  $X_i = \#$  of throws needed to get  $N_i$  after first time we see  $N_{i-1}$ .

## Example 2

Throw a fair die until you have seen all 6 numbers.

How many times do you throw the die on average?

Let  $N_i$  be the  $i$ th new number we see.

Let  $X_1 = 1$ :

For  $i > 1$ :  $X_i = \#$  of throws needed to get  
 $N_i$  after first time we see  $N_{i-1}$ .

Example

2 2 5 2 3 3 2 3 6 5 2 6 5 1 2 1 5 6 1 3 4

## Example 2

Throw a fair die until you have seen all 6 numbers.

How many times do you throw the die on average?

Let  $N_i$  be the  $i$ th new number we see.

Let  $X_1 = 1$ :

For  $i > 1$ :  $X_i = \#$  of throws needed to get  $N_i$  after first time we see  $N_{i-1}$ .

Example

	2	2	5	2	3	3	2	3	6	5	2	6	5	1	2	1	5	6	1	3	4
$i$	1	2	3						4					5							6
$N_i$	2	5	3						6					1							4
$X_i$	1	2	2						4					5							7

## Example 2 (cont'd)

## Example 2 (cont'd)

$$X = X_1 + X_2 + X_3 + X_4 + X_5 + X_6$$

## Example 2 (cont'd)

$$X = X_1 + X_2 + X_3 + X_4 + X_5 + X_6$$

$$\Rightarrow E(X) = E(X_1) + E(X_2) + E(X_3) + E(X_4) + E(X_5) + E(X_6)$$

## Example 2 (cont'd)

$$X = X_1 + X_2 + X_3 + X_4 + X_5 + X_6$$

$$\Rightarrow E(X) = E(X_1) + E(X_2) + E(X_3) + E(X_4) + E(X_5) + E(X_6)$$

so we need to calculate all of the  $E(X_i)$

## Example 2 (cont'd)

$$X = X_1 + X_2 + X_3 + X_4 + X_5 + X_6$$

$$\Rightarrow E(X) = E(X_1) + E(X_2) + E(X_3) + E(X_4) + E(X_5) + E(X_6)$$

so we need to calculate all of the  $E(X_i)$

$i = 1$ :  $X_1 = 1$  so  $E(X_1) = 1$ .



## Example 2 (cont'd)

$$X = X_1 + X_2 + X_3 + X_4 + X_5 + X_6$$

$$\Rightarrow E(X) = E(X_1) + E(X_2) + E(X_3) + E(X_4) + E(X_5) + E(X_6)$$

so we need to calculate all of the  $E(X_i)$

$i = 1$ :  $X_1 = 1$  so  $E(X_1) = 1$ .

$i = 2$ : Once  $N_1$  is chosen,  $X_2$  is the number of times we need to throw the die until we see something that is *not*  $N_1$ . Since being "*not*  $N_1$ " occurs with probability  $\frac{5}{6}$ ,  $X_2$  is geometric with  $p = \frac{5}{6}$  so  $E(X_2) = \frac{1}{p} = \frac{6}{5}$ .

## Example 2 (cont'd)

$$X = X_1 + X_2 + X_3 + X_4 + X_5 + X_6$$

$$\Rightarrow E(X) = E(X_1) + E(X_2) + E(X_3) + E(X_4) + E(X_5) + E(X_6)$$

so we need to calculate all of the  $E(X_i)$

$i = 1$ :  $X_1 = 1$  so  $E(X_1) = 1$ .

$i = 2$ : Once  $N_1$  is chosen,  $X_2$  is the number of times we need to throw the die until we see something that is *not*  $N_1$ . Since being "*not*  $N_1$ " occurs with probability  $\frac{5}{6}$ ,  $X_2$  is geometric with  $p = \frac{5}{6}$  so  $E(X_2) = \frac{1}{p} = \frac{6}{5}$ .

$i = 3$ : Similarly, once  $N_1, N_2$  are chosen,  $X_3$  is the number of times we need to throw the die until we see something that is *not*  $N_1, N_2$ . Since being "*not*  $N_1, N_2$ " occurs with probability  $\frac{4}{6}$ ,  $X_3$  is geometric with  $p = \frac{4}{6}$  so  $E(X_3) = \frac{1}{p} = \frac{6}{4}$ .

## Example 2 (cont'd)

$$X = X_1 + X_2 + X_3 + X_4 + X_5 + X_6$$

## Example 2 (cont'd)

$$X = X_1 + X_2 + X_3 + X_4 + X_5 + X_6$$

General  $i$  In the general case, once  $N_1, N_2, \dots, N_{i-1}$  are chosen,  $X_i$  is the number of times we need to throw the die until we see something that is *not*  $N_1, N_2, \dots, N_{i-1}$ . Since being “*not*  $N_1, N_2, \dots, N_{i-1}$ ” occurs with probability  $\frac{6-(i-1)}{6}$ ,  $X_i$  is geometric with  $p = \frac{6-(i-1)}{6}$  so  $E(X_i) = \frac{1}{p} = \frac{6}{6-(i-1)}$ .

$$\Rightarrow E(X) = E(X_1) + E(X_2) + E(X_3) + E(X_4) + E(X_5) + E(X_6)$$

## Example 2 (cont'd)

$$X = X_1 + X_2 + X_3 + X_4 + X_5 + X_6$$

General  $i$  In the general case, once  $N_1, N_2, \dots, N_{i-1}$  are chosen,  $X_i$  is the number of times we need to throw the die until we see something that is *not*  $N_1, N_2, \dots, N_{i-1}$ . Since being “*not*  $N_1, N_2, \dots, N_{i-1}$ ” occurs with probability  $\frac{6-(i-1)}{6}$ ,  $X_i$  is geometric with  $p = \frac{6-(i-1)}{6}$  so  $E(X_i) = \frac{1}{p} = \frac{6}{6-(i-1)}$ .

$$\begin{aligned} \Rightarrow E(X) &= E(X_1) + E(X_2) + E(X_3) + E(X_4) + E(X_5) + E(X_6) \\ &= \frac{6}{6} + \frac{6}{5} + \dots + \frac{6}{1} = 6 \sum_{i=1}^6 \frac{1}{i} = 6 \cdot \frac{49}{20} = \frac{147}{10}. \end{aligned}$$

Compare this to previous problem in which we needed  $6 \cdot 6$  flips on average, to see the numbers in order.

## Example 3

## Example 3

This is known as the coupon collectors problem.

## Example 3

This is known as the coupon collectors problem.

There are  $n$  coupons.



## Example 3

This is known as the coupon collectors problem.

There are  $n$  coupons.

If you collect all of them, you win a prize.

## Example 3

This is known as the coupon collectors problem.

There are  $n$  coupons.

If you collect all of them, you win a prize.

Each time you go to the store, you get a random coupon.

## Example 3

This is known as the coupon collectors problem.

There are  $n$  coupons.

If you collect all of them, you win a prize.

Each time you go to the store, you get a random coupon.

How long do you need (in expectation) to collect all of the coupons?

## Example 3

This is known as the coupon collectors problem.

There are  $n$  coupons.

If you collect all of them, you win a prize.

Each time you go to the store, you get a random coupon.

How long do you need (in expectation) to collect all of the coupons?

$$X = X_1 + X_2 + \dots + X_n.$$

## Example 3

This is known as the coupon collectors problem.

There are  $n$  coupons.

If you collect all of them, you win a prize.

Each time you go to the store, you get a random coupon.

How long do you need (in expectation) to collect all of the coupons?

$$X = X_1 + X_2 + \dots + X_n.$$

$$X_1 = 1.$$

For  $i > 1$  :  $X_i$  = time needed to receive  $i$ th new coupon after having received  $(i - 1)$ st new coupon.

## Example 3 (cont'd)

### Example 3 (cont'd)

$X_i$  is geometric with  $p = (n - (i - 1))/n$ .

### Example 3 (cont'd)

$X_i$  is geometric with  $p = (n - (i - 1))/n$ .

So,

$$E(X_i) = \frac{n}{n - (i - 1)}$$



### Example 3 (cont'd)

$X_i$  is geometric with  $p = (n - (i - 1))/n$ .

So,

$$E(X_i) = \frac{n}{n - (i - 1)}$$

$$E(X) = \sum_{i=1}^n E(X_i) = \sum_{i=1}^n \frac{n}{n - (i - 1)} = n \sum_{i=1}^n \frac{1}{i}$$

We just showed that

$$E(X) = n \sum_{i=1}^n \frac{1}{i}$$

$H_n = \sum_{i=1}^n 1/i$  has a special name. It is called the  $n^{\text{th}}$  **harmonic number**.

It is also known that  $\forall n \ |H_n - \ln n| \leq 2$ .

So  $H_n$  grows like  $\ln n$  and

$E(X)$  grows like  $n \ln n$ .