

# Implicit Function

Siu-Wing Cheng

Room 3514

Phone: 2358-6973

[scheng@cse.ust.hk](mailto:scheng@cse.ust.hk)

<http://www.cse.ust.hk/faculty/scheng>

Department of Computer Science and Engineering  
The Hong Kong University of Science and Technology  
Hong Kong

# Implicit Function in $\mathbb{R}^2$

In general, a function  $f(x_1, x_2) = 0$  describes a curve in  $\mathbb{R}^2$ .

Examples:

- $x_1^2 + x_2^2 = 1$ .
- $x_1^2 - x_2 = 0$ .

# Implicit Function in $\mathbb{R}^2$

In general, a function  $f(x_1, x_2) = 0$  describes a curve in  $\mathbb{R}^2$ .

Examples:

- $x_1^2 + x_2^2 = 1$ .
- $x_1^2 - x_2 = 0$ .

What are the sets of points  $f(x_1, x_2) > 0$  and  $f(x_1, x_2) < 0$ ?

# Implicit Function in $\mathbb{R}^3$

In general, a function  $f(x_1, x_2, x_3) = 0$  describes a surface in  $\mathbb{R}^3$ .

Examples:

- $x_1^2 + x_2^2 + x_3^2 = 1$ .
- $x_1^2 + x_2^2 - x_3 = 0$ .

# Implicit Function in $\mathbb{R}^3$

In general, a function  $f(x_1, x_2, x_3) = 0$  describes a surface in  $\mathbb{R}^3$ .

Examples:

- $x_1^2 + x_2^2 + x_3^2 = 1$ .
- $x_1^2 + x_2^2 - x_3 = 0$ .

What are the sets of points  $f(x_1, x_2) > 0$  and  $f(x_1, x_2) < 0$ ?

# Approximation Surface in $\mathbb{R}^3$

Given a set  $P$  of point samples, our goal is to construct an implicit function  $f(x_1, x_2, x_3) = 0$  so that  $f$  approximates the unknown surface.

# Approximation Surface in $\mathbb{R}^3$

Given a set  $P$  of point samples, our goal is to construct an implicit function  $f(x_1, x_2, x_3) = 0$  so that  $f$  approximates the unknown surface.

Once we know  $f$ , we can then use  $f$  to compute the restricted Delaunay triangulation. The triangulation gives us a triangulated surface that approximates the unknown surface.

# Approximation Surface in $\mathbb{R}^3$

Desirable properties of  $f$ :



# Approximation Surface in $\mathbb{R}^3$

Desirable properties of  $f$ :

- The surface  $f(x_1, x_2, x_3) = 0$  should pass close to the given point samples.

Desirable properties of  $f$ :

- The surface  $f(x_1, x_2, x_3) = 0$  should pass close to the given point samples.
- The surface normals of  $f(x_1, x_2, x_3) = 0$  in the vicinity of the given point samples should approximate the surface normals of the unknown surface at these point samples.

Desirable properties of  $f$ :

- The surface  $f(x_1, x_2, x_3) = 0$  should pass close to the given point samples.
- The surface normals of  $f(x_1, x_2, x_3) = 0$  in the vicinity of the given point samples should approximate the surface normals of the unknown surface at these point samples.
- The function  $f(x_1, x_2, x_3)$  can be evaluated quickly.

A vector  $\mathbf{n}$  in  $\mathbb{R}^3$  has a direction and a magnitude. It is represented by three coordinates  $(n_1, n_2, n_3)$  just like a point.

The representation  $(n_1, n_2, n_3)$  means that  $\mathbf{n}$  can be viewed as the line segment directed from the origin to the point  $(n_1, n_2, n_3)$ . However, we are actually free to translate  $\mathbf{n}$ . So  $\mathbf{n}$  is not fixed in any particular position.

The magnitude of  $\mathbf{n}$  is equal to  $\sqrt{n_1^2 + n_2^2 + n_3^2}$ . This can be viewed as the length of the vector.

Given two points  $x, y \in \mathbb{R}^3$ , we can use  $x - y$  to denote the vector directed from  $y$  to  $x$ . Its representation is

$$(x_1 - y_1, x_2 - y_2, x_3 - y_3).$$

# Surface Normal

Given a point  $p$  on a surface,

Given a point  $p$  on a surface,

- A surface normal at  $p$  is a vector that is orthogonal to the surface at  $p$ .

Given a point  $p$  on a surface,

- A surface normal at  $p$  is a vector that is orthogonal to the surface at  $p$ .
- A unit normal at  $p$  is a normal with unit length.



Given a point  $p$  on a surface,

- A surface normal at  $p$  is a vector that is orthogonal to the surface at  $p$ .
- A unit normal at  $p$  is a normal with unit length.
- A unit outward normal at  $p$  is a unit normal that points outside the surface.

# A Motivating Example

Suppose that the unknown surface is a plane. Let  $p$  be a given point sample on the plane.

Suppose that we known the normal  $\mathbf{n}_p$  to the plane at the point  $p$ . Given a point  $x \in \mathbb{R}^3$ , how would you test whether  $x$  lies the plane or not?

# A Motivating Example

Suppose that the unknown surface is a plane. Let  $p$  be a given point sample on the plane.

Suppose that we known the normal  $\mathbf{n}_p$  to the plane at the point  $p$ . Given a point  $x \in \mathbb{R}^3$ , how would you test whether  $x$  lies the plane or not?

The point  $x$  lies on the plane iff the inner product  $\langle \mathbf{n}_p, (x - p) \rangle = 0$ .

# A Motivating Example

Suppose that the unknown surface is a plane. Let  $p$  be a given point sample on the plane.

Suppose that we known the normal  $\mathbf{n}_p$  to the plane at the point  $p$ . Given a point  $x \in \mathbb{R}^3$ , how would you test whether  $x$  lies the plane or not?

The point  $x$  lies on the plane iff the inner product  $\langle \mathbf{n}_p, (x - p) \rangle = 0$ .

The inner product  $\langle \mathbf{n}_p, (x - p) \rangle$  is equal to

$$|\mathbf{n}_p| \cdot |x - p| \cdot \cos \angle \mathbf{n}_p, (x - p).$$

It can also be evaluated as

$$n_{p,1}(x_1 - p_1) + n_{p,2}(x_2 - p_2) + n_{p,3}(x_3 - p_3).$$

# A Motivating Example

Since any surface is quite flat in a small neighborhood around  $p$ , given a point  $x$  very close to  $p$ , we can use

$$\langle \mathbf{n}_p, (x - p) \rangle = 0$$

to tell if  $x$  is on the surface or not.

# A Motivating Example

Since any surface is quite flat in a small neighborhood around  $p$ , given a point  $x$  very close to  $p$ , we can use

$$\langle \mathbf{n}_p, (x - p) \rangle = 0$$

to tell if  $x$  is on the surface or not.

If a point  $x$  is far from  $p$ , we should find another given point sample  $q$  very close to  $x$ , compute  $\langle \mathbf{n}_q, (x - q) \rangle$ , and make the decision.

# Local Support and Weight Function

Define a weight function  $w : P \times \mathbb{R}^3 \rightarrow \mathbb{R}$ :

$$w(p, x) = \frac{e^{-(d(p,x)/r(x))^2}}{\sum_{q \in P} e^{-(d(q,x)/r(x))^2}}$$

where  $r(x)$  is the distance from  $x$  to the fifth nearest sample point in  $P$ .

# Local Support and Weight Function

Define a weight function  $w : P \times \mathbb{R}^3 \rightarrow \mathbb{R}$ :

$$w(p, x) = \frac{e^{-(d(p,x)/r(x))^2}}{\sum_{q \in P} e^{-(d(q,x)/r(x))^2}}$$

where  $r(x)$  is the distance from  $x$  to the fifth nearest sample point in  $P$ .

Intuitively speaking,

- $w(p, x)$  is a value between 0 and 1.
- $w(p, x)$  increases as  $x$  moves towards  $p$ .
- $w(p, x)$  drops rapidly as  $x$  moves away from  $p$ .



# An Implicit Function

$$f(x) = \sum_{p \in P} \langle \mathbf{n}_p, (x - p) \rangle \cdot w(p, x)$$

# An Implicit Function

$$f(x) = \sum_{p \in P} \langle \mathbf{n}_p, (x - p) \rangle \cdot w(p, x)$$

Remaining issue: How to obtain  $\mathbf{n}_p$ ?

$$f(x) = \sum_{p \in P} \langle \mathbf{n}_p, (x - p) \rangle \cdot w(p, x)$$

Remaining issue: How to obtain  $\mathbf{n}_p$ ?

Can be done using Principal Component Analysis.  
Details skipped.