

This small handout is a worked example to help clarify the difference between

- Sample spaces

and

- Probability distributions on the sample spaces

This small handout is a worked example to help clarify the difference between

- Sample spaces
- and
- Probability distributions on the sample spaces

In particular, you will see two **different** probability distributions on the **same** sample space.

This small handout is a worked example to help clarify the difference between

- Sample spaces

and

- Probability distributions on the sample spaces

In particular, you will see two **different** probability distributions on the **same** sample space.

You will also see an example illustrating why the ***inclusion-exclusion formula*** is actually a statement about sets, and does not depend upon the actual weights of the probability distribution.

Consider the sample space of 3 coin flips (8 outcomes)

$$E = \{\text{first flip is Head}\} = \{HHT, HTT, HHH, HTH\}$$

$$F = \{\text{last flip is Head}\} = \{THH, TTH, HTH, HHH\}$$

$$E \cap F = \{\text{first \& last flips are Heads}\} = \{HHH, HTH\}$$

Consider the sample space of 3 coin flips (8 outcomes)


$E = \{\text{first flip is Head}\}$

$= \{HHT, HTT, HHH, HTH\}$

$F = \{\text{last flip is Head}\}$

$= \{THH, TTH, HTH, HHH\}$

$E \cap F = \{\text{first \& last flips are Heads}\} = \{HHH, HTH\}$



$$|E| = 4$$

$$|F| = 4$$

$$|E \cap F| = 2$$

Consider the sample space of 3 coin flips (8 outcomes)

$E = \{\text{first flip is Head}\}$


$F = \{\text{last flip is Head}\}$

$E \cap F = \{\text{first \& last flips are Heads}\}$

$= \{HHT, HTT, HHH, HTH\}$

$= \{THH, TTH, HTH, HHH\}$

$= \{HHH, HTH\}$



$$|E| = 4$$

$$|F| = 4$$

$$|E \cap F| = 2$$

On the left we first look
at the *size* of subsets.

Consider the sample space of 3 coin flips (8 outcomes)

$E = \{\text{first flip is Head}\}$


$F = \{\text{last flip is Head}\}$

$E \cap F = \{\text{first \& last flips are Heads}\}$

$= \{HHT, HTT, HHH, HTH\}$

$= \{THH, TTH, HTH, HHH\}$

$= \{HHH, HTH\}$



$$|E| = 4$$

$$|F| = 4$$

$$|E \cap F| = 2$$

On the left we first look at the *size* of subsets.

Now, assume that the coin is fair so each outcome has the same probability, $\frac{1}{8}$, of occurring.

Consider the sample space of 3 coin flips (8 outcomes)

$E = \{\text{first flip is Head}\}$


$F = \{\text{last flip is Head}\}$

$E \cap F = \{\text{first \& last flips are Heads}\}$

$= \{HHT, HTT, HHH, HTH\}$

$= \{THH, TTH, HTH, HHH\}$


$= \{HHH, HTH\}$



$$|E| = 4$$

$$|F| = 4$$

$$|E \cap F| = 2$$




$$P(E) = 4 \frac{1}{8} = \frac{1}{2}$$

$$P(F) = 4 \frac{1}{8} = \frac{1}{2}$$

$$P(E \cap F) = 2 \frac{1}{8} = \frac{1}{4}$$

On the left we first look at the *size* of subsets.

Now, assume that the coin is fair so each outcome has the same probability, $\frac{1}{8}$, of occurring.




$$P(E) = 4 \frac{1}{8} = \frac{1}{2}$$

$$P(F) = 4 \frac{1}{8} = \frac{1}{2}$$

$$P(E \cap F) = 2 \frac{1}{8} = \frac{1}{4}$$

We just saw the probabilities for fair coins.

This created a *uniform* distribution on the sample space.



$$P(E) = 4 \frac{1}{8} = \frac{1}{2}$$


$$P(F) = 4 \frac{1}{8} = \frac{1}{2}$$

$$P(E \cap F) = 2 \frac{1}{8} = \frac{1}{4}$$

We just saw the probabilities for fair coins.

This created a *uniform* distribution on the sample space.

Now assume that the coin is *not* fair. Instead $P(H) = \frac{1}{3}$. In this case we get a *different* probability distribution on the *same* sample space.




$$P(E) = 4 \frac{1}{8} = \frac{1}{2}$$

$$P(F) = 4 \frac{1}{8} = \frac{1}{2}$$

$$P(E \cap F) = 2 \frac{1}{8} = \frac{1}{4}$$

We just saw the probabilities for fair coins.

This created a *uniform* distribution on the sample space.




$$P(E) = \frac{1+2+2+4}{27} = \frac{1}{3}$$

$$P(F) = \frac{1+2+2+4}{27} = \frac{1}{3}$$

$$P(E \cap F) = \frac{1+2}{27} = \frac{1}{9}$$

Now assume that the coin is *not* fair. Instead $P(H) = \frac{1}{3}$. In this case we get a *different* probability distribution on the *same* sample space.

As a final note, we point out that the inclusion-exclusion formula is really a statement about the **sample space** and *does not depend* upon the actual weights of the **probability distribution**




As a final note, we point out that the inclusion-exclusion formula is really a statement about the **sample space** and *does not depend* upon the actual weights of the **probability distribution**

As an example, consider the $n = 2$ case to the right.

Working through the sets, we see that, **independent of the actual probability weights,**


$$P(E \cup F) = P(E) + P(F) - P(E \cap F)$$



As a final note, we point out that the inclusion-exclusion formula is really a statement about the **sample space** and *does not depend* upon the actual weights of the **probability distribution**

As an example, consider the $n = 2$ case to the right.

Working through the sets, we see that, **independent of the actual probability weights,**



$$P(E \cup F) = P(E) + P(F) - P(E \cap F)$$

$$P(E) = P(\{HHT\}) + P(\{HTT\}) + P(\{HHH\}) + P(\{HTH\})$$

$$P(F) = P(\{THH\}) + P(\{TTH\}) + P(\{HHH\}) + P(\{HTH\})$$

$$P(E \cap F) = P(\{HHH\}) + P(\{HTH\})$$

$$P(E \cup F) = P(\{HHT\}) + P(\{HTT\}) + P(\{THH\}) + P(\{TTH\}) + P(\{HHH\}) + P(\{HTH\})$$