

COMP170

Discrete Mathematical Tools for Computer Science

Recursion, Recurrences and Induction

Version 2.0: Last updated, May 13, 2007

Discrete Math for Computer Science

K. Bogart, C. Stein and R.L. Drysdale

Section 4.2, pp. 143-153

Recursion, Recurrences and Induction

- Recursion
- Recurrences
- Iterating a Recurrence
- Geometric Series
- First-Order Linear Recurrences

Recursion

Recursion

- Recursive computer programs or algorithms often lead to inductive analyses

Recursion

- Recursive computer programs or algorithms often lead to inductive analyses
- A classic example of this is the Towers of Hanoi problem

Towers of Hanoi

Towers of Hanoi



Towers of Hanoi



- 3 pegs; n disks of different sizes.
- A **legal move** takes a disk from one peg and moves it onto another peg so that it is not on top of a smaller disk
- Problem: Find a (efficient) way to move all of the disks from one peg to another


Towers of Hanoi



Towers of Hanoi




legal move




Towers of Hanoi



legal move




legal move




Towers of Hanoi




legal move



legal move




not legal




Towers of Hanoi




legal move




legal move



not legal



legal move



Towers of Hanoi

Problem

Towers of Hanoi

Problem

Start with n disks
on leftmost peg



Towers of Hanoi

Problem

Start with n disks
on leftmost peg

using only
legal moves




Towers of Hanoi

Problem

Start with n disks
on leftmost peg

move all disks to
rightmost peg.

using only
legal moves




Towers of Hanoi

Problem

Start with n disks
on leftmost peg

move all disks to
rightmost peg.

using only
legal moves



Given $i, j \in \{1, 2, 3\}$ let $\overline{\{i, j\}} = \{1, 2, 3\} - \{i\} - \{j\}$

i.e., $\overline{\{1, 2\}} = 3$, $\overline{\{1, 3\}} = 2$, $\overline{\{2, 3\}} = 1$.

Towers of Hanoi

General Solution

Towers of Hanoi

General Solution

Recursion Base:

If $n = 1$ moving one disk from i to j is easy.
Just move it.




Towers of Hanoi

General Solution

Recursion Base:

If $n = 1$ moving one disk from i to j is easy.
Just move it.



Towers of Hanoi




To move $n > 1$ disks
from i to j

Towers of Hanoi



To move $n > 1$ disks
from i to j

1)




move top $\frac{n-1}{\{i, j\}}$ disks
from i to $\{i, j\}$

Towers of Hanoi




To move $n > 1$ disks
from i to j

1)



move top $\frac{n-1}{2}$ disks
from i to $\{i, j\}$

2)




move largest disk
from i to j

Towers of Hanoi




To move $n > 1$ disks
from i to j

1)



move top $\frac{n-1}{2}$ disks
from i to $\{i, j\}$

2)



move largest disk
from i to j

3)



move top $\frac{n-1}{2}$ disks
from $\{i, j\}$ to j .

To move n disks from i to j

i) move top $\overline{n - 1}$ disks
from i to $\overline{\{i, j\}}$

ii) move largest disk
from i to j

iii) move top $\overline{n - 1}$ disks
from $\overline{\{i, j\}}$ to j .

- To prove **Correctness** of solution we are implicitly using induction

To move n disks from i to j

i) move top $n - 1$ disks
from i to $\overline{\{i, j\}}$

ii) move largest disk
from i to j

iii) move top $n - 1$ disks
from $\overline{\{i, j\}}$ to j .

- To prove **Correctness** of solution we are implicitly using induction
- $p(n)$ is statement that algorithm is correct for n

To move n disks from i to j

i) move top $n - 1$ disks
from i to $\overline{\{i, j\}}$

ii) move largest disk
from i to j

iii) move top $n - 1$ disks
from $\overline{\{i, j\}}$ to j .

- To prove **Correctness** of solution we are implicitly using induction

- $p(n)$ is statement that algorithm is correct for n

To move n disks from i to j

i) move top $n - 1$ disks from i to $\overline{\{i, j\}}$

ii) move largest disk from i to j

iii) move top $n - 1$ disks from $\overline{\{i, j\}}$ to j .

- $p(1)$ is statement that algorithm works for $n = 1$ disks, which is obviously true

- To prove **Correctness** of solution we are implicitly using induction
- $p(n)$ is statement that algorithm is correct for n

To move n disks from i to j

i) move top $n - 1$ disks from i to $\overline{\{i, j\}}$

ii) move largest disk from i to j

iii) move top $n - 1$ disks from $\overline{\{i, j\}}$ to j .

- $p(1)$ is statement that algorithm works for $n = 1$ disks, which is obviously true
- $p(n - 1) \Rightarrow p(n)$ is “recursion” statement that if our algorithm works for $n - 1$ disks, then we can build a correct solution for n disks

Running Time

$M(n)$ is number of disk moves needed for n disks

To move n disks from i to j

i) move top $n - 1$ disks from i to $\overline{\{i, j\}}$

ii) move largest disk from i to j

iii) move top $n - 1$ disks from $\overline{\{i, j\}}$ to j .

Running Time

$M(n)$ is number of disk moves needed for n disks

To move n disks from i to j

i) move top $n - 1$ disks from i to $\overline{\{i, j\}}$

ii) move largest disk from i to j

iii) move top $n - 1$ disks from $\overline{\{i, j\}}$ to j .

- $M(1) = 1$

Running Time

$M(n)$ is number of disk moves needed for n disks

To move n disks from i to j

i) move top $n - 1$ disks from i to $\overline{\{i, j\}}$

ii) move largest disk from i to j

iii) move top $n - 1$ disks from $\overline{\{i, j\}}$ to j .

- $M(1) = 1$
- If $n > 1$, then $M(n) = 2M(n - 1) + 1$

- We saw that $M(1) = 1$ and that

- We saw that $M(1) = 1$ and that
- $M(n) = 2M(n - 1) + 1$ for $n > 1$.

- We saw that $M(1) = 1$ and that
- $M(n) = 2M(n - 1) + 1$ for $n > 1$.
- Iterating the recurrence gives
$$M(1) = 1, \quad M(2) = 3, \quad M(3) = 7,$$
$$M(4) = 15, \quad M(5) = 31, \quad \dots$$

- We saw that $M(1) = 1$ and that
- $M(n) = 2M(n - 1) + 1$ for $n > 1$.
- Iterating the recurrence gives
$$M(1) = 1, \quad M(2) = 3, \quad M(3) = 7,$$
$$M(4) = 15, \quad M(5) = 31, \quad \dots$$
- We guess that $M(n) = 2^n - 1$.

- We saw that $M(1) = 1$ and that
- $M(n) = 2M(n - 1) + 1$ for $n > 1$.
- Iterating the recurrence gives
$$M(1) = 1, \quad M(2) = 3, \quad M(3) = 7, \\ M(4) = 15, \quad M(5) = 31, \dots$$
- We guess that $M(n) = 2^n - 1$.
 - We'll prove this by induction

- We saw that $M(1) = 1$ and that
- $M(n) = 2M(n - 1) + 1$ for $n > 1$.
- Iterating the recurrence gives
$$M(1) = 1, \quad M(2) = 3, \quad M(3) = 7, \\ M(4) = 15, \quad M(5) = 31, \dots$$
- We guess that $M(n) = 2^n - 1$.
 - We'll prove this by induction
 - Later, we'll see how to solve without guessing

Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n-1) + 1 & \text{otherwise.} \end{cases}$$

we show that $M(n) = 2^n - 1$

Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n-1) + 1 & \text{otherwise.} \end{cases}$$

we show that $M(n) = 2^n - 1$

Proof: (by induction),

Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n-1) + 1 & \text{otherwise.} \end{cases}$$

we show that $M(n) = 2^n - 1$

Proof: (by induction),

The base case $n = 1$ is true, since $2^1 - 1 = 1$.

Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n-1) + 1 & \text{otherwise.} \end{cases}$$

we show that $M(n) = 2^n - 1$

Proof: (by induction),

The base case $n = 1$ is true, since $2^1 - 1 = 1$.

For the inductive step, assume that

$$M(n-1) = 2^{n-1} - 1 \text{ for } n > 1.$$

Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n-1) + 1 & \text{otherwise.} \end{cases}$$

we show that $M(n) = 2^n - 1$

Proof: (by induction),

The base case $n = 1$ is true, since $2^1 - 1 = 1$.

For the inductive step, assume that

$M(n-1) = 2^{n-1} - 1$ for $n > 1$. Then

$$M(n) = 2M(n-1) + 1$$

Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n-1) + 1 & \text{otherwise.} \end{cases}$$

we show that $M(n) = 2^n - 1$

Proof: (by induction),

The base case $n = 1$ is true, since $2^1 - 1 = 1$.

For the inductive step, assume that

$M(n-1) = 2^{n-1} - 1$ for $n > 1$. Then

$$M(n) = 2M(n-1) + 1 \quad \text{def}$$

Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n-1) + 1 & \text{otherwise.} \end{cases}$$

we show that $M(n) = 2^n - 1$

Proof: (by induction),

The base case $n = 1$ is true, since $2^1 - 1 = 1$.

For the inductive step, assume that

$M(n-1) = 2^{n-1} - 1$ for $n > 1$. Then

$$\begin{aligned} M(n) &= 2M(n-1) + 1 && \text{def} \\ &= 2(2^{n-1} - 1) + 1 \end{aligned}$$

Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n-1) + 1 & \text{otherwise.} \end{cases}$$

we show that $M(n) = 2^n - 1$

Proof: (by induction),

The base case $n = 1$ is true, since $2^1 - 1 = 1$.

For the inductive step, assume that

$M(n-1) = 2^{n-1} - 1$ for $n > 1$. Then

$$\begin{aligned} M(n) &= 2M(n-1) + 1 && \text{def} \\ &= 2(2^{n-1} - 1) + 1 && \text{ind hyp} \end{aligned}$$

Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n-1) + 1 & \text{otherwise.} \end{cases}$$

we show that $M(n) = 2^n - 1$

Proof: (by induction),

The base case $n = 1$ is true, since $2^1 - 1 = 1$.

For the inductive step, assume that

$M(n-1) = 2^{n-1} - 1$ for $n > 1$. Then

$$\begin{aligned} M(n) &= 2M(n-1) + 1 && \text{def} \\ &= 2(2^{n-1} - 1) + 1 && \text{ind hyp} \\ &= 2^n - 1. && \text{ind conc} \end{aligned}$$

Note that we used induction **twice**

Note that we used induction **twice**

The first time was to **derive** Correctness of Algorithm and the **recurrence**

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n - 1) + 1 & \text{otherwise.} \end{cases}$$

Note that we used induction **twice**

The first time was to **derive** Correctness of Algorithm and the **recurrence**

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n-1) + 1 & \text{otherwise.} \end{cases}$$

The second time was to **derive the closed form solution**

$$M(n) = 2^n - 1$$

of the recurrence

Francois Edouard Anatole Lucas

b. 1842, d. 1891

French mathematician.

Best known for his results in
number theory.

He is also famous for being a
creator of mathematical puzzles,
among the most well-known of
which is the Tower of Hanoi puzzle
(1883).



Recursion, Recurrences and Induction

- Recursion
- Recurrences
- Iterating a Recurrence
- Geometric Series
- First-Order Linear Recurrences

A **recurrence equation** or **recurrence** for a function defined on the set of integers greater than or equal to some number b is one that tells us how to compute the n th value from some or all the first $(n - 1)$ values.

To completely specify a function on the basis of a recurrence, we have to give the **initial condition(s)** (also called the *base case(s)*) for the recurrence.

A **recurrence equation** or **recurrence** for a function defined on the set of integers greater than or equal to some number b is one that tells us how to compute the n th value from some or all the first $(n - 1)$ values.

To completely specify a function on the basis of a recurrence, we have to give the **initial condition(s)** (also called the *base case(s)*) for the recurrence.

$$M(n) = \begin{cases} 1 & \text{if } n = 1, \\ 2M(n - 1) + 1 & \text{otherwise.} \end{cases} \quad \text{Towers of Hanoi}$$

$$F(n) = \begin{cases} 1 & \text{if } n = 0, 1, \\ F(n - 1) + F(n - 2) & \text{otherwise.} \end{cases} \quad \text{Fibonacci Numbers}$$

Example 2:

Example 2:

Let $S(n)$ be the number of subsets of a set of size n .
What is the formula for $S(n)$?

The empty set, of size $n = 0$ has only one subset (itself), so $S(0) = 1$.

It is not difficult to see that

$$S(1) = 2, S(2) = 4, S(3) = 8,$$

Example 2:

Let $S(n)$ be the number of subsets of a set of size n .
What is the formula for $S(n)$?

The empty set, of size $n = 0$ has only one subset (itself), so $S(0) = 1$.

It is not difficult to see that
 $S(1) = 2$, $S(2) = 4$, $S(3) = 8$,

We “guess” that $S(n) = 2^n$ but, in order to prove formula, we’ll need to think recursively.

Example 2:

Let $S(n)$ be the number of subsets of a set of size n .
What is the formula for $S(n)$?

The empty set, of size $n = 0$ has only one subset (itself), so $S(0) = 1$.

It is not difficult to see that
 $S(1) = 2$, $S(2) = 4$, $S(3) = 8$,

We “guess” that $S(n) = 2^n$ but, in order to prove formula, we’ll need to think recursively.

Consider the eight subsets of $\{1, 2, 3\}$:
 $\emptyset, \{1\}, \{2\}, \{1, 2\}, \{3\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}$.

\emptyset	$\{1\}$	$\{2\}$	$\{1, 2\}$
$\{3\}$	$\{1, 3\}$	$\{2, 3\}$	$\{1, 2, 3\}$

$$\begin{array}{cccc}
 \emptyset & \{1\} & \{2\} & \{1, 2\} \\
 \{3\} & \{1, 3\} & \{2, 3\} & \{1, 2, 3\}
 \end{array}$$

First four subsets do not contain 3, but second four do.

$$\begin{array}{cccc}
 \emptyset & \{1\} & \{2\} & \{1, 2\} \\
 \{3\} & \{1, 3\} & \{2, 3\} & \{1, 2, 3\}
 \end{array}$$

First four subsets do not contain 3, but second four do.

First four subsets are exactly the subsets of $\{1, 2\}$, while second four are the subsets of $\{1, 2\}$ with 3 added into each one.

$$\begin{array}{cccc}
 \emptyset & \{1\} & \{2\} & \{1, 2\} \\
 \{\textcolor{red}{3}\} & \{1, \textcolor{red}{3}\} & \{2, \textcolor{red}{3}\} & \{1, 2, \textcolor{red}{3}\}
 \end{array}$$

First four subsets do not contain $\textcolor{red}{3}$, but second four do.

First four subsets are exactly the subsets of $\{1, 2\}$, while second four are the subsets of $\{1, 2\}$ with $\textcolor{red}{3}$ added into each one.

So, we get a subset of $\{1, 2, \textcolor{red}{3}\}$ either by taking a subset of $\{1, 2\}$ or by adjoining $\textcolor{red}{3}$ to a subset of $\{1, 2\}$.

$$\begin{array}{cccc}
 \emptyset & \{1\} & \{2\} & \{1, 2\} \\
 \{\textcolor{red}{3}\} & \{1, \textcolor{red}{3}\} & \{2, \textcolor{red}{3}\} & \{1, 2, \textcolor{red}{3}\}
 \end{array}$$

First four subsets do not contain $\textcolor{red}{3}$, but second four do.

First four subsets are exactly the subsets of $\{1, 2\}$, while second four are the subsets of $\{1, 2\}$ with $\textcolor{red}{3}$ added into each one.

So, we get a subset of $\{1, 2, \textcolor{red}{3}\}$ either by taking a subset of $\{1, 2\}$ or by adjoining $\textcolor{red}{3}$ to a subset of $\{1, 2\}$.

This suggests that the recurrence for the number of subsets of an n -element set $(\{1, 2, \dots, n\})$ is

$$S(n) = \begin{cases} 2S(n-1) & \text{if } n \geq 1, \\ 1 & \text{if } n = 0. \end{cases}$$

Proof (inductive) of correctness of this recurrence:

Proof (inductive) of correctness of this recurrence:

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain element n .

Proof (inductive) of correctness of this recurrence:

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain element n .

Each subset S containing n can be constructed in a unique fashion by adding n to the subset $S - \{n\}$ not containing n .

Proof (inductive) of correctness of this recurrence:

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain element n .

Each subset S containing n can be constructed in a unique fashion by adding n to the subset $S - \{n\}$ not containing n .

Each subset S not containing n can be constructed by removing n from the unique set $S \cup \{n\}$ containing n .

Proof (inductive) of correctness of this recurrence:

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain element n .

Each subset S containing n can be constructed in a unique fashion by adding n to the subset $S - \{n\}$ not containing n .

Each subset S not containing n can be constructed by removing n from the unique set $S \cup \{n\}$ containing n .

So, the number of subsets containing n is exactly the same as the number of subsets not containing n .

Proof (inductive) of correctness of this recurrence:

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain element n .

Each subset S containing n can be constructed in a unique fashion by adding n to the subset $S - \{n\}$ not containing n .

Each subset S not containing n can be constructed by removing n from the unique set $S \cup \{n\}$ containing n .

So, the number of subsets containing n is exactly the same as the the number of subsets not containing n .

The number of subsets not containing n is just the number of subsets of the $(n - 1)$ -element set $\{1, 2, \dots, n - 1\}$ which is $S(n - 1)$.

Proof (inductive) of correctness of this recurrence:

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain element n .

Each subset S containing n can be constructed in a unique fashion by adding n to the subset $S - \{n\}$ not containing n .

Each subset S not containing n can be constructed by removing n from the unique set $S \cup \{n\}$ containing n .

So, the number of subsets containing n is exactly the same as the the number of subsets not containing n .

The number of subsets not containing n is just the number of subsets of the $(n - 1)$ -element set $\{1, 2, \dots, n - 1\}$ which is $S(n - 1)$.

Thus, if $n > 0$, then $S(n) = 2S(n - 1)$.

Proof (inductive) of correctness of this recurrence:

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain element n .

Each subset S containing n can be constructed in a unique fashion by adding n to the subset $S - \{n\}$ not containing n .

Each subset S not containing n can be constructed by removing n from the unique set $S \cup \{n\}$ containing n .

So, the number of subsets containing n is exactly the same as the the number of subsets not containing n .

The number of subsets not containing n is just the number of subsets of the $(n - 1)$ -element set $\{1, 2, \dots, n - 1\}$ which is $S(n - 1)$.

Thus, if $n > 0$, then $S(n) = 2S(n - 1)$.

We already observed that \emptyset has only one subset (itself), so $S(0) = 1$ and we have proved the correctness of the recurrence.

If

$$S(n) = \begin{cases} 2S(n-1) & \text{if } n \geq 1, \\ 1 & \text{if } n = 0. \end{cases}$$

Then, $S(n) = 2^n$ for all $n \geq 0$.

If

$$S(n) = \begin{cases} 2S(n-1) & \text{if } n \geq 1, \\ 1 & \text{if } n = 0. \end{cases}$$

Then, $S(n) = 2^n$ for all $n \geq 0$.

Proof: by induction

If

$$S(n) = \begin{cases} 2S(n-1) & \text{if } n \geq 1, \\ 1 & \text{if } n = 0. \end{cases}$$

Then, $S(n) = 2^n$ for all $n \geq 0$.

Proof: by induction

i) if $n = 0$ then $S(0) = 2^0 = 1$.

If

$$S(n) = \begin{cases} 2S(n-1) & \text{if } n \geq 1, \\ 1 & \text{if } n = 0. \end{cases}$$

Then, $S(n) = 2^n$ for all $n \geq 0$.

Proof: by induction

i) if $n = 0$ then $S(0) = 2^0 = 1$.

ii) If the statement is true for $n - 1$ then $S(n - 1) = 2^{n-1}$ so

If

$$S(n) = \begin{cases} 2S(n-1) & \text{if } n \geq 1, \\ 1 & \text{if } n = 0. \end{cases}$$

Then, $S(n) = 2^n$ for all $n \geq 0$.

Proof: by induction

i) if $n = 0$ then $S(0) = 2^0 = 1$.

ii) If the statement is true for $n - 1$ then $S(n - 1) = 2^{n-1}$ so

$$S(n) = 2S(n-1) = 2 \cdot 2^{n-1} = 2^n$$

and we are done!

Example 3:

When paying off a loan with initial amount A and monthly payment M at an interest rate of p percent, the total amount $T(n)$ of the loan still due after n months is computed by adding $p/12$ percent to the amount due after $n - 1$ months and then subtracting the monthly payment M .

Convert this description into a recurrence for the amount owed after n months.

Example 3:

When paying off a loan with initial amount A and monthly payment M at an interest rate of p percent, the total amount $T(n)$ of the loan still due after n months is computed by adding $p/12$ percent to the amount due after $n - 1$ months and then subtracting the monthly payment M .

Convert this description into a recurrence for the amount owed after n months.

Answer

$$T(n) = \left(1 + \frac{0.01p}{12}\right) \cdot T(n - 1) - M, \text{ with } T(0) = A.$$

Example 3:

When paying off a loan with initial amount A and monthly payment M at an interest rate of p percent, the total amount $T(n)$ of the loan still due after n months is computed by adding $p/12$ percent to the amount due after $n - 1$ months and then subtracting the monthly payment M .

Convert this description into a recurrence for the amount owed after n months.

Answer

$$T(n) = \left(1 + \frac{0.01p}{12}\right) \cdot T(n - 1) - M, \text{ with } T(0) = A.$$

We will now see a general tool for deriving closed form solution to these type of recurrence relations

Recursion, Recurrences and Induction

- Recursion
- Recurrences
- Iterating a Recurrence
- Geometric Series
- First-Order Linear Recurrences

Iterating a Recurrence

Iterating a Recurrence

Let $T(n) = rT(n-1) + a$,
where r and a are constants.

Iterating a Recurrence

Let $T(n) = rT(n-1) + a$,
where r and a are constants.

Find a recurrence that expresses

$T(n)$ in terms of $T(n-2)$

$T(n)$ in terms of $T(n-3)$

$T(n)$ in terms of $T(n-4)$

⋮

Iterating a Recurrence

Let $T(n) = rT(n-1) + a$,
where r and a are constants.

Find a recurrence that expresses

$T(n)$ in terms of $T(n-2)$

$T(n)$ in terms of $T(n-3)$

$T(n)$ in terms of $T(n-4)$

⋮

Can you generalize this to find a closed form solution to

$$T(n) = rT(n-1) + a?$$

Note that $T(n) = rT(n-1) + a$, implies that,
 $\forall i < n, T(n-i) = rT((n-i)-1) + a$.

Note that $T(n) = rT(n-1) + a$, implies that,
 $\forall i < n, T(n-i) = rT((n-i)-1) + a$.

Then

$$T(n) = rT(n-1) + a$$

Note that $T(n) = rT(n-1) + a$, implies that,
 $\forall i < n, T(n-i) = rT((n-i)-1) + a$.

Then

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r(rT(n-2) + a) + a \end{aligned}$$

Note that $T(n) = rT(n-1) + a$, implies that,
 $\forall i < n, T(n-i) = rT((n-i)-1) + a$.

Then

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r(rT(n-2) + a) + a \\ &= r^2T(n-2) + ra + a \end{aligned}$$

Note that $T(n) = rT(n-1) + a$, implies that,
 $\forall i < n, T(n-i) = rT((n-i)-1) + a$.

Then

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r(rT(n-2) + a) + a \\ &= r^2T(n-2) + ra + a \\ &= r^2(rT(n-3) + a) + ra + a \end{aligned}$$

Note that $T(n) = rT(n-1) + a$, implies that,
 $\forall i < n, T(n-i) = rT((n-i)-1) + a$.

Then

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r(rT(n-2) + a) + a \\ &= r^2T(n-2) + ra + a \\ &= r^2(rT(n-3) + a) + ra + a \\ &= r^3T(n-3) + r^2a + ra + a \end{aligned}$$

Note that $T(n) = rT(n-1) + a$, implies that,

$$\forall i < n, T(n-i) = rT((n-i)-1) + a.$$

Then

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r(rT(n-2) + a) + a \\ &= r^2T(n-2) + ra + a \\ &= r^2(rT(n-3) + a) + ra + a \\ &= r^3T(n-3) + r^2a + ra + a \\ &= r^3(rT(n-4) + a) + r^2a + ra + a \end{aligned}$$

Note that $T(n) = rT(n-1) + a$, implies that,

$$\forall i < n, T(n-i) = rT((n-i)-1) + a.$$

Then

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r(rT(n-2) + a) + a \\ &= r^2T(n-2) + ra + a \\ &= r^2(rT(n-3) + a) + ra + a \\ &= r^3T(n-3) + r^2a + ra + a \\ &= r^3(rT(n-4) + a) + r^2a + ra + a \\ &= r^4T(n-4) + r^3a + r^2a + ra + a. \end{aligned}$$

Note that $T(n) = rT(n-1) + a$, implies that,
 $\forall i < n, T(n-i) = rT((n-i)-1) + a$.

Then

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r(rT(n-2) + a) + a \\ &= r^2T(n-2) + ra + a \\ &= r^2(rT(n-3) + a) + ra + a \\ &= r^3T(n-3) + r^2a + ra + a \\ &= r^3(rT(n-4) + a) + r^2a + ra + a \\ &= r^4T(n-4) + r^3a + r^2a + ra + a. \end{aligned}$$

From this, we can “guess” that

$$T(n) = r^n T(0) + a \sum_{i=0}^{n-1} r^i$$

Note that $T(n) = rT(n-1) + a$, implies that,

$$\forall i < n, T(n-i) = rT((n-i)-1) + a.$$

Then

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r(rT(n-2) + a) + a \\ &= r^2T(n-2) + ra + a \\ &= r^2(rT(n-3) + a) + ra + a \\ &= r^3T(n-3) + r^2a + ra + a \\ &= r^3(rT(n-4) + a) + r^2a + ra + a \\ &= r^4T(n-4) + r^3a + r^2a + ra + a. \end{aligned}$$

From this, we can “guess” that

$$T(n) = r^n T(0) + a \sum_{i=0}^{n-1} r^i = r^n b + a \sum_{i=0}^{n-1} r^i.$$

The method we used to guess the solution is called **iterating the recurrence**, because we repeatedly (iteratively) use the recurrence.

The method we used to guess the solution is called **iterating the recurrence**, because we repeatedly (iteratively) use the recurrence.

Another approach is to iterate from the “bottom-up” instead of “top-down”.

The method we used to guess the solution is called **iterating the recurrence**, because we repeatedly (iteratively) use the recurrence.

Another approach is to iterate from the “bottom-up” instead of “top-down”.

$$T(0) = b$$

The method we used to guess the solution is called **iterating the recurrence**, because we repeatedly (iteratively) use the recurrence.

Another approach is to iterate from the “bottom-up” instead of “top-down”.

$$T(0) = b$$

$$T(1) = rT(0) + a = rb + a$$

The method we used to guess the solution is called **iterating the recurrence**, because we repeatedly (iteratively) use the recurrence.

Another approach is to iterate from the “bottom-up” instead of “top-down”.

$$T(0) = b$$

$$T(1) = rT(0) + a = rb + a$$

$$T(2) = rT(1) + a = r(rb + a) + a = r^2b + ra + a$$

The method we used to guess the solution is called **iterating the recurrence**, because we repeatedly (iteratively) use the recurrence.

Another approach is to iterate from the “bottom-up” instead of “top-down”.

$$T(0) = b$$

$$T(1) = rT(0) + a = rb + a$$

$$T(2) = rT(1) + a = r(rb + a) + a = r^2b + ra + a$$

$$T(3) = rT(2) + a = r^3b + r^2a + ra + a$$

The method we used to guess the solution is called **iterating the recurrence**, because we repeatedly (iteratively) use the recurrence.

Another approach is to iterate from the “bottom-up” instead of “top-down”.

$$T(0) = b$$

$$T(1) = rT(0) + a = rb + a$$

$$T(2) = rT(1) + a = r(rb + a) + a = r^2b + ra + a$$

$$T(3) = rT(2) + a = r^3b + r^2a + ra + a$$

This could lead to the same guess

$$T(n) = r^n b + a \sum_{i=0}^{n-1} r^i.$$

Recursion, Recurrences and Induction

- Recursion
- Recurrences
- Iterating a Recurrence
- Geometric Series
- First-Order Linear Recurrences

Geometric Series

Geometric Series

$\sum_{i=0}^{n-1} r^i$ is a finite geometric series with common ratio r .

$\sum_{i=0}^{n-1} ar^i$ is a finite geometric series with common ratio r
and initial value a .

Geometric Series

$\sum_{i=0}^{n-1} r^i$ is a finite geometric series with common ratio r .

$\sum_{i=0}^{n-1} ar^i$ is a finite geometric series with common ratio r
and initial value a .

Geometric Series

$\sum_{i=0}^{n-1} r^i$ is a finite geometric series with common ratio r .

$\sum_{i=0}^{n-1} ar^i$ is a finite geometric series with common ratio r and initial value a .

It is known that, for all $r \neq 1$,

$$\sum_{i=0}^{n-1} r^i = \frac{1 - r^n}{1 - r}.$$

Geometric Series

$\sum_{i=0}^{n-1} r^i$ is a finite geometric series with common ratio r .

$\sum_{i=0}^{n-1} ar^i$ is a finite geometric series with common ratio r and initial value a .

It is known that, for all $r \neq 1$,

$$\sum_{i=0}^{n-1} r^i = \frac{1 - r^n}{1 - r}.$$

Note: We will see another proof of this soon.

Theorem 4.1

If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Theorem 4.1

If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Proof by induction:

Theorem 4.1

If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Proof by induction:

$$T(0) = r^0 b + a \frac{1 - r^0}{1 - r} = b.$$

Theorem 4.1

If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Proof by induction:

$$T(0) = r^0 b + a \frac{1 - r^0}{1 - r} = b.$$

So, the formula is true when $n = 0$.

Theorem 4.1

If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Proof by induction:

$$T(0) = r^0 b + a \frac{1 - r^0}{1 - r} = b.$$

So, the formula is true when $n = 0$.

Now assume that $n > 0$ and

$$T(n-1) = r^{n-1} b + a \frac{1 - r^{n-1}}{1 - r}.$$

Then we have $T(n) = rT(n-1) + a$

Then we have $T(n) = rT(n-1) + a$

$$= r \left(r^{n-1}b + a \frac{1 - r^{n-1}}{1 - r} \right) + a$$

Then we have

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r \left(r^{n-1}b + a \frac{1-r^{n-1}}{1-r} \right) + a \\ &= r^n b + \frac{ar - ar^n}{1-r} + a \end{aligned}$$

Then we have

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r \left(r^{n-1}b + a \frac{1-r^{n-1}}{1-r} \right) + a \\ &= r^n b + \frac{ar - ar^n}{1-r} + a \\ &= r^n b + \frac{ar - ar^n + a - ar}{1-r} \end{aligned}$$

Then we have

$$\begin{aligned}
 T(n) &= rT(n-1) + a \\
 &= r \left(r^{n-1}b + a \frac{1-r^{n-1}}{1-r} \right) + a \\
 &= r^n b + \frac{ar - ar^n}{1-r} + a \\
 &= r^n b + \frac{ar - ar^n + a - ar}{1-r} \\
 &= r^n b + a \frac{1-r^n}{1-r}.
 \end{aligned}$$

Then we have

$$\begin{aligned}
 T(n) &= rT(n-1) + a \\
 &= r \left(r^{n-1}b + a \frac{1-r^{n-1}}{1-r} \right) + a \\
 &= r^n b + \frac{ar - ar^n}{1-r} + a \\
 &= r^n b + \frac{ar - ar^n + a - ar}{1-r} \\
 &= r^n b + a \frac{1-r^n}{1-r}.
 \end{aligned}$$

Therefore, by the principle of mathematical induction, our formula holds for all integers $n \geq 0$.

Theorem 4.1

If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Theorem 4.1

If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Example:

Theorem 4.1

If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Example:

$$T(n) = 3T(n-1) + 2 \quad \text{with} \quad T(0) = 5$$

Theorem 4.1

If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Example:

$$T(n) = 3T(n-1) + 2 \quad \text{with} \quad T(0) = 5$$

Plugging $r = 3$, $a = 2$, $b = 5$ into the formula, gives

$$T(n) = 3^n \cdot 5 + 2 \frac{1 - 3^n}{1 - 3} = 3^n \cdot 6 - 1$$

Corollary 4.2: The formula for the sum of a geometric series with $r \neq 1$ is

$$\sum_{i=0}^{n-1} r^i = \frac{1 - r^n}{1 - r}$$

Corollary 4.2: The formula for the sum of a geometric series with $r \neq 1$ is

$$\sum_{i=0}^{n-1} r^i = \frac{1 - r^n}{1 - r}$$

Proof:

Let $T(0) = 0$, and $T(n) = \sum_{i=0}^{n-1} r^i$ for $n > 0$.

Corollary 4.2: The formula for the sum of a geometric series with $r \neq 1$ is

$$\sum_{i=0}^{n-1} r^i = \frac{1 - r^n}{1 - r}$$

Proof:

Let $T(0) = 0$, and $T(n) = \sum_{i=0}^{n-1} r^i$ for $n > 0$.

Then $T(n) = rT(n-1) + 1$.

Applying Theorem 4.1 with $b = 0$ and $a = 1$ gives

$$T(n) = \frac{1 - r^n}{1 - r}$$

Lemma 4.3: Let $r \neq 1$ be a positive value independent of n .
Let $t(n)$ be the largest term in the geometric series

$$\sum_{i=0}^{n-1} r^i$$

Then the value of the geometric series is $O(t(n))$.

Lemma 4.3: Let $r \neq 1$ be a positive value independent of n . Let $t(n)$ be the largest term in the geometric series

$$\sum_{i=0}^{n-1} r^i$$

Then the value of the geometric series is $O(t(n))$.

Proof: There are two cases.

- i) $r < 1$: in which case, $t(n) = r^0 = 1$.
- ii) $r > 1$: in which case $t(n) = r^{n-1}$

Lemma 4.3: Let $r \neq 1$ be a positive value independent of n . Let $t(n)$ be the largest term in the geometric series

$$\sum_{i=0}^{n-1} r^i$$

Then the value of the geometric series is $O(t(n))$.

Proof: There are two cases.

- i) $r < 1$: in which case, $t(n) = r^0 = 1$.
- ii) $r > 1$: in which case $t(n) = r^{n-1}$

(i) is easy because in this case

$$\sum_{i=0}^{n-1} r^i = \frac{1 - r^n}{1 - r} < \frac{1}{1 - r}$$

Lemma 4.3: Let $r \neq 1$ be a positive value independent of n . Let $t(n)$ be the largest term in the geometric series

$$\sum_{i=0}^{n-1} r^i$$

Then the value of the geometric series is $O(t(n))$.

Proof: There are two cases.

i) $r < 1$: in which case, $t(n) = r^0 = 1$.

ii) $r > 1$: in which case $t(n) = r^{n-1}$

(i) is easy because in this case

$$\sum_{i=0}^{n-1} r^i = \frac{1 - r^n}{1 - r} < \frac{1}{1 - r} \quad \text{which is } O(1) = O(t(n))$$

In case (ii), $r > 1$, $t(n) = r^{n-1}$ and

$$\sum_{i=0}^{n-1} r^i = \frac{1 - r^n}{1 - r} = \frac{r^n - 1}{r - 1} < \frac{r^n}{r - 1} = r^{n-1} \frac{r}{r - 1}$$

In case (ii), $r > 1$, $t(n) = r^{n-1}$ and

$$\sum_{i=0}^{n-1} r^i = \frac{1 - r^n}{1 - r} = \frac{r^n - 1}{r - 1} < \frac{r^n}{r - 1} = r^{n-1} \frac{r}{r - 1}$$

Thus, $\sum_{i=0}^{n-1} r^i = O(r^{n-1}) = O(t(n))$.

Recursion, Recurrences and Induction

- Recursion
- Recurrences
- Iterating a Recurrence
- Geometric Series
- First-Order Linear Recurrences

First-Order Linear Recurrences

First-Order Linear Recurrences

A recurrence of the form $T(n) = f(n)T(n-1) + g(n)$ is called a **first-order linear recurrence**.

First-Order Linear Recurrences

A recurrence of the form $T(n) = f(n)T(n-1) + g(n)$ is called a **first-order linear recurrence**.

- **First Order** because it is only dependent upon going back one step, i.e., $T(n-1)$.

First-Order Linear Recurrences

A recurrence of the form $T(n) = f(n)T(n-1) + g(n)$ is called a **first-order linear recurrence**.

- **First Order** because it is only dependent upon going back one step, i.e., $T(n-1)$.

If it was dependent upon $T(n-2)$, it would be a **second-order** recurrence, e.g., $T(n) = T(n-1) + 2T(n-2)$.

First-Order Linear Recurrences

A recurrence of the form $T(n) = f(n)T(n-1) + g(n)$ is called a **first-order linear recurrence**.

- **First Order** because it is only dependent upon going back one step, i.e., $T(n-1)$.

If it was dependent upon $T(n-2)$, it would be a **second-order** recurrence, e.g., $T(n) = T(n-1) + 2T(n-2)$.

- **Linear** because $T(n-1)$ only appears to the first power.

First-Order Linear Recurrences

A recurrence of the form $T(n) = f(n)T(n-1) + g(n)$ is called a **first-order linear recurrence**.

- **First Order** because it is only dependent upon going back one step, i.e., $T(n-1)$.

If it was dependent upon $T(n-2)$, it would be a **second-order** recurrence, e.g., $T(n) = T(n-1) + 2T(n-2)$.

- **Linear** because $T(n-1)$ only appears to the first power.

Something like $T(n) = (T(n-1))^2 + 3$ would be a **non-linear** first-order recurrence relation.

$$T(n) = f(n)T(n-1) + g(n)$$

When $f(n)$ is a constant, say r , the general solution is almost as easy to write as in Theorem 4.1.

$$T(n) = f(n)T(n-1) + g(n)$$

When $f(n)$ is a constant, say r , the general solution is almost as easy to write as in Theorem 4.1.

Iterating the recurrence gives

$$T(n) = rT(n-1) + g(n)$$

$$T(n) = f(n)T(n-1) + g(n)$$

When $f(n)$ is a constant, say r , the general solution is almost as easy to write as in Theorem 4.1.

Iterating the recurrence gives

$$\begin{aligned} T(n) &= rT(n-1) + g(n) \\ &= r(rT(n-2) + g(n-1)) + g(n) \\ &= r^2T(n-2) + rg(n-1) + g(n) \\ &= r^3T(n-3) + r^2g(n-2) + rg(n-1) + g(n) \\ &\vdots \end{aligned}$$

$$T(n) = f(n)T(n-1) + g(n)$$

When $f(n)$ is a constant, say r , the general solution is almost as easy to write as in Theorem 4.1.

Iterating the recurrence gives

$$\begin{aligned} T(n) &= rT(n-1) + g(n) \\ &= r(rT(n-2) + g(n-1)) + g(n) \\ &= r^2T(n-2) + rg(n-1) + g(n) \\ &= r^3T(n-3) + r^2g(n-2) + rg(n-1) + g(n) \\ &\vdots \\ &= r^nT(0) + \sum_{i=0}^{n-1} r^i g(n-i) \end{aligned}$$

Theorem 4.5 For any positive constants a and r , and any function g defined on the nonnegative integers, the solution to the first-order linear recurrence

$$T(n) = \begin{cases} rT(n-1) + g(n) & \text{if } n > 0, \\ a & \text{if } n = 0, \end{cases}$$

is

$$T(n) = r^n a + \sum_{i=1}^n r^{n-i} g(i). \quad (*)$$

Theorem 4.5 For any positive constants a and r , and any function g defined on the nonnegative integers, the solution to the first-order linear recurrence

$$T(n) = \begin{cases} rT(n-1) + g(n) & \text{if } n > 0, \\ a & \text{if } n = 0, \end{cases}$$

is

$$T(n) = r^n a + \sum_{i=1}^n r^{n-i} g(i). \quad (*)$$

Proof by induction:

Theorem 4.5 For any positive constants a and r , and any function g defined on the nonnegative integers, the solution to the first-order linear recurrence

$$T(n) = \begin{cases} rT(n-1) + g(n) & \text{if } n > 0, \\ a & \text{if } n = 0, \end{cases}$$

is

$$T(n) = r^n a + \sum_{i=1}^n r^{n-i} g(i). \quad (*)$$

Proof by induction:

Because the sum $(*)$ has no terms when $n = 0$, the formula gives $T(0) = a$ and, so, is valid when $n = 0$.

Theorem 4.5 For any positive constants a and r , and any function g defined on the nonnegative integers, the solution to the first-order linear recurrence

$$T(n) = \begin{cases} rT(n-1) + g(n) & \text{if } n > 0, \\ a & \text{if } n = 0, \end{cases}$$

is

$$T(n) = r^n a + \sum_{i=1}^n r^{n-i} g(i). \quad (*)$$

Proof by induction:

Because the sum $(*)$ has no terms when $n = 0$, the formula gives $T(0) = a$ and, so, is valid when $n = 0$.

We now assume that n is positive and

$$T(n-1) = r^{n-1} a + \sum_{i=1}^{n-1} r^{(n-1)-i} g(i).$$

Using the definition of the recurrence and the inductive hypothesis, we get that

$$T(n) = rT(n-1) + g(n)$$

Using the definition of the recurrence and the inductive hypothesis, we get that

$$\begin{aligned} T(n) &= rT(n-1) + g(n) \\ &= r \left(r^{n-1}a + \sum_{i=1}^{n-1} r^{(n-1)-i} g(i) \right) + g(n) \end{aligned}$$

Using the definition of the recurrence and the inductive hypothesis, we get that

$$\begin{aligned} T(n) &= rT(n-1) + g(n) \\ &= r \left(r^{n-1}a + \sum_{i=1}^{n-1} r^{(n-1)-i} g(i) \right) + g(n) \\ &= r^n a + \sum_{i=1}^{n-1} r^{(n-1)+1-i} g(i) + g(n) \end{aligned}$$

Using the definition of the recurrence and the inductive hypothesis, we get that

$$\begin{aligned}T(n) &= rT(n-1) + g(n) \\&= r \left(r^{n-1}a + \sum_{i=1}^{n-1} r^{(n-1)-i} g(i) \right) + g(n) \\&= r^n a + \sum_{i=1}^{n-1} r^{(n-1)+1-i} g(i) + g(n) \\&= r^n a + \sum_{i=1}^{n-1} r^{n-i} g(i) + g(n)\end{aligned}$$

Using the definition of the recurrence and the inductive hypothesis, we get that

$$\begin{aligned}T(n) &= rT(n-1) + g(n) \\&= r \left(r^{n-1}a + \sum_{i=1}^{n-1} r^{(n-1)-i} g(i) \right) + g(n) \\&= r^n a + \sum_{i=1}^{n-1} r^{(n-1)+1-i} g(i) + g(n) \\&= r^n a + \sum_{i=1}^{n-1} r^{n-i} g(i) + g(n) \\&= r^n a + \sum_{i=1}^n r^{n-i} g(i).\end{aligned}$$

Using the definition of the recurrence and the inductive hypothesis, we get that

$$\begin{aligned}T(n) &= rT(n-1) + g(n) \\&= r \left(r^{n-1}a + \sum_{i=1}^{n-1} r^{(n-1)-i} g(i) \right) + g(n) \\&= r^n a + \sum_{i=1}^{n-1} r^{(n-1)+1-i} g(i) + g(n) \\&= r^n a + \sum_{i=1}^{n-1} r^{n-i} g(i) + g(n) \\&= r^n a + \sum_{i=1}^n r^{n-i} g(i).\end{aligned}$$

Therefore, by the principle of mathematical induction, the solution to the recurrence is given by (*) for all nonnegative integers n .

Example: Solve $T(n) = 4T(n - 1) + 2^n$ with $T(0) = 6$.

Example: Solve $T(n) = 4T(n - 1) + 2^n$ with $T(0) = 6$.

Using Theorem 4.5

$$T(n) = 6 \cdot 4^n + \sum_{i=1}^n 4^{n-i} \cdot 2^i$$

Example: Solve $T(n) = 4T(n - 1) + 2^n$ with $T(0) = 6$.

Using Theorem 4.5

$$\begin{aligned} T(n) &= 6 \cdot 4^n + \sum_{i=1}^n 4^{n-i} \cdot 2^i \\ &= 6 \cdot 4^n + 4^n \sum_{i=1}^n 4^{-i} \cdot 2^i \end{aligned}$$

Example: Solve $T(n) = 4T(n-1) + 2^n$ with $T(0) = 6$.

Using Theorem 4.5

$$\begin{aligned} T(n) &= 6 \cdot 4^n + \sum_{i=1}^n 4^{n-i} \cdot 2^i \\ &= 6 \cdot 4^n + 4^n \sum_{i=1}^n 4^{-i} \cdot 2^i \\ &= 6 \cdot 4^n + 4^n \sum_{i=1}^n \left(\frac{1}{2}\right)^i \end{aligned}$$

Example: Solve $T(n) = 4T(n-1) + 2^n$ with $T(0) = 6$.

Using Theorem 4.5

$$\begin{aligned} T(n) &= 6 \cdot 4^n + \sum_{i=1}^n 4^{n-i} \cdot 2^i \\ &= 6 \cdot 4^n + 4^n \sum_{i=1}^n 4^{-i} \cdot 2^i \\ &= 6 \cdot 4^n + 4^n \sum_{i=1}^n \left(\frac{1}{2}\right)^i \\ &= 6 \cdot 4^n + 4^n \cdot \frac{1}{2} \cdot \sum_{i=0}^{n-1} \left(\frac{1}{2}\right)^i \end{aligned}$$

Example: Solve $T(n) = 4T(n-1) + 2^n$ with $T(0) = 6$.

Using Theorem 4.5

$$\begin{aligned}T(n) &= 6 \cdot 4^n + \sum_{i=1}^n 4^{n-i} \cdot 2^i \\&= 6 \cdot 4^n + 4^n \sum_{i=1}^n 4^{-i} \cdot 2^i \\&= 6 \cdot 4^n + 4^n \sum_{i=1}^n \left(\frac{1}{2}\right)^i \\&= 6 \cdot 4^n + 4^n \cdot \frac{1}{2} \cdot \sum_{i=0}^{n-1} \left(\frac{1}{2}\right)^i \\&= 6 \cdot 4^n + \left(1 - \left(\frac{1}{2}\right)^n\right) \cdot 4^n\end{aligned}$$

Example: Solve $T(n) = 4T(n-1) + 2^n$ with $T(0) = 6$.

Using Theorem 4.5

$$\begin{aligned}T(n) &= 6 \cdot 4^n + \sum_{i=1}^n 4^{n-i} \cdot 2^i \\&= 6 \cdot 4^n + 4^n \sum_{i=1}^n 4^{-i} \cdot 2^i \\&= 6 \cdot 4^n + 4^n \sum_{i=1}^n \left(\frac{1}{2}\right)^i \\&= 6 \cdot 4^n + 4^n \cdot \frac{1}{2} \cdot \sum_{i=0}^{n-1} \left(\frac{1}{2}\right)^i \\&= 6 \cdot 4^n + \left(1 - \left(\frac{1}{2}\right)^n\right) \cdot 4^n \\&= 7 \cdot 4^n - 2^n.\end{aligned}$$

Example: Solve $T(n) = 3T(n - 1) + n$ with $T(0) = 10$.

Using Theorem 4.5

$$\begin{aligned} T(n) &= 10 \cdot 3^n + \sum_{i=1}^n 3^{n-i} \cdot i \\ &= 10 \cdot 3^n + 3^n \sum_{i=1}^n i \cdot 3^{-i} \end{aligned}$$

We now need the following well known theorem
(can be proven by induction or see book for another proof)

Theorem 4.6

For any real number $x \neq 1$,

$$\sum_{i=1}^n ix^i = \frac{nx^{n+2} - (n+1)x^{n+1} + x}{(1-x)^2}.$$

Example: Solve $T(n) = 3T(n - 1) + n$ with $T(0) = 10$.

Using Theorem 4.5

$$\begin{aligned} T(n) &= 10 \cdot 3^n + \sum_{i=1}^n 3^{n-i} \cdot i \\ &= 10 \cdot 3^n + 3^n \sum_{i=1}^n i \cdot 3^{-i} \end{aligned}$$

Example: Solve $T(n) = 3T(n - 1) + n$ with $T(0) = 10$.

Using Theorem 4.5

$$\begin{aligned} T(n) &= 10 \cdot 3^n + \sum_{i=1}^n 3^{n-i} \cdot i \\ &= 10 \cdot 3^n + 3^n \sum_{i=1}^n i \cdot 3^{-i} \\ &= 10 \cdot 3^n + 3^n \left(-\frac{3}{2}(n+1)3^{-(n+1)} - \frac{3}{4}3^{-(n+1)} + \frac{3}{4} \right) \end{aligned}$$

Example: Solve $T(n) = 3T(n - 1) + n$ with $T(0) = 10$.

Using Theorem 4.5

$$\begin{aligned}T(n) &= 10 \cdot 3^n + \sum_{i=1}^n 3^{n-i} \cdot i \\&= 10 \cdot 3^n + 3^n \sum_{i=1}^n i \cdot 3^{-i} \\&= 10 \cdot 3^n + 3^n \left(-\frac{3}{2}(n+1)3^{-(n+1)} - \frac{3}{4}3^{-(n+1)} + \frac{3}{4} \right) \\&= \frac{43}{4}3^n - \frac{n+1}{2} - \frac{1}{4}\end{aligned}$$