

Language

Definition

A *language* over alphabet A is a set $L \subseteq A^*$. Example for $A = \{0,1\}$:

- ▶ a finite language like $L = \{1, 10, 1001\}$ or the empty language \emptyset
- ▶ infinite but very difficult to describe (there are random languages: there exist more languages as subsets of A^* than there are finite descriptions)
- ▶ infinite but having some nice structure, where words follow a certain “pattern” that we can describe precisely and check efficiently \leftarrow these are our focus

$L_2 = \{01, 0101, 010101, \dots\}$ = those non-empty words that are of the form $01\dots 01$ where the block 01 is repeated some finite positive number of times. Using notation $(01)^n$ for a word consisting of block 01 repeated n times, we can write $L_2 = \{(01)^n \mid n \geq 1\}$.

Languages are sets, so we can take their union (\cup), intersection (\cap), and apply other set operations on languages.

Languages \emptyset and $\{\varepsilon\}$ are very different: \emptyset is a set that contains no words, whereas $\{\varepsilon\}$ contains precisely one word, the word of length zero.

More Language Operations

In addition to set operations, languages support operations defined on their words.

Definition (Language concatenation)

Given $L_1 \subseteq A^*$ and $L_2 \subseteq A^*$, define $L_1 \cdot L_2 = \{w_1 \cdot w_2 \mid w_1 \in L_1, w_2 \in L_2\}$

Example: $\{\varepsilon, a, aa\} \cdot \{b, bb\} = \{b, bb, ab, abb, aab, aabb\}$

In other words, $w \in L_1 L_2$ iff w can be split into $w_1 \in L_1$ and $w_2 \in L_2$ s.t. $w = w_1 w_2$.

More Language Operations

In addition to set operations, languages support operations defined on their words.

Definition (Language concatenation)

Given $L_1 \subseteq A^*$ and $L_2 \subseteq A^*$, define $L_1 \cdot L_2 = \{w_1 \cdot w_2 \mid w_1 \in L_1, w_2 \in L_2\}$

Example: $\{\epsilon, a, aa\} \cdot \{b, bb\} = \{b, bb, ab, abb, aab, aabb\}$

In other words, $w \in L_1 L_2$ iff w can be split into $w_1 \in L_1$ and $w_2 \in L_2$ s.t. $w = w_1 w_2$.

Definition (Language exponentiation)

Given $L \subseteq A^*$, define

$$\begin{aligned} L^0 &= \{\epsilon\} \\ L^{n+1} &= L \cdot L^n \end{aligned}$$

More Language Operations

In addition to set operations, languages support operations defined on their words.

Definition (Language concatenation)

Given $L_1 \subseteq A^*$ and $L_2 \subseteq A^*$, define $L_1 \cdot L_2 = \{w_1 \cdot w_2 \mid w_1 \in L_1, w_2 \in L_2\}$

Example: $\{\varepsilon, a, aa\} \cdot \{b, bb\} = \{b, bb, ab, abb, aab, aabb\}$

In other words, $w \in L_1 L_2$ iff w can be split into $w_1 \in L_1$ and $w_2 \in L_2$ s.t. $w = w_1 w_2$.

Definition (Language exponentiation)

Given $L \subseteq A^*$, define

$$\begin{aligned} L^0 &= \{\varepsilon\} \\ L^{n+1} &= L \cdot L^n \end{aligned}$$

Theorem

Given $L \subseteq A^*$, we have $L^n = \{w_1 \dots w_n \mid w_1, \dots, w_n \in L\}$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

► $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\}$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\}$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\}$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\}$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\} = \{w_1 \varepsilon \mid w_1 \in L\}$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\} = \{w_1 \varepsilon \mid w_1 \in L\} = L$
- ▶ $\{\varepsilon\}L$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\} = \{w_1 \varepsilon \mid w_1 \in L\} = L$
- ▶ $\{\varepsilon\}L = L$
- ▶ $\emptyset\{\varepsilon\}$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\} = \{w_1 \varepsilon \mid w_1 \in L\} = L$
- ▶ $\{\varepsilon\}L = L$
- ▶ $\emptyset\{\varepsilon\} = \emptyset$
- ▶ Is LL equal to $\{w^2 \mid w \in L\}$?

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\} = \{w_1 \varepsilon \mid w_1 \in L\} = L$
- ▶ $\{\varepsilon\}L = L$
- ▶ $\emptyset\{\varepsilon\} = \emptyset$
- ▶ Is LL equal to $\{w^2 \mid w \in L\}$? No! The latter only has words of the shape ww .
- ▶ $\{\varepsilon\}^n$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\} = \{w_1 \varepsilon \mid w_1 \in L\} = L$
- ▶ $\{\varepsilon\}L = L$
- ▶ $\emptyset\{\varepsilon\} = \emptyset$
- ▶ Is LL equal to $\{w^2 \mid w \in L\}$? No! The latter only has words of the shape ww .
- ▶ $\{\varepsilon\}^n = \{w_1 \dots w_n \mid w_1 \in \{\varepsilon\}, \dots, w_n \in \{\varepsilon\}\}$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\} = \{w_1 \varepsilon \mid w_1 \in L\} = L$
- ▶ $\{\varepsilon\}L = L$
- ▶ $\emptyset\{\varepsilon\} = \emptyset$
- ▶ Is LL equal to $\{w^2 \mid w \in L\}$? No! The latter only has words of the shape ww .
- ▶ $\{\varepsilon\}^n = \{w_1 \dots w_n \mid w_1 \in \{\varepsilon\}, \dots, w_n \in \{\varepsilon\}\} = \{\varepsilon^n\}$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\} = \{w_1 \varepsilon \mid w_1 \in L\} = L$
- ▶ $\{\varepsilon\}L = L$
- ▶ $\emptyset\{\varepsilon\} = \emptyset$
- ▶ Is LL equal to $\{w^2 \mid w \in L\}$? No! The latter only has words of the shape ww .
- ▶ $\{\varepsilon\}^n = \{w_1 \dots w_n \mid w_1 \in \{\varepsilon\}, \dots, w_n \in \{\varepsilon\}\} = \{\varepsilon^n\} = \{\varepsilon\}$

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\} = \{w_1 \varepsilon \mid w_1 \in L\} = L$
- ▶ $\{\varepsilon\}L = L$
- ▶ $\emptyset\{\varepsilon\} = \emptyset$
- ▶ Is LL equal to $\{w^2 \mid w \in L\}$? No! The latter only has words of the shape ww .
- ▶ $\{\varepsilon\}^n = \{w_1 \dots w_n \mid w_1 \in \{\varepsilon\}, \dots, w_n \in \{\varepsilon\}\} = \{\varepsilon^n\} = \{\varepsilon\}$

Note the difference between:

- ▶ the empty language \emptyset , which contains no words
- ▶ the language $\{\varepsilon\}$, which contains exactly one word, ε

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\} = \{w_1 \varepsilon \mid w_1 \in L\} = L$
- ▶ $\{\varepsilon\}L = L$
- ▶ $\emptyset\{\varepsilon\} = \emptyset$
- ▶ Is LL equal to $\{w^2 \mid w \in L\}$? No! The latter only has words of the shape ww .
- ▶ $\{\varepsilon\}^n = \{w_1 \dots w_n \mid w_1 \in \{\varepsilon\}, \dots, w_n \in \{\varepsilon\}\} = \{\varepsilon^n\} = \{\varepsilon\}$

Note the difference between:

- ▶ the empty language \emptyset , which contains no words
- ▶ the language $\{\varepsilon\}$, which contains exactly one word, ε

Q: for any L_1 and L_2 , does $L_1 L_2 = L_2 L_1$ hold? Can it possibly hold?

Expanding the Definition

If L is an arbitrary language, compute each of the following:

- ▶ $L\emptyset = \{w_1 w_2 \mid w_1 \in L, w_2 \in \emptyset\} = \emptyset$
- ▶ $\emptyset L = \emptyset$
- ▶ $L\{\varepsilon\} = \{w_1 w_2 \mid w_1 \in L, w_2 \in \{\varepsilon\}\} = \{w_1 w_2 \mid w_1 \in L, w_2 = \varepsilon\} = \{w_1 \varepsilon \mid w_1 \in L\} = L$
- ▶ $\{\varepsilon\}L = L$
- ▶ $\emptyset\{\varepsilon\} = \emptyset$
- ▶ Is LL equal to $\{w^2 \mid w \in L\}$? No! The latter only has words of the shape ww .
- ▶ $\{\varepsilon\}^n = \{w_1 \dots w_n \mid w_1 \in \{\varepsilon\}, \dots, w_n \in \{\varepsilon\}\} = \{\varepsilon^n\} = \{\varepsilon\}$

Note the difference between:

- ▶ the empty language \emptyset , which contains no words
- ▶ the language $\{\varepsilon\}$, which contains exactly one word, ε

Q: for any L_1 and L_2 , does $L_1 L_2 = L_2 L_1$ hold? Can it possibly hold?

Answer: no (ex: $L_1 = \{a\}$, $L_2 = \{b\}$) and yes (ex: $L_1, L_2 \in \{\emptyset, \{\varepsilon\}\}$, $L_1 = L_2$, ...).

Concatenation of Languages

Let A be alphabet. Consider the set of all languages $L \subseteq A^*$

Is this a monoid?

Concatenation of Languages

Let A be alphabet. Consider the set of all languages $L \subseteq A^*$

Is this a monoid?

- ▶ Is there a neutral element?

Concatenation of Languages

Let A be alphabet. Consider the set of all languages $L \subseteq A^*$

Is this a monoid?

- ▶ Is there a neutral element? $L\{\varepsilon\} = L$, $\{\varepsilon\}L = L$, so $\{\varepsilon\}$ is one
- ▶ Which law needs to hold? Does it hold?

Concatenation of Languages

Let A be alphabet. Consider the set of all languages $L \subseteq A^*$

Is this a monoid?

- ▶ Is there a neutral element? $L\{\varepsilon\} = L$, $\{\varepsilon\}L = L$, so $\{\varepsilon\}$ is one
- ▶ Which law needs to hold? Does it hold?

Associativity: we need $L_1 \cdot (L_2 \cdot L_3) = (L_1 \cdot L_2) \cdot L_3$

Concatenation of Languages

Let A be alphabet. Consider the set of all languages $L \subseteq A^*$

Is this a monoid?

- ▶ Is there a neutral element? $L\{\varepsilon\} = L$, $\{\varepsilon\}L = L$, so $\{\varepsilon\}$ is one
- ▶ Which law needs to hold? Does it hold?

Associativity: we need $L_1 \cdot (L_2 \cdot L_3) = (L_1 \cdot L_2) \cdot L_3$

$$L_1 \cdot (L_2 \cdot L_3) = \{w_1(w_2w_3) \mid w_1 \in L_1 \wedge (w_2 \in L_2 \wedge w_3 \in L_3)\}$$

Concatenation of Languages

Let A be alphabet. Consider the set of all languages $L \subseteq A^*$

Is this a monoid?

- ▶ Is there a neutral element? $L\{\varepsilon\} = L$, $\{\varepsilon\}L = L$, so $\{\varepsilon\}$ is one
- ▶ Which law needs to hold? Does it hold?

Associativity: we need $L_1 \cdot (L_2 \cdot L_3) = (L_1 \cdot L_2) \cdot L_3$

$$L_1 \cdot (L_2 \cdot L_3) = \{w_1(w_2w_3) \mid w_1 \in L_1 \wedge (w_2 \in L_2 \wedge w_3 \in L_3)\}$$

$$(L_1 \cdot L_2) \cdot L_3 = \{(w_1w_2)w_3 \mid (w_1 \in L_1 \wedge w_2) \in L_2 \wedge w_3 \in L_3\}$$

Concatenation of Languages

Let A be alphabet. Consider the set of all languages $L \subseteq A^*$

Is this a monoid?

- ▶ Is there a neutral element? $L\{\varepsilon\} = L$, $\{\varepsilon\}L = L$, so $\{\varepsilon\}$ is one
- ▶ Which law needs to hold? Does it hold?

Associativity: we need $L_1 \cdot (L_2 \cdot L_3) = (L_1 \cdot L_2) \cdot L_3$

$$L_1 \cdot (L_2 \cdot L_3) = \{w_1(w_2w_3) \mid w_1 \in L_1 \wedge (w_2 \in L_2 \wedge w_3 \in L_3)\}$$

$$(L_1 \cdot L_2) \cdot L_3 = \{(w_1w_2)w_3 \mid (w_1 \in L_1 \wedge w_2) \in L_2 \wedge w_3 \in L_3\}$$

It is a monoid.

Concatenation of Languages

Let A be alphabet. Consider the set of all languages $L \subseteq A^*$

Is this a monoid?

- ▶ Is there a neutral element? $L\{\varepsilon\} = L$, $\{\varepsilon\}L = L$, so $\{\varepsilon\}$ is one
- ▶ Which law needs to hold? Does it hold?

Associativity: we need $L_1 \cdot (L_2 \cdot L_3) = (L_1 \cdot L_2) \cdot L_3$

$$L_1 \cdot (L_2 \cdot L_3) = \{w_1(w_2w_3) \mid w_1 \in L_1 \wedge (w_2 \in L_2 \wedge w_3 \in L_3)\}$$

$$(L_1 \cdot L_2) \cdot L_3 = \{(w_1w_2)w_3 \mid (w_1 \in L_1 \wedge w_2) \in L_2 \wedge w_3 \in L_3\}$$

It is a monoid.

Concatenation of Languages

Let A be alphabet. Consider the set of all languages $L \subseteq A^*$

Is this a monoid?

- ▶ Is there a neutral element? $L\{\varepsilon\} = L$, $\{\varepsilon\}L = L$, so $\{\varepsilon\}$ is one
- ▶ Which law needs to hold? Does it hold?

Associativity: we need $L_1 \cdot (L_2 \cdot L_3) = (L_1 \cdot L_2) \cdot L_3$

$$L_1 \cdot (L_2 \cdot L_3) = \{w_1(w_2w_3) \mid w_1 \in L_1 \wedge (w_2 \in L_2 \wedge w_3 \in L_3)\}$$

$$(L_1 \cdot L_2) \cdot L_3 = \{(w_1w_2)w_3 \mid (w_1 \in L_1 \wedge w_2) \in L_2 \wedge w_3 \in L_3\}$$

It is a monoid.

Does the cancellation law hold? (Reminder: it means $x \star y = z \star y \implies x = z$)

Concatenation of Languages

Let A be alphabet. Consider the set of all languages $L \subseteq A^*$

Is this a monoid?

- ▶ Is there a neutral element? $L\{\varepsilon\} = L$, $\{\varepsilon\}L = L$, so $\{\varepsilon\}$ is one
- ▶ Which law needs to hold? Does it hold?

Associativity: we need $L_1 \cdot (L_2 \cdot L_3) = (L_1 \cdot L_2) \cdot L_3$

$$L_1 \cdot (L_2 \cdot L_3) = \{w_1(w_2w_3) \mid w_1 \in L_1 \wedge (w_2 \in L_2 \wedge w_3 \in L_3)\}$$

$$(L_1 \cdot L_2) \cdot L_3 = \{(w_1w_2)w_3 \mid (w_1 \in L_1 \wedge w_2) \in L_2 \wedge w_3 \in L_3\}$$

It is a monoid.

Does the cancellation law hold? (Reminder: it means $x \star y = z \star y \implies x = z$)

No, because there is an *absorbing* element, \emptyset .

For all L_1, L_2 we have $L_1\emptyset = \emptyset = L_2\emptyset$, but not necessarily $L_1 = L_2$

Representing Languages Through Programs

In general not possible: some formal languages are not *recursively enumerable* sets.

Reasonably powerful representation: computable characteristic functions.

A language $L \subseteq A^*$ is given by its *characteristic function* $f_L : A^* \rightarrow \{0,1\}$
defined by $f_L(w) = 1$ for $w \in L$ and $f_L(w) = 0$ for $w \notin L$.

Representing Languages Through Programs

In general not possible: some formal languages are not *recursively enumerable* sets.

Reasonably powerful representation: computable characteristic functions.

A language $L \subseteq A^*$ is given by its *characteristic function* $f_L : A^* \rightarrow \{0,1\}$
defined by $f_L(w) = 1$ for $w \in L$ and $f_L(w) = 0$ for $w \notin L$.

case class Lang[A](contains: List[A] \Rightarrow Boolean)

Representing Languages Through Programs

In general not possible: some formal languages are not *recursively enumerable* sets.

Reasonably powerful representation: computable characteristic functions.

A language $L \subseteq A^*$ is given by its *characteristic function* $f_L : A^* \rightarrow \{0,1\}$ defined by $f_L(w) = 1$ for $w \in L$ and $f_L(w) = 0$ for $w \notin L$.

```
case class Lang[A](contains: List[A]  $\Rightarrow$  Boolean)
```

Example: build the language $L_2 = \{(01)^n \mid n \geq 1\}$.

```
def f(w: List[Int]): Boolean = w match  
  case Cons(0, Cons(1, Nil()))  $\Rightarrow$  true  
  case Cons(0, Cons(1, wRest))  $\Rightarrow$  f(wRest)  
  case _  $\Rightarrow$  false  
val L2 = Lang(f)
```

```
L2.contains(0::1::0::1::Nil()) // true
```

Representing Language Concatenation

We can use code to express concatenation of computable languages.

```
def concat(L1: Lang[A], L2: Lang[A]): Lang[A] =  
  def f(w: List[A]) =  
    val n = w.length  
    def checkFrom(i: Int) =  
      require(0 <= i && i <= n)  
      (L1.contains(w.take(i)) && L2.contains(w.drop(i))) ||  
      (i < n && checkFrom(i + 1))  
    checkFrom(0, w.length)  
  Lang(f) // return the language with characteristic function f  
  
// take and drop are defined on lists:  
(a :: b :: c :: d :: e :: Nil()).take(2) = a :: b :: Nil()  
(a :: b :: c :: d :: e :: Nil()).drop(2) = c :: d :: e :: Nil()
```

Repetition of a Language: Kleene Star

Definition (Kleene star)

Given $L \subseteq A^*$, define

$$L^* = \bigcup_{n \geq 0} L^n$$

Repetition of a Language: Kleene Star

Definition (Kleene star)

Given $L \subseteq A^*$, define

$$L^* = \bigcup_{n \geq 0} L^n$$

Theorem

For $L \subseteq A^$, for every $w \in A^*$ we have $w \in L^*$ if and only if*

$$\exists n \geq 0. \exists w_1, \dots, w_n \in L. w = w_1 \dots w_n$$

Repetition of a Language: Kleene Star

Definition (Kleene star)

Given $L \subseteq A^*$, define

$$L^* = \bigcup_{n \geq 0} L^n$$

Theorem

For $L \subseteq A^$, for every $w \in A^*$ we have $w \in L^*$ if and only if*

$$\exists n \geq 0. \exists w_1, \dots, w_n \in L. w = w_1 \dots w_n$$

$$\{a\}^* = \{\varepsilon, a, aa, aaa, \dots\}$$

$$\{a, bb\}^* = \{\varepsilon, a, bb, abb, bba, aa, bbbb, aabb, \dots\} \text{ (describe this language)}$$

Repetition of a Language: Kleene Star

Definition (Kleene star)

Given $L \subseteq A^*$, define

$$L^* = \bigcup_{n \geq 0} L^n$$

Theorem

For $L \subseteq A^$, for every $w \in A^*$ we have $w \in L^*$ if and only if*

$$\exists n \geq 0. \exists w_1, \dots, w_n \in L. w = w_1 \dots w_n$$

$$\{a\}^* = \{\varepsilon, a, aa, aaa, \dots\}$$

$$\{a, bb\}^* = \{\varepsilon, a, bb, abb, bba, aa, bbbb, aabb, \dots\} \text{ (describe this language)}$$

- words whose all contiguous blocks of b -s have even length

Can L^* be finite for some L ? If so, describe all such L

Repetition of a Language: Kleene Star

Definition (Kleene star)

Given $L \subseteq A^*$, define

$$L^* = \bigcup_{n \geq 0} L^n$$

Theorem

For $L \subseteq A^*$, for every $w \in A^*$ we have $w \in L^*$ if and only if

$$\exists n \geq 0. \exists w_1, \dots, w_n \in L. w = w_1 \dots w_n$$

$$\{a\}^* = \{\varepsilon, a, aa, aaa, \dots\}$$

$$\{a, bb\}^* = \{\varepsilon, a, bb, abb, bba, aa, bbbb, aabb, \dots\} \text{ (describe this language)}$$

- words whose all contiguous blocks of b -s have even length

Can L^* be finite for some L ? If so, describe all such L

- $\{\varepsilon\}^* = \{\varepsilon\}$, $\emptyset^* = \{\varepsilon\}$; all others have at least one word of length ≥ 1 , so L^* is infinite

Star and the Empty Word

Concatenating with an empty word has no effect, so we have the following:

$$L^* = (L \setminus \{\epsilon\})^* = \{\epsilon\} \cup \bigcup_{n \geq 1} (L \setminus \{\epsilon\})^n$$

Moreover, $w \in L^*$ if and only if either $w = \epsilon$ or, for some n where $1 \leq n \leq |w|$,

$$w = w_1 \dots w_n$$

where $w_i \in L$ and $|w_i| \geq 1$ for all i where $1 \leq i \leq n$.

Star and the Empty Word

Concatenating with an empty word has no effect, so we have the following:

$$L^* = (L \setminus \{\varepsilon\})^* = \{\varepsilon\} \cup \bigcup_{n \geq 1} (L \setminus \{\varepsilon\})^n$$

Moreover, $w \in L^*$ if and only if either $w = \varepsilon$ or, for some n where $1 \leq n \leq |w|$,

$$w = w_1 \dots w_n$$

where $w_i \in L$ and $|w_i| \geq 1$ for all i where $1 \leq i \leq n$.

- ▶ we omit ε because it leaves concatenation the same
- ▶ we can assume $n \leq |w|$ because all blocks have length at least one

Star and the Empty Word

Concatenating with an empty word has no effect, so we have the following:

$$L^* = (L \setminus \{\varepsilon\})^* = \{\varepsilon\} \cup \bigcup_{n \geq 1} (L \setminus \{\varepsilon\})^n$$

Moreover, $w \in L^*$ if and only if either $w = \varepsilon$ or, for some n where $1 \leq n \leq |w|$,

$$w = w_1 \dots w_n$$

where $w_i \in L$ and $|w_i| \geq 1$ for all i where $1 \leq i \leq n$.

- ▶ we omit ε because it leaves concatenation the same
- ▶ we can assume $n \leq |w|$ because all blocks have length at least one

If L is computable (has a computable characteristic function), is L^* also computable?

Star and the Empty Word

Concatenating with an empty word has no effect, so we have the following:

$$L^* = (L \setminus \{\varepsilon\})^* = \{\varepsilon\} \cup \bigcup_{n \geq 1} (L \setminus \{\varepsilon\})^n$$

Moreover, $w \in L^*$ if and only if either $w = \varepsilon$ or, for some n where $1 \leq n \leq |w|$,

$$w = w_1 \dots w_n$$

where $w_i \in L$ and $|w_i| \geq 1$ for all i where $1 \leq i \leq n$.

- ▶ we omit ε because it leaves concatenation the same
- ▶ we can assume $n \leq |w|$ because all blocks have length at least one

If L is computable (has a computable characteristic function), is L^* also computable?

- ▶ try all possible ways of splitting w
- ▶ if $k = |w|$, for each point between the letters of w you can decide to split there or not, so there are 2^{k-1} ways to split: $w = \square \underbrace{|\square| \dots |\square|}_{k-1} \square$
- ▶ there is a better way - see exercises

Starring: $\{a, ab\}$

Let $A = \{a, b\}$ and $L = \{a, ab\}$.

Come up with a property “...” that describes the language L^* :

$$L^* = \{w \in A^* \mid \dots\}$$

Prove that the property and L^* denote the same language.

Starring: $\{a, ab\}$

Let $A = \{a, b\}$ and $L = \{a, ab\}$.

Come up with a property “...” that describes the language L^* :

$$L^* = \{w \in A^* \mid \dots\}$$

Prove that the property and L^* denote the same language.

Properties:

- ▶ does not begin with b
- ▶ does not contain bb

in one phrase: there is an “ a ” before every “ b ”