

COMP 170 – Fall 2006

Midterm 1 Solution

Q1. Suppose that we have a student hall with 9 rooms labelled A, B, \dots, I .

How many ways are there to assign 10 students to the 9 hall rooms so that every student is assigned and no hall room is empty?

This is exactly the number of onto functions from S_{10} to S_9 (S_i , being a set with i elements)

This is exactly the number of onto functions from S_{10} to S_9 (S_i , being a set with i elements)

Note that there should be exactly one pair of students get assigned to the same room, while all the others get assigned an individual room.

There are $\binom{10}{2}$ ways of choosing that pair.

This is exactly the number of onto functions from S_{10} to S_9 (S_i , being a set with i elements)

Note that there should be exactly one pair of students get assigned to the same room, while all the others get assigned an individual room.

There are $\binom{10}{2}$ ways of choosing that pair.

Once the pair is chosen, there are $9!$ ways of assigning the students to the rooms.

This is exactly the number of onto functions from S_{10} to S_9 (S_i , being a set with i elements)

Note that there should be exactly one pair of students get assigned to the same room, while all the others get assigned an individual room.

There are $\binom{10}{2}$ ways of choosing that pair.

Once the pair is chosen, there are $9!$ ways of assigning the students to the rooms.

So the answer is

$$\binom{10}{2} 9! = 16329600$$

Q2. Suppose you have 5 standard six-sided dice; one red, one green, one yellow, one black, one orange. Each die has the numbers 1-6 on its sides. Roll all of the dice. An “outcome” of a roll is a number for each die.

An outcome contains a

- (i) *two-of-a-kind* if there are at least two dice showing the same number;
- (ii) *three-of-a-kind* if there are at least three dice showing the same number;
- (iii) *four-of-a-kind* if there are at least four dice showing the same number;
- (iv) *five-of-a-kind* if all five dice show the same number.

(a) How many possible outcomes are there?

(a) How many possible outcomes are there?

$$6^5 = 7776$$

(a) How many possible outcomes are there?

$$6^5 = 7776$$

(b) How many possible outcomes contain a *five-of-a-kind*?

(a) How many possible outcomes are there?

$$6^5 = 7776$$

(b) How many possible outcomes contain a *five-of-a-kind*?

$$6$$

(c) How many possible outcomes contain a *four-of-a-kind* but no *five-of-a-kind*?

(c) How many possible outcomes contain a *four-of-a-kind* but no *five-of-a-kind*?

Each such outcome is uniquely determined by

- (i) the value of the *four-of-a-kind*,
- (ii) the color of the die that is not part of the *four-of-a-kind* and
- (iii) the value of that die. So the answer is

$$6 \cdot 5 \cdot 5 = 150.$$

(d) How many possible outcomes contain a *three-of-a-kind* but no *four-of-a-kind*?

(d) How many possible outcomes contain a *three-of-a-kind* but no *four-of-a-kind*?

Each such outcome is uniquely determined by

- (i) the value of the *three-of-a-kind*,
- (ii) the colors of the die that are **not** part of the *three-of-a-kind* and
- (iii) the values of those dice. So the answer is

$$6 \cdot \binom{5}{2} \cdot 5^2 = 1500.$$

(e) How many possible outcomes contain a *three-of-a-kind* AND a *two-of-a-kind* that have different dice values but no *five-of-a-kind*?

(e) How many possible outcomes contain a *three-of-a-kind* AND a *two-of-a-kind* that have different dice values but no *five-of-a-kind*?

Each such outcome is uniquely determined by

- (i) the value of the *three-of-a-kind*,
- (ii) the colors of the die that are **not** part of the *three-of-a-kind* and
- (iii) the values of those dice. So the answer is

$$6 \cdot \binom{5}{2} \cdot 5 = 300.$$

Q3. Is the following formula correct for all $n \geq 1$?

$$\sum_{i=0}^n 3^i \binom{n}{i} = 2^n \sum_{i=0}^n \binom{n}{i}.$$

If yes, prove it. If no, give a value of n for which the formula is incorrect.

Q3. Is the following formula correct for all $n \geq 1$?

$$\sum_{i=0}^n 3^i \binom{n}{i} = 2^n \sum_{i=0}^n \binom{n}{i}.$$

If yes, prove it. If no, give a value of n for which the formula is incorrect.

The proof is that the both equal 4^n .

This can be derived using two applications of the binomial theorem.

$$4^n = (3 + 1)^n = \sum_{i=0}^n 3^i 1^{n-i} \binom{n}{i}$$

$$4^n = (3 + 1)^n = \sum_{i=0}^n 3^i 1^{n-i} \binom{n}{i}$$

and

$$4^n = 2^n 2^n = 2^n \sum_{i=0}^n \binom{n}{i}$$

where the $2^n = \sum_{i=0}^n \binom{n}{i}$ was derived in class.

$$4^n = (3 + 1)^n = \sum_{i=0}^n 3^i 1^{n-i} \binom{n}{i}$$

and

$$4^n = 2^n 2^n = 2^n \sum_{i=0}^n \binom{n}{i}$$

where the $2^n = \sum_{i=0}^n \binom{n}{i}$ was derived in class.

Alternatively, the second equality can be derived as

$$4^n = (2 + 2)^n = \sum_{i=0}^n 2^i 2^{n-i} \binom{n}{i} = 2^n \sum_{i=0}^n \binom{n}{i}$$

Q4. A *permutation* of the letters
A, B, C, D, E, F, G, H is a list (string)
of the eight letters in some order.

A permutation *contains* the substring
ABC if the letters appear in the given or-
der. For example, permutations
DEFABCGH and HGABCFED
contain the substring ABC but
ADBECFGH doesn't.

(a) How many *permutations* of the letters
A, B, C, D, E, F, G, H contain the substring
ABC?

(a) How many *permutations* of the letters
A, B, C, D, E, F, G, H contain the substring
ABC?

To solve this problem consider substring **ABC** as
an indivisible item and each of the other individual
5 letters as a separate indivisible item.

Then the answer is the number of ways of writing a
permutation of 6 items (the five letters and **ABC**)
which is

$$6! = 720.$$

(b) How many *permutations* of the letters *A,B,C,D,E,F,G,H* contain *A* before *B* and *B* before *C*?

(b) How many *permutations* of the letters **A,B,C,D,E,F,G,H** contain **A** before **B** and **B** before **C**?

Such a permutation is uniquely defined by

(i) choosing the locations of the three letters **A,B,C** (once the locations are chosen we know that **A** must go in the first, **B** in the 2nd and **C** in the 3rd) and

(ii) the locations of the five remaining letters in the 5 remaining spaces.

This is therefore

$$\binom{8}{3} \cdot 5! = \frac{8!}{3!} = 6720.$$

Q5. An *anagram* is a distinct ordering of the letters. For example, the word **eat** has six anagrams “eat”, “eta”, “ate”, “aet”, “tea”, “tae”, while the word “**too**” has only three anagrams “too”, “oto” and “oot”.

(a) How many anagrams does the word “mammal” have?

(a) How many anagrams does the word “mammal” have?

This is the number of ways to label 6 items so that

- (i) 3 are labeled 'm',
- (ii) 2 are labelled 'a' and
- (iii) 1 is labelled 'l'

which is

$$\binom{6}{3} \cdot \binom{3}{2} \cdot \binom{1}{1} = \frac{6!}{3! \cdot 2! \cdot 1!} = 60$$

(b) How many anagrams does the word
“mississippi” have?

(b) How many anagrams does the word
“mississippi” have?

This is the number of ways to label 11 items so that

- (i) 1 is labeled 'm',
 - (ii) 4 are labelled 'i',
 - (iii) 4 are labelled 's' and
 - (iv) 2 are labelled 'p'
- which is

$$\binom{11}{1} \cdot \binom{10}{4} \cdot \binom{6}{4} \cdot \binom{2}{2} = \frac{11!}{1! \cdot 4! \cdot 4! \cdot 2!} = 34650$$

Q6. Give a combinatorial proof of the identity, for all $n \geq 9$,

$$\binom{n}{2} \binom{n-2}{3} \binom{n-5}{4} = \binom{n}{4} \binom{n-4}{3} \binom{n-7}{2}.$$

Q6. Give a combinatorial proof of the identity, for all $n \geq 9$,

$$\binom{n}{2} \binom{n-2}{3} \binom{n-5}{4} = \binom{n}{4} \binom{n-4}{3} \binom{n-7}{2}.$$

“Consider the problem of how to color n items so that 2 are red, 3 are green, 4 are blue and the remaining $n - 9$ are yellow.”

The left hand side of the inequality obviously counts this.

“Old problem was how to color n items so that 2 are red, 3 are green, 4 are blue and the remaining $n - 9$ are yellow.

Notice that the problem is the same if we change the order in which we ask the question.

“Consider the problem of how to color n items so that 4 are blue, 3 are green, 2 are red and the remaining $n - 9$ are yellow.”

This is what the right hand side of the equation is counting, so the two sides are the same.

Q7. Consider the following statement

$$\gcd(j, k) = \gcd(k - j, j).$$

Is this statement always true for k, j with $k \geq j \geq 0$?
Either *prove* that it is true for all k, j with $k \geq j \geq 0$,
or give values for k, j with $k \geq j \geq 0$ such that
 $\gcd(j, k) \neq \gcd(k - j, j)$.

We will show that

d is a common divisor of j and k

if and only if

d is a common divisor of $k - j$ and j .

This shows that

the set of common divisors of j and k

is exactly the same as

the set of common divisors of $k - j$ and j .

This immediately implies that the

greatest common divisor of j and k

is the same as the

greatest common divisor of $k - j$ and j .

To prove the *if* direction

suppose that d is a common divisor of j and k .

Then, for some a and b

$$j = a \cdot d \quad \text{and} \quad k = b \cdot d$$

so

$$k - j = (b - a)d$$

and

d is a divisor of $k - j$.

Thus d is a common divisor of $k - j$ and j .

To prove the *only if* direction

suppose that d is a common divisor of $k - j$ and j .

Then, for some c and a

$$j = a \cdot d \quad \text{and} \quad k - j = c \cdot d$$

so

$$k = (k - j) + j = (c + a)d$$

and

d is a divisor of k .

Thus d is a common divisor of k and j .

Q8. (a) Consider the equation

$$3 \cdot_{100} x = 13$$

Does this equation have a solution x in Z_{100} ?

Q8. (a) Consider the equation

$$3 \cdot_{100} x = 13$$

Does this equation have a solution x in Z_{100} ?

(b) Consider the equation

$$15 \cdot_{100} x = 13$$

Does this equation have a solution x in Z_{100} ?

(a) Yes.

To start, first notice that $3 \cdot (-33) + 1 \cdot 100 = 1$.

(This can either be derived using the extended GCD algorithm or just by observation.)

This tells us that the multiplicative inverse of 3 in Z_{100} is $-33 \bmod 100 = 67$.

So, the answer is

$$x = 67 \cdot_{100} 13 = 871 \bmod 100 = 71.$$

As a reality check, note that

$$3 \cdot_{100} 71 = 213 \bmod 100 = 13$$

so $x = 71$ really does solve our problem.

(b) No.

This can be proven by contradiction.

Suppose there was some solution x .

Then $15 \cdot_{100} x = 13$ so there is some q such that $15 \cdot x = q \cdot 100 + 13$ or

$$15 \cdot x - q \cdot 100 = 13.$$

But the left hand side of this equation is divisible by 5 while the right hand side is not, leading to a contradiction.

(b) No.

This can be proven by contradiction.

Suppose there was some solution x .

Then $15 \cdot_{100} x = 13$ so there is some q such that $15 \cdot x = q \cdot 100 + 13$ or

$$15 \cdot x - q \cdot 100 = 13.$$

But the left hand side of this equation is divisible by 5 while the right hand side is not, leading to a contradiction.

*Note: the fact that $\gcd(15, 100) = 5 \neq 1$ does **not** prove that $15 \cdot_{100} x = 13$ has no answer.*