

Research Statement

Henry Kvinge

1 Introduction

My research interests in mathematics focus on the connections between representation theory, combinatorics, and probability theory. I am particularly interested in the interaction between combinatorial representation theory and categorical representation theory. In this research statement I will describe some of my past and current projects which include:

- investigating a surprising connection between graphical Heisenberg categories and symmetric functions and stochastic growth processes.
- Generalizing the combinatorial/probabilistic framework for studying asymptotic processes on symmetric groups to a broader family of algebras
- the study of crystal graphs associated to affine type quantum groups and their connections to the representation theory of cyclotomic quiver Hecke algebras.

I am also doing work in geometric data analysis, machine learning, and compressive sensing. I briefly summarize some of these projects in Section 3.

2 Representation theory and combinatorics

2.1 The categorification program

Initiated by Crane and Frenkel [5], the idea underlying the categorification program is that many of the fundamental structures in algebra, geometry, and topology are the shadow of richer (often undiscovered) higher categories. Categorification, the process of revealing these higher analogues, then involves lifting set-theoretic constructions to category-theoretic constructions, roughly: replacing sets with categories, functions with functors, and equalities with isomorphisms. Though Khovanov's categorification of the Jones polynomial via Khovanov homology [16] may be the most famous example, far more familiar examples exist (for example, singular homology as a categorification of Betti numbers). Categorifying the fundamental algebras from representation theory has been a major, ongoing project in the field. Furthermore, theory developed in the course of this program has led to results of independent interest, such as the proof of one of the Kazhdan-Lusztig conjectures for all Coxeter types [6] via Soergel bimodules [37].

As consequence of this progress, a host of new categories have been discovered which sit “above” and “control” many of the classical objects in representation theory such as Hecke algebras, Heisenberg algebras, and quantum groups. These categories are rich in structure, and a major drive in my research has been using these categorical constructions to better understand the connections between representation theory and other areas of math, particularly combinatorics and probability theory. My first research project described below studies how categorifications of infinite dimensional Heisenberg algebras both provide graphical realizations of analogues of the symmetric functions and also naturally provide the framework for

stochastic growth processes motivated by representation theory and combinatorics [24, 25]. In my second project, the categorification of quantum groups, via quiver Hecke algebras, is used to show how a fundamental isomorphism of combinatorially defined crystals can in fact be understood to be a consequence of induction/restriction functors from representation theory [26].

2.2 Heisenberg categories, symmetric functions, and growth processes

A *tower of algebras* $\{A_n\}_{n \geq 0}$ is a collection of nested algebras $A_0 \subseteq A_1 \subseteq A_2 \subseteq \dots$. The classic example is the symmetric group algebras $\{\mathbb{C}[S_n]\}_{n \geq 0}$. A result of Geissinger [8] showed that the direct sum of all Grothendieck groups of symmetric groups $\bigoplus_{n \geq 0} K_0(\mathbb{C}[S_n]\text{-mod})$ (which is isomorphic to the algebra of symmetric functions) carries the structure of the Fock space representation of the infinite dimensional Heisenberg algebra \mathfrak{h} . In particular, the elements of \mathfrak{h} act by linear operators coming from induction and restriction functors between symmetric group representations. This observation was the first hint of a deeper story in which a vastly richer *Heisenberg category* \mathcal{H} , acts via genuine induction and restriction functors on the module categories $\bigoplus_{n \geq 0} A_n\text{-mod}$ for a tower of algebras $\{A_n\}_{n \geq 0}$.

$$\begin{array}{ccc}
 \mathcal{H} & \xrightarrow{\text{categorical representation}} & \bigoplus_{n \geq 0} A_n\text{-mod} \\
 \downarrow \text{zigzag } K_0 & & \downarrow \text{zigzag } K_0 \\
 \mathfrak{h} & \xrightarrow{\text{representation}} & \bigoplus_{n \geq 0} K_0(A_n\text{-mod})
 \end{array}$$

The first Heisenberg category was defined by Khovanov [17] with $A_n = \mathbb{C}[S_n]$. Subsequently, Heisenberg categories associated to a range of towers of algebras have been discovered, including: Hecke algebras [29], Sergeev algebras [4], wreath product algebras [34], and degenerate cyclotomic Hecke algebras [30, 3]. In all these examples, \mathcal{H} can be defined diagrammatically, i.e. through an isotopy invariant calculus of planar diagrams with local relations and decorations.


All Heisenberg categories are additive monoidal categories and hence their center, the endomorphisms of the monoidal identity $Z(\mathcal{H}) := \text{End}_{\mathcal{H}}(\mathbf{1})$, is a commutative algebra. In such diagrammatically defined categories, the center has a helpful diagrammatic definition as the algebra of all closed diagrams.

One of my ongoing projects is centered around studying the commutative algebra $Z(\mathcal{H})$ corresponding to different towers of algebras. This object is rich in combinatorial and representation-theoretic data. We showed that the center of Khovanov's Heisenberg category, $Z(\mathcal{H}_{\mathbb{C}[S_n]})$, is isomorphic to Λ^* , the algebra of shifted symmetric functions [24]. A deformation of the classical symmetric functions, Λ^* was discovered by Kerov and Olshanski

[15] and further investigated by Okounkov and Olshanski [32] among many others. It has connections to Lie theory [32], Gromov-Witten invariants [33], and asymptotic representation theory [10].

As a consequence of our algebra isomorphism $Z(\mathcal{H}_{\mathbb{C}[S_n]}) \cong \Lambda^*$, we were able to describe an isotopy invariant graphical calculus for Λ^* using closed diagrams and a set of local relations. We showed how individual generators such as shifted Schur functions and properties of Λ^* such as involutions, have a natural graphical interpretation.

Surprisingly, under the isomorphism $Z(\mathcal{H}_{\mathbb{C}[S_n]}) \cong \Lambda^*$, the simplest generating diagrams (which are clockwise or counterclockwise oriented *bubbles* with k dots on them)



are shifted symmetric functions encoding moments for a family of probability measures, Kerov's *transition* and *co-transition measures*, parametrized by Young diagrams. This same probability measure was used as a tool to study the famous Plancherel growth process [13, 14]. This unexpected connection points to a deeper relationship between the combinatorics of categorical representation theory and the combinatorics of asymptotic representation theory. Exploring this connection more closely is one of the themes of my current work.

The results described above do not seem to be a phenomenon restricted to symmetric groups. In [25], we investigated the center of the twisted Heisenberg category $\mathcal{H}_{\mathbb{S}_n}$ of Cautis and Sussan [4] which is associated to the tower of Sergeev algebras $\{\mathbb{S}_n\}_{n \geq 0}$. We showed that $Z(\mathcal{H}_{\mathbb{S}_n})$ is isomorphic to the subalgebra of the symmetric functions generated by odd power sums $\mathbb{C}[p_1, p_3, \dots]$, and similar connections to combinatorics and asymptotic representation theory hold. Notably, the bubble generators of $Z(\mathcal{H}_{\mathbb{S}_n})$ also correspond to functions encoding transition probabilities of a stochastic growth process (this time on the graph of all strict Young diagrams).

Given the observations described above, one long-term goal of my research is to use algebraic, combinatorial, and functorial tools to generalize some of the rich mathematics appearing in the asymptotic representation theory of symmetric groups to more general towers of algebras. For example, it would be interesting to understand how groundbreaking results such as the limit shape of Young diagrams under the Plancherel growth process generalizes other towers of algebras. A first step in this direction consists in generalizing one particular framework used in asymptotic representation theory called a *coherent system*. First introduced by Borodin-Olshanski in [2] to study infinite-dimensional diffusion processes, a coherent system on a graded graph is a sequence of probability measures $\{M_n\}_{n \geq 0}$ which respect certain up/down transition functions p_\uparrow, p_\downarrow between graded components. One natural coherent system on the graded graph of Young diagrams is the sequence of Plancherel measures with up/down transition functions defined via induction and restriction functors applied irreducible symmetric group representations.

In [20], we show that when a tower of algebras $\{A_n\}_{n \geq 0}$ is a *free Frobenius tower* (that is, each algebra A_{n+1} is both a Frobenius extension of A_n and a free (A_n, A_n) -bimodule) then:

- There are generally two coherent systems which generalize this “Plancherel” coherent system. In one, the down transition functions are controlled by the dimension of

indecomposable projective modules and up transition functions are controlled by the dimensions of simple modules, and in the other the reverse is true. Of course, when $\{A_n\}_{n \geq 0}$ is semisimple, the above distinction is hidden.

- Data for one of these coherent systems is captured by two families of central elements in $\{Z(A_n)\}_{n \geq 0}$.


The goal of the framework above is to open the door to the study of new towers of algebras. Some other projects that I am either currently working on or plan to work on in the near future are:

- *Centers of Heisenberg categories, asymptotic representation theory, and new analogues of symmetric functions:* The centers of Heisenberg categories allow one to capture the structure of the centers of all algebras in a tower, i.e. $\{Z(A_n)\}_{n \geq 0}$, simultaneously. In previous examples, this construction has yielded analogues of the symmetric functions with rich combinatorial structure (for example: shifted symmetric functions). I would like to understand what analogues can be obtained from more exotic towers of algebras such as degenerate cyclotomic Hecke algebras or Frobenius wreath product algebras.
- *General representation-theoretic growth processes and Heisenberg categories:* W

2.3 Crystal structure in representation categories

Crystals were first discovered by Kashiwara as combinatorial objects attached to certain quantum group $U_q(\mathfrak{g})$ representations that appear in the limit $q \rightarrow 0$ [12]. Since they can be described via colored directed graphs, crystals are substantially simpler than their corresponding representations, and in many cases allow us to translate questions in algebra to questions in combinatorics.

Crystals can also arise when a quantum group acts on a category via endofunctors. The first example of this was found when it was noticed that graphs defined via the branching rules of affine and cyclotomic Hecke algebras carry the structure of a crystal graph [27, 1, 9]. Only in hindsight was it realized that this is the result of a hidden categorical action of a quantum group. Subsequently many other representation categories with categorical action of a quantum group have been shown to carry a crystal structure. Examples include the category of finite-dimensional quiver Hecke algebra representations [28] and category \mathcal{O} for the cyclotomic rDAHA [36].



When a representation category carries a crystal structure one can try to understand how the combinatorics of the crystal lift to the representation theory. Such crystal-theoretic considerations allowed Lauda-Vazirani to classify all simple quiver Hecke algebra representations [28].




Figure 1: Young's lattice is a combinatorial model for the crystal corresponding to the highest weight representation V_{Λ_0} of $U_q(\mathfrak{sl}_\infty)$.

2.4 Results on crystal structure in categories

Khovanov and Lauda [18], [19] and independently Rouquier [35] invented *quiver Hecke algebras* R (also known as *Khovanov-Lauda-Rouquier (KLR) algebras*) to categorify the upper half $U_q^+(\mathfrak{g})$ of the quantum group for any fixed symmetrizable Kac-Moody algebra \mathfrak{g} .

For each integral dominant weight Λ , the algebra R has a finite-dimensional quotient called a *cyclotomic quiver Hecke algebra* R^Λ . Lauda-Vazirani showed that the collection of simple R^Λ -modules carries the structure of $B(\Lambda)$ which is the crystal of the corresponding highest weight representation of $U_q(\mathfrak{g})$. Here nodes are the simple R^Λ -modules and Kashiwara operators \tilde{f}_i are given by refined induction functors between the simple modules [28].

When \mathfrak{g} is of classical affine type, one class of crystals not previously appearing in the context of quiver Hecke algebras is the *Kirillov-Reshetikhin (KR) crystals* [31]. These are denoted by $B^{r,s}$ where $r \in I \setminus \{0\}$ for Dynkin indexing set I and $s \in \mathbb{Z}_{>0}$. A notable property of $B^{1,1}$ in particular is that for certain choices of fundamental weight Λ_j there is an isomorphism of crystals

$$\varphi : B^{1,1} \otimes B(\Lambda_j) \xrightarrow{\sim} B(\Lambda_i), \quad (1)$$

for some $i \in I$ [11]. This isomorphism is particularly exciting since it allows one to study the (infinite) crystal $B(\Lambda_i)$ through tensor products of the finite crystals $B^{1,1}$.

Given that R^{Λ_i} -mod (respectively R^{Λ_j} -mod) carries the structure of $B(\Lambda_i)$ (resp. $B(\Lambda_j)$), it is natural to ask whether (1) actually reflects representation-theoretic structure in R^{Λ_i} -mod. Vazirani and I showed that this is the case in [26]. Recall that each simple R^{Λ_i} -module M^\bullet corresponds to a node \bullet in $B(\Lambda_i)$, and similarly each simple R^{Λ_j} -module M^\bullet corresponds to a node \bullet in $B(\Lambda_j)$. We show that each assignment of the crystal isomorphism (1) is actually the shadow of a canonically defined surjective R^{Λ_i} -module homomorphism ϕ .

- *Searching for pure signals in hyperspectral imagery via the geometry of the Grassmannian*: Unlike RGB images, which only sample from three different spectral bands of light (red, green, and blue), hyperspectral imagery can sometimes sample more than 200. While hyperspectral images thus have strong discriminatory power, it can be difficult to extract information from them because they are often very high dimensional. One particularly important problem is understanding how to isolate the “most pure” signals from a hyperspectral image. Geometrically, this corresponds to picking out those spatial locations in the image whose spectral curve sits on the convex hull formed by all spectral curves in the image. In [7] we show how if one considers local patches rather than individual pixels, this endmember extraction problem is most naturally realized on a Grassmann manifold. We propose an algorithm to solve the problem in this setting.
- *Compressive sensing algorithm design for a single-pixel camera*¹: Over the course of the last year, I have been involved with two projects with an industry partner which involve developing compressive sensing (CS) algorithms for single pixel imaging devices. This has involved choosing a CS sampling framework which captures the structure of images and also fits hardware constraints, developing and testing CS reconstruction algorithms which can reconstruct images in real-time with significant device noise, and writing a software package that implements all these algorithms for use in the field. We are currently in the process of writing three conference papers related to this work.

My current and future work in this area focuses on bringing concepts and methods from pure math to bear on problems in data analytics and machine learning. Two projects currently in progress include:

- *Schubert varieties and attribute discovery for machine learning*: One foundational problem in machine learning and data analysis is attribute discovery within a data set. We have evidence to suggest that when data can be represented on a Grassmann manifold, then Schubert varieties are a natural framework in which to differentiate fundamental attributes associated. We currently in the process of not only making this association rigorous, but also adapting various optimization algorithms from Euclidean space to the Grassmannian framework with Schubert constraints.
- *Decomposition of the regular representation and dimensionality reduction*: The multi-dimensional scaling (MDS) algorithm is a foundational algorithm which can be used both for dimensionality reduction and more generally for realization of a finite configuration of points from an abstract metric space in Euclidean space. It is thus a basic algorithm used in many areas of science and engineering. In the special case when the points of interest live on a group, then the algorithm can be given a representation theoretic interpretation. This observation both sheds new light on the algorithm itself and suggests alternative implementations.

References

- [1] Susumu Ariki, *On the decomposition numbers of the Hecke algebra of $G(m, 1, n)$* , J. Math. Kyoto Univ. **36** (1996), no. 4, 789–808.

¹Work described in this section is ITAR-protected and hence some details have been omitted.

- [2] Alexei Borodin and Grigori Olshanski, *Infinite-dimensional diffusions as limits of random walks on partitions*, Probab. Theory Related Fields **144** (2009), no. 1-2, 281–318.
- [3] J. Brundan, *On the definition of Heisenberg category*, To appear in Algebraic Combinatorics (2017), arXiv:1709.06589.
- [4] Sabin Cautis and Joshua Sussan, *On a categorical Boson-Fermion correspondence*, Comm. Math. Phys. **336** (2015), no. 2, 649–669.
- [5] Louis Crane and Igor B. Frenkel, *Four-dimensional topological quantum field theory, Hopf categories, and the canonical bases*, J. Math. Phys. **35** (1994), no. 10, 5136–5154, Topology and physics.
- [6] Ben Elias and Geordie Williamson, *The Hodge theory of Soergel bimodules*, Ann. of Math. (2) **180** (2014), no. 3, 1089–1136.
- [7] Elin Farnell, Henry Kvinge, Michael Kirby, and Chris Peterson, *Endmember extraction on the Grassmannian*, 2018 IEEE Data Science Workshop (DSW), June 2018, pp. 71–75.
- [8] Ladnor Geissinger, *Hopf algebras of symmetric functions and class functions*, Combinatoire et représentation du groupe symétrique (Actes Table Ronde C.N.R.S., Univ. Louis-Pasteur Strasbourg, Strasbourg, 1976), Springer, Berlin, 1977, pp. 168–181. Lecture Notes in Math., Vol. 579.
- [9] Ian Grojnowski, *Affine sl_p controls the representation theory of the symmetric group and related Hecke algebras*, arXiv:9907129 (1999).
- [10] Vladimir Ivanov and Grigori Olshanski, *Kerov’s central limit theorem for the Plancherel measure on Young diagrams*, Symmetric functions 2001: surveys of developments and perspectives, NATO Sci. Ser. II Math. Phys. Chem., vol. 74, Kluwer Acad. Publ., Dordrecht, 2002, pp. 93–151.
- [11] Seok-Jin Kang, Masaki Kashiwara, Kailash C. Misra, Tetsuji Miwa, Toshiki Nakashima, and Atsushi Nakayashiki, *Perfect crystals of quantum affine Lie algebras*, Duke Math. J. **68** (1992), no. 3, 499–607.
- [12] Masaki Kashiwara, *On crystal bases of the Q -analogue of universal enveloping algebras*, Duke Math. J. **63** (1991), no. 2, 465–516.
- [13] Sergei Kerov, *Transition probabilities of continual Young diagrams and the Markov moment problem*, Funktsional. Anal. i Prilozhen. **27** (1993), no. 2, 32–49, 96.
- [14] ———, *A differential model for the growth of Young diagrams*, Proceedings of the St. Petersburg Mathematical Society, Vol. IV, Amer. Math. Soc. Transl. Ser. 2, vol. 188, Amer. Math. Soc., Providence, RI, 1999, pp. 111–130.
- [15] Sergei Kerov and Grigori Olshanski, *Polynomial functions on the set of Young diagrams*, C. R. Acad. Sci. Paris Sér. I Math. **319** (1994), no. 2, 121–126.
- [16] Mikhail Khovanov, *A categorification of the Jones polynomial*, Duke Math. J. **101** (2000), no. 3, 359–426.

- [17] ———, *Heisenberg algebra and a graphical calculus*, Fund. Math. **225** (2014), no. 1, 169–210.
- [18] Mikhail Khovanov and Aaron D. Lauda, *A diagrammatic approach to categorification of quantum groups. I*, Represent. Theory **13** (2009), 309–347.
- [19] ———, *A diagrammatic approach to categorification of quantum groups II*, Trans. Amer. Math. Soc. **363** (2011), no. 5, 2685–2700.
- [20] Henry Kvinge, *Coherent systems of probability measures on graphs for representations of free Frobenius towers*, arXiv:? (2018).
- [21] Henry Kvinge, Elin Farnell, Michael Kirby, and Chris Peterson, *A GPU-oriented algorithm design for secant-based dimensionality reduction*, 2018 17th International Symposium on Parallel and Distributed Computing (ISPDC), June 2018, pp. 69–76.
- [22] ———, *Monitoring the shape of weather, soundscapes, and dynamical systems: a new statistic for dimension-driven data analysis on large data sets*, accepted for publication in the Proceedings of the IEEE International Conference on Big Data 2018, August 2018.
- [23] Henry Kvinge, Elin Farnell, Michael Kirby, and Chris Peterson, *Too many secants: a hierarchical approach to secant-based dimensionality reduction on large data sets*, to appear in Proceedings of the 2018 IEEE High Performance Extreme Computing Conference, IEEE, 2018.
- [24] Henry Kvinge, Anthony M. Licata, and Stuart Mitchell, *Khovanov’s Heisenberg category, moments in free probability, and shifted symmetric functions*, arXiv:1610.04571 (2016), To appear in Algebraic Combinatorics.
- [25] Henry Kvinge, Can Ozan Oğuz, and Michael Reeks, *The center of the twisted Heisenberg category, factorial Schur Q -functions, and transition functions on the Schur graph*, arXiv:1712.09626 (2017), To appear in the Journal of Algebraic Combinatorics.
- [26] Henry Kvinge and Monica Vazirani, *A combinatorial categorification of the tensor product of the Kirillov-Reshetikhin crystal $B^{1,1}$ and a fundamental crystal*, Algebras and Representation Theory (2017).
- [27] Alain Lascoux, Bernard Leclerc, and Jean-Yves Thibon, *Hecke algebras at roots of unity and crystal bases of quantum affine algebras*, Comm. Math. Phys. **181** (1996), no. 1, 205–263.
- [28] Aaron D. Lauda and Monica Vazirani, *Crystals from categorified quantum groups*, Adv. Math. **228** (2011), no. 2, 803–861.
- [29] Anthony Licata and Alistair Savage, *Hecke algebras, finite general linear groups, and Heisenberg categorification*, Quantum Topol. **4** (2013), no. 2, 125–185.
- [30] Marco Mackaay and Alistair Savage, *Degenerate cyclotomic Hecke algebras and higher level Heisenberg categorification*, J. Algebra **505** (2018), 150–193.

- [31] Masato Okado and Anne Schilling, *Existence of Kirillov-Reshetikhin crystals for nonexceptional types*, Represent. Theory **12** (2008), 186–207.
- [32] Andrei Okounkov and Grigori Olshanski, *Shifted Schur functions*, Algebra i Analiz **9** (1997), no. 2, 73–146.
- [33] Andrei Okounkov and Rahul Pandharipande, *Gromov-Witten theory, Hurwitz theory, and completed cycles*, Ann. of Math. (2) **163** (2006), no. 2, 517–560.
- [34] Daniele Rosso and Alistair Savage, *A general approach to Heisenberg categorification via wreath product algebras*, Math. Z. **286** (2017), no. 1-2, 603–655.
- [35] Raphaël Rouquier, *2-Kac-Moody algebras*, arXiv:0812.5023 (2008).
- [36] Peng Shan, *Crystals of Fock spaces and cyclotomic rational double affine Hecke algebras*, Ann. Sci. Éc. Norm. Supér. (4) **44** (2011), no. 1, 147–182.
- [37] Wolfgang Soergel, *Kategorie \mathcal{O} , perverse Garben und Moduln über den Koinvarianten zur Weylgruppe*, J. Amer. Math. Soc. **3** (1990), no. 2, 421–445.