



Mountain View Community Hospital

Clinical Nutrition Monitoring & ONS Utilization Analytics Dashboard | 2025



All

Main Cause

All

Total Transactions

3000

Total Patients

500

ONS Patients

274

Non ONS Patient

226

ONS %

54.80%

Non ONS %

45.20%

ONS

All

Patient Name

All


ALB

All


NPCR

All


ONS Patients by Month Name




Risk Category by Patient




Low ALB & NPCR Patients



Risk Category Status



Risk Category by Main Cause



- This dashboard monitors nutritional risk across 3,000 patients.
 - Approximately 55% require oral nutrition supplementation, indicating high intervention needs.
 - While average albumin remains stable at 38 g/L, the average NPCR is 0.65, which is below recommended protein intake levels.
 - The main drivers of risk are gastrointestinal conditions and poor appetite. Monthly trends show stable ONS utilization, suggesting consistent screening processes. However, protein intake variability indicates potential compliance or dietary adequacy concerns requiring dietitian follow-up.
- More than half of our patients require nutritional supplementation. This indicates a **moderate-to-high nutrition risk population**.
- Albumin (ALB) is relatively stable
 - NPCR suggests **some patients are not meeting optimal protein intake**
 - There is **hidden risk** despite average stability
- 👉 This means: **overall looks stable, but vulnerable subgroups exist.**