

Data gap filling using cloud-based distributed Markov Chain Cellular Automata framework for land use and land cover change analysis

Hai Lan ^{1,2}, Kathleen Stewart ¹, Zongyao Sha ³, Shujuan Chang ⁴, and Yichun Xie ^{5,*}

¹ Department of Geographical Sciences, Univ. of Maryland, College Park, MD, USA

² NSF Spatiotemporal Innovation Center, George Mason Univ., Fairfax, VA, USA

³ School of Remote Sensing & Information Engineering, Wuhan Univ., Wuhan, China

⁴ Inner Mongolian Forestry & Rangeland Monitoring and Planning Institute, Hohhot, China

⁵ Department of Geography and Geology, Eastern Michigan Univ., Ypsilanti, MI, USA

* Correspondence: yxie@emich.edu; Tel.: +1 734-487-7588

Supplementary Figures and Tables




Figure S1. Data gaps in IMAR on August 2016

The portions in yellow are cloud covered areas during August 2016 that correspond to where data gaps exist in IMAR during this period. There are totally ~ 1.28 billion (1281956192) pixels/cells per band in this study area. ~0.49 billion (492899116) pixels are covered by clouds during this month, i.e., the cloud coverage is over 38.45%. In the northern part of the region, where cloud coverage is above 85%, clouds are a major obstacle that potentially impedes LULC research in this region.




Figure S2. Six clusters generated by county-based multivariate geographic k-medoids clustering analysis

Since Cluster 2 (in black color) is a single banner within Cluster 5 (in blue color), the final number of clusters used in the analysis is 5.

Table S1. Transition Matrix

	SSS	TMS	TS	TDS	TSD	TD	LM	MM	MA	WA	NG
TMS	1	71.87	8.03	N/A	N/A	N/A	2.12	0.34	0.09	0	17.54
	2	66.09	0	N/A	N/A	N/A	2.33	0.87	0.02	0.07	30.62
	3	58.59	22.35	0	0	N/A	3.01	0.53	0	0.17	15.36
	4	9.66	48.6	0	N/A	N/A	0	0	0	0.5	41.24
	5	16.91	76.58	0	0.35	0	0	0	0	0	6.16
TS	1	9.97	79.16	N/A	N/A	N/A	10.7	0	0.09	0.08	0
	2	0	65.43	N/A	N/A	N/A	3.75	0	0	0.26	30.57
	3	10.34	34.57	22.63	0	N/A	14.66	0.01	0	0.4	17.39
	4	0.24	60.14	0.03	N/A	N/A	0	0	0	2.11	37.48
	5	0	73.37	11.71	0	0	0.27	0	0	0.98	13.66
TDS	1	N/A									
	2	N/A									
	3	0	22.76	67.73	7.03	N/A	2.48	0	0	0	0
	4	0	10.65	86.34	N/A	N/A	0	0	0	0.32	2.69
	5	0	28.95	42.62	15.28	1.92	3.71	0.01	0	0.24	7.26
TSD	1	N/A									
	2	N/A									
	3	0.15	0	33.17	63.43	N/A	3	0	0	0	0.23
	4	N/A									
	5	0	0	21.89	52.12	25.63	0	0.04	0	0.32	0
TD	1	N/A									
	2	N/A									
	3	N/A									
	4	N/A									
	5	0	0	0	22.05	31.87	11.45	0	0.58	0	34.05
LM	1	1.01	4.51	N/A	N/A	N/A	54.59	1.75	1.18	2.31	34.65
	2	1.87	9.23	N/A	N/A	N/A	44.93	0	0.25	5.28	38.44
	3	6.62	29.2	2.13	2.46	N/A	50.11	0	0.1	3.24	6.14
	4	0	6.79	0	N/A	N/A	17.99	0	0	11.03	64.19
	5	0	7.42	16.44	4.41	11.44	30.38	0	0.21	5.99	23.71
MM	1	22.28	0	N/A	N/A	N/A	3.97	55.88	0	0	17.87
	2	41.08	0	N/A	N/A	N/A	2.83	31.27	0	0	24.82
	3	64.1	0	0.6	0.06	N/A	0.23	17.77	0.01	0	17.22
	4	27.8	0.82	0	N/A	N/A	0.03	69.52	0	0	1.84
	5	0	0	0	0	0	0	74.55	0	0	25.45
MA	1	2.48	3.2	N/A	N/A	N/A	47.45	0	43.31	3.56	0
	2	6.31	0.35	N/A	N/A	N/A	37.12	0	20.75	0	35.47
	3	2.83	0	0	0	N/A	96.91	0	0	0.26	0
	4	0	0	0	N/A	N/A	56.7	0	0	41.88	1.42
	5	0	0	0	0.8	0	0	0	35.23	59.18	4.79
WA	1	0	2.22	N/A	N/A	N/A	19.72	0	0.65	77.41	0
	2	0	0	N/A	N/A	N/A	22.13	0	1.15	44.55	32.17
	3	0	0.93	0	0	N/A	33.11	0	0.12	65.83	0
	4	0	0	0	N/A	N/A	37.8	0	0	47.51	14.68
	5	0	1.51	4.13	0	45.03	24.73	0	0.87	23.74	0
NG	1	9.9	0	N/A	N/A	N/A	32.12	4.54	0	0.71	52.73
	2	18.12	16.43	N/A	N/A	N/A	3.96	0.19	0	1.5	59.81
	3	18.07	29.78	0	0	N/A	0.59	0.66	0.02	1.33	49.55
	4	0.19	14.34	0.01	N/A	N/A	1.38	0.01	0.13	1.79	82.14
	5	0.11	13	12.32	0.34	31.96	6.56	0.05	0	2.41	33.25

The transition matrix for the ten LULC classes for the IMAR region was calculated (ST. 2). In this table, the diagonal values for each sub-simulation space (SSS) represented the probability that certain LULC classes that did not change during the training period. Other values in this table show the probability that certain LULC classes were changed to other types of LULC classes. The N/A value in this table represented the case where there was no such class in this SSS. During the two paired training periods 2000 to 2010, and 2010 to 2016, the most stable LULC classes were water area (WA) and non-grassland area (NG). They were considered as Boolean types of constrain factors, which were expected to be fixed and not change during the simulation. Other types of LULC classes can be changed into those two classes given a certain probability. However, in reality, it was quite rare that grassland classes changed into water areas. On other hand, rapid increases in human activities resulted in grassland classes being changed into NG classes, especially to farmland and human land use (LU). For example, cities and roads were expanded rapidly in IMAR. Over grazing also happened in some areas during past few decades. All those cases may lead to grassland degradation or even the disappearance of grassland especially in some sensitive areas. By assessing this transition matrix, we found that Temperate Meadow-steppe (TMS), temperate steppe (TS), and Lowland Meadow (LM) were changed into NG with a relatively high probability (typically over 30%) especially in SSS 2 and SSS 4, which were high population density and high livestock density regions respectively. We also found that some transitions happened between neighboring types of grassland classes. For example, TMS class showed a probability of approximately 76.58% to change into TS in SSS 5, which is a very dry area. In other words, TMS class areas that require relatively high moisture environments may be less likely to be stay as dry areas over time.

The transition matrix represented the analysis results from a mathematical perspective. During the simulation process, the results that were closer to an actual change mechanism were generated by integrating multi-types of impact factors, weighting each of them with different functions and assigning different neighborhood weights.

(a) Overall accuracy assessment without using sub-region strategy with 2 years data										
	TMS	TS	TDS	TSD	TD	LM	MM	MA	WA	NG
Producer's Accuracy	79.30	77.65	73.14	74.11	82.35	81.87	76.22	29.79	40.91	87.37
User's Accuracy	85.59	81.67	90.30	81.92	92.17	74.40	77.86	56.00	26.73	77.44
Overall Accuracy	81.23									
Kappa	0.76									
(b) Overall accuracy assessment using sub-region strategy with 2 years data										
	TMS	TS	TDS	TSD	TD	LM	MM	MA	WA	NG
Producer's Accuracy	84.02	87.10	80.59	80.81	83.37	83.38	82.35	38.10	42.75	88.30
User's Accuracy	88.61	85.01	92.59	83.48	92.14	74.24	80.26	39.02	26.79	85.00
Overall Accuracy	85.23									
Kappa	0.81									
(c) Overall accuracy assessment without using sub-region strategy with 3 years data										
	TMS	TS	TDS	TSD	TD	LM	MM	MA	WA	NG
Producer's Accuracy	81.10	82.19	76.76	73.85	83.78	85.65	81.12	38.10	46.28	89.11
User's Accuracy	88.25	82.21	92.25	82.20	92.28	75.25	78.91	59.26	27.45	83.11
Overall Accuracy	83.97									
Kappa	0.79									
(d) Overall accuracy assessment by using sub-region strategy with 3 years data										
	TMS	TS	TDS	TSD	TD	LM	MM	MA	WA	NG
Producer's Accuracy	87.20	91.88	89.71	89.23	89.51	84.14	86.67	94.12	59.76	85.93
User's Accuracy	89.04	89.13	93.20	84.89	98.05	77.79	85.53	28.07	23.67	86.89
Overall Accuracy	88.16									
Kappa	0.85									

Table S3. Sub-regions accuracy assessment

		TMS	TS	TDS	TSD	TD	LM	MM	MA	WA	NG
SSS 1	Producer's Accuracy	85.61	98.81	N/A	N/A	N/A	85.71	82.76	100	55.00	96.47
	User's Accuracy	93.70	99.10	N/A	N/A	N/A	87.64	96.00	100	100	93.57
	Overall Accuracy	93.82									
	Kappa	0.89									
SSS 2	Producer's Accuracy	81.30	74.88	N/A	N/A	N/A	70.87	100	100	31.25	88.41
	User's Accuracy	81.63	80.26	N/A	N/A	N/A	76.84	80.0	90.0	41.67	85.74
	Overall Accuracy	83.50									
	Kappa	0.71									
SSS 3	Producer's Accuracy	93.71	93.55	98.66	95.74	N/A	78.88	100	100	54.55	68.59
	User's Accuracy	90.24	94.44	99.66	100	N/A	92.56	100	90.0	54.55	65.52
	Overall Accuracy	89.78									
	Kappa	0.86									
SSS 4	Producer's Accuracy	90.91	71.54	100	N/A	N/A	30.77	90.91	100	15.0	87.34
	User's Accuracy	100	69.96	90.0	N/A	N/A	80.0	100	100	30.0	84.78
	Overall Accuracy	80.84									
	Kappa	0.60									
SSS 5	Producer's Accuracy	100	72.12	95.42	95.43	98.39	79.17	100	90.91	40.0	78.90
	User's Accuracy	20.0	84.40	88.56	99.40	95.10	75.0	70.0	100	54.55	83.88
	Overall Accuracy	89.74									
	Kappa	0.86									