

Unravelling the time-course of listener adaptation to an unfamiliar talker

Maryann Tan^{1, 2} & T. Florian Jaeger^{2,3}

¹ Centre for Research on Bilingualism, University of Stockholm

² Brain and Cognitive Sciences, University of Rochester

³ Computer Science, University of Rochester

Author Note

We are grateful to ### ommitted for review ###

Correspondence concerning this article should be addressed to Maryann Tan, Department of Bilingualism, Stockholm University, Sweden. E-mail: maryann.tan@biling.su.se

10 Abstract

11 We investigate constraints on adaptive speech perception during the initial encounters with
12 unfamiliar speech patterns. Such adaptive changes are now considered important to spoken
13 language understanding, overcoming substantial cross-talker variability in the realization of
14 speech categories. We present evidence from a novel incremental exposure-test paradigm to assess
15 how previously experienced cross-talker variability guides (and thus constrains) listeners'
16 adaptation. Specifically, we ask adaptation is constrained weakly—slower and sublinear, but
17 continued adaptation with increasing exposure—or strongly—adaptation only up to a point, after
18 which additional exposure has no benefits (at least not prior to, e.g., sleep). The results
19 contribute to a proposed theoretical distinction between two hypotheses about the mechanisms
20 underlying the initial moments of adaptation, model learning vs. model selection.

21 *Keywords:* speech perception; adaptation; incremental changes; distributional learning

22 Word count: X

Unravelling the time-course of listener adaptation to an unfamiliar talker

1 TO-DO

1.1 Highest priority

- MARYANN
- Figure out why slopes aren't identical across conditions under our current way of averaging exposure across all participants.
- Try an add line to table 2 to separate the unlearning hypothesis from the others (low priority). Add +40 vs baseline sub-heading

1.2 Medium priority

- MARYANN
- Fix a lot of the outstanding XXXes. Fill in the references – in library.bib
- FLORIAN
- think about table 1 and 2: how to change the wording on tables to actually refer to intercepts rather than PSEs or change the figures? Changing current representations of analyses to improve intuitive-ity.
- write overview of results
- restructure results presentation.
- write SI sections with proofs

1.2.1 Lower Priority

- MARYANN
- Combine data from exposure and test, use all together instead of coding block, code trial and code it as a smooth. That means using GAMM – that may require taking lapse (try it

first without lapses because the GAMM takes care of the lapse. The RE will be expressed differently. It has to follow the GAMM syntax.) The primary thing we want to smooth over is “block”, but could theoretically smooth over VOT and Block.

- Florian

- compare IBBU predictions over blocks with human behavioural data

1.3 To do later

- Everyone: Eat ice-cream and perhaps have a beer.

1 Introduction

One of the hallmarks of human speech perception is its adaptivity. Listeners' interpretation of acoustic input can change within minutes of exposure to an unfamiliar talker, supporting robust speech recognition across talkers (Bradlow & Bent, 2008; Clarke & Garrett, 2004; Xie, Liu, & Jaeger, 2021; Xie, Weatherholtz, et al., 2018). Recent reviews have identified distributional learning of marginal cue statistics ('normalization,' Apfelbaum & McMurray, 2015; McMurray & Jongman, 2011) or the statistics of cue-to-category mappings as an important mechanism affording this adaptivity ('representational learning,' Clayards, Tanenhaus, Aslin, & Jacobs, 2008; Davis & Sohoglu, 2020; Idemaru & Holt, 2011; Kleinschmidt & Jaeger, 2015; for review, Schertz & Clare, 2020; Xie, Jaeger, & Kurumada, 2023). This hypothesis has gained considerable influence over the past decade, with findings that changes in listener perception are qualitatively predicted by the statistics of exposure stimuli (Bejjanki, Beck, Lu, & Pouget, 2011; Clayards et al., 2008; Idemaru & Holt, 2020; Kleinschmidt & Jaeger, 2012; Munson, 2011; Nixon, Rij, Mok, Baayen, & Chen, 2016; Tan, Xie, & Jaeger, 2021; Theodore & Monto, 2019; for important caveats, see Harmon, Idemaru, & Kapatsinski, 2019).

More recently the means by which to test and differentiate between closely related hypotheses founded on distributional learning have been facilitated by the formalisation of speech perception as rational inference with Bayesian belief-updating (Ideal Adaptor framework; Kleinschmidt and Jaeger (2015)). In an ideal adaptor account speech sounds are represented as distributions of acoustic cues under a given linguistic category or context. At any given moment a listener holds prior beliefs about cue distributions formed over his or her long-term experience with talkers. Because cue distributions differ across talkers and contexts the listener infers the intended category of a novel talker by integrating prior beliefs about cue-category distributions with the present talker's input. How quickly a listener adapts to the speech of the talker depends on the degree of confidence the listener has about prior cue distributions. A greater confidence in priors requires a higher amount of evidence from the talker in order for the listener to update his beliefs about the cue distributions such that it aligns with that of the talker's. Conversely, low confidence in prior distributions would require less evidence for adaptation to happen. On these principles various learning models can be built to simulate the amount and manner of exposure to

81 talker inputs, allowing specific predictions of perceptual changes from each additional piece of
82 input under different parametric assumptions (Hitczenko & Feldman, 2016 for examples;
83 Kleinschmidt & Jaeger, 2015; Xie et al., 2023).

84 A characteristic prediction in an ideal adaptor account of adaptation is that one should be
85 able to observe graded (or incremental) outcomes in line with the distribution of the acoustic
86 input as well as the frequency of the input (amount of evidence). Recently Cummings and
87 Theodore (2023) sought to test this core tenet with a series of perceptual recalibration
88 experiments. Listeners were exposed to an ambiguous sound mid-way between two categories, /s/
89 and /sh/ either in an /s/-biasing or /sh/-biasing word contexts. They were then tested on their
90 categorisation behaviour on a continuum of ambiguous /s-/sh/ sounds. Depending on the
91 context in which listeners heard the ambiguous sound, they will tend to categorise the sounds
92 more often in the direction of the biasing context. By manipulating the number of times an
93 ambiguous sound between /s/ and /sh/ was heard between participants and within each biasing
94 context (1, 4, 10 or 20 occurrences) they showed that the size of the putative perceptual
95 recalibration effect correlated with the frequency of exposure. This kind of statistically modulated
96 change is rarely demonstrated in perceptual recalibration not least because of how those studies
97 are typically designed. It is more common to dispense exposure in one block followed by a block
98 of test trials because the effect of interest is simply whether the average categorisation functions
99 of the two groups diverge or not post-exposure. While Cummings and Theodore (2023) found
100 graded effects between-participants, it should in principle be possible to detect graded change
101 within participants as well had they been given the full extent of critical trials but tested after
102 each incremental input. Such a build-up in perceptual recalibration was in fact tested in
103 Vroomen, Linden, De Gelder, and Bertelson (2007). Participants were exposed to an ambiguous
104 sound between /aba/ and /ada/ paired with videos of a person either clearly articulating “aba” or
105 “ada”. Interspersed between the first 64 exposure trials of the experiment were test trials to check
106 for incremental build-up in the recalibration effect. In their modelling of this study Kleinschmidt
107 and Jaeger (2011) found the ideal adaptor to fit the data well indicating that listeners integrate
108 each observed piece of evidence with their priors and updated their beliefs in a way that was
109 quantitatively predicted by the model.

So far we have discussed findings of gradient changes in perceptual recalibration studies that repeatedly expose listeners to tokens made from a single synthesised ambiguous token, or in the case of (**cummings-theodore?**), tokens that are a blend of the cues from the /s/ and /sh/ utterances of the critical exposure words. The main manipulation is the number of exposures to the ambiguous sounds with the predicted effect being the quantity of difference between groups or in the case of incremental testing, the quantity of difference between trials.

In the distributional learning experimental paradigm clusters of cue values are constructed and centered around specific mean values with specific variances along a single acoustic dimension to simulate within talker variability. These cues are mapped onto two adjacent sound categories and played to listeners in the context of minimal pair words who are tasked to categorise what they believe they heard. It is the the locations of the means of these distributions or category variances that are manipulated between groups. Through exposure to these distributions, supervised or unsupervised, and with sufficient numbers of trials listeners are expected to learn the talker's distributions and therefore show categorization patterns that correspond to the exposure distribution. Kleinschmidt and Jaeger (2016) for instance exposed L1-US English listeners to recordings of /b/-/p/ minimal pair words like *beach* and *peach* that were acoustically manipulated. Separate groups of listeners were exposed to distributions of voice onset times (VOTs)—the primary cue distinguishing words like *beach* and *peach*—that were shifted by up to +30 ms, relative to what one might expect from a ‘typical’ talker (Figure 1A). In line with the distributional learning hypothesis, listeners’ category boundary or point of subjective equality (PSE)—i.e., the VOT for which listeners are equally likely to respond “b” or “p”—shifted in the same direction as the exposure distribution (Figure 1B). Also in line with the distributional learning hypothesis, these shifts were larger the further the exposure distributions were shifted. An incremental Bayesian belief updating model was fit to the data which showed that within each exposure group listeners gradually increased the shift of their categorisations as they heard more trials (Kleinschmidt, 2020).

This initial finding of incremental belief updating within the distributional learning paradigm while noteworthy, is not without its limitations. Distributional learning studies have not typically been set up as exposure-test configurations. All exposure trials are in a sense test

139 trials as well. This means that the estimated categorisation function is fit over a number of trials
140 decided on by the researcher – for e.g. Kleinschmidt and Jaeger (2016) apportioned their 222
141 trials into 6 segments. Accordingly, past studies have not implemented standard test trials across
142 groups.

143 The present study aims to build on the pioneering work of Clayards et al. (2008);
144 Kleinschmidt and Jaeger (2016); Theodore and Monto (2019); Kleinschmidt (2020) with some
145 design innovations that we believe affords a productive test of the core claims of the ideal
146 adaptor. We aim to replicate the findings of Kleinschmidt and Jaeger (2016) by manipulating
147 exposure distributions between groups while empirically investigating incremental learning by
148 introducing a novel test-exposure-test design. Another novelty this study possibly brings is a
149 document of how soon, from the onset of exposure, does the distributional learning effect emerge.
150 This is something that remains opaque in previous studies because of their lack of test blocks.
151 Given the substantial amount of evidence that adaptation takes place rapidly (e.g. 5 mins from
152 exposure in L2 accent adaptation; 4 - 10 trials in lexically guided perceptual recalibration) we
153 might expect listeners to also show learning effects very early on in an experiment. On the other
154 hand, it is possible that distributional learning effects may show up later than one would expect
155 in perceptual recalibration since listeners have a presumably more challenging task of inferring
156 the means of two categories over a range of cues.

157 In experimental work researchers often have to consider the generalizability of their results
158 which leads to questions about ecological validity. There is a trade-off between ecological validity
159 of experimental design and the desired degree of control over the variables. Questions about
160 ecological validity of prior work in distributional learning pertains to 2 features. First, the stimuli
161 which were generated with a synthesiser, had an obvious machine-like quality (Clayards et al.,
162 2008; Kleinschmidt & Jaeger, 2016). Secondly, the pairs of distributions of voiced and voiceless
163 categories were always identical in their variances [see also theodore-monto2019] which adds to
164 the artificiality of the experiment. In our design description we show how we can begin to
165 improve on these features through the stimuli and the setting of exposure conditions.

166 **END OF INTRODUCTION

167 For example, influential models of adaptive speech perception predict proportional, rather




Figure 1. Design and results of Kleinschmidt and Jaeger (2016) replotted. **Panel A:** Different groups of participants were exposed to different shifts in the mean VOT of /b/ and /p/. **Panel B:** categorization functions of individual participants depending on the exposure condition (shift in VOT means of /b/ and /p/). For reference, the black dashed line shows the categorization function of the 0-shift condition. The colored dashed lines shows the categorization function expected for an ideal observer that has fully learned the exposure distributions. **Panel C:** Mean and 95% CI of participants' points of subjective equality (PSEs), relative to the PSE of the ideal observers.

than sublinear, shifts (for proof, see SI ??). This is the case both for incremental Bayesian belief-updating model (Kleinschmidt & Jaeger, 2011) and general purpose normalization accounts (McMurray & Jongman, 2011)—models that have been found to explain listeners' behavior well in experiments with less substantial changes in exposure. There are, however, proposals that can accommodate this finding. Some proposals distinguish between two types of mechanisms that might underlie representational changes, *model learning* and *model selection* (Xie, Weatherholtz, et al., 2018, p. 229). The former refers to the learning of a new category representations—for example, learning a new generative model for the talker (Kleinschmidt & Jaeger, 2015, pt. II) or storage of new talker-specific exemplars (Johnson, 1997; Sumner, 2011). Xie and colleagues hypothesized that this process might be much slower than is often assumed in the literature,

potentially requiring multiple days of exposure and memory consolidation during sleep (see also Fenn & Hambrick, 2013; Tamminen, Davis, Merckx, & Rastle, 2012; Xie, Earle, & Myers, 2018). Rapid adaptation that occurs within minutes of exposure might instead be achieved by selecting between *existing* talker-specific representations that were learned from previous speech input—e.g., previously learned talker-specific generative models (see mixture model in Kleinschmidt & Jaeger, 2015, pp. 180–181) or previously stored exemplars from other talkers (Johnson, 1997). Model learning and model selection both offer explanations for the sublinear effects observed in Kleinschmidt and Jaeger (2016). But they suggest different predictions for the evolution of this effect over the course of exposure.

Under the hypothesis of model learning, sublinear shifts in PSEs can be explained by assuming a hierarchical prior over talker-specific generative models ($p(\Theta)$ in Kleinschmidt & Jaeger, 2015, p. 180). This prior would ‘shrink’ adaptation towards listeners’ priors—similar to the effect of random by-subject or by-item effects in generalized linear mixed-effect models, which shrink group-level effect estimates towards the population mean of the data (Baayen, Davidson, & Bates, 2008). Critically, as long as these priors attribute non-zero probability to even extreme shifts (e.g., the type of Gaussian prior used in mixed-effects models), this predicts listeners’ PSEs will continue to change with increasing exposure until they have converged against the PSE that is ideal for the exposure statistics. In contrast, the hypothesis of model selection predicts that rapid adaptation is more strictly constrained by previous experience: listeners can only adapt their categorization functions up to a point that corresponds to (a mixture of) previously learned talker-specific generative models. This would imply that at least the earliest moments of adaptation are subject to a hard limit (Figure 2): exposure helps listeners to adapt their interpretation to more closely aligned with the statistics of the input, but only to a certain point.

The present study employs a novel incremental exposure-test paradigm to address two questions. We test whether the sublinear effects of exposure observed in recent work replicate for exposure that (somewhat) more closely resembles the type of speech input listeners receive on a daily basis. And, we evaluate the predictions of the model learning and selection hypotheses against human perception. We take this question to be of interest beyond the specific hypotheses we contrast: whether there are hard limits to the benefits of exposure to unfamiliar speech




Figure 2. Contrasting predictions of model learning and model selection hypotheses about the incremental effects of exposure on listeners’ categorization function. Both hypothesis predict incremental adaptation towards the statistics of the input, as well as constraints on this adaptation. The two hypotheses differ, however, in that model selection predicts a hard limit on how far listeners’ can adapt during initial encounters with an unfamiliar talker.

patterns ultimately has consequences for education and medical treatment.

All data and code for this article can be downloaded from <https://osf.io/hxyc4/>. The article is written in R markdown, allowing readers to replicate our analyses with the press of a button using freely available software (R, R Core Team, 2022; RStudio Team, 2020), while changing any of the parameters of our models (see SI, ??).

2 Experiment

We revise the standard paradigm used to investigate distributional learning in speech perception. Previous work has employed ‘batch testing’ designs, in which changes in categorization responses are assessed only after extended exposure to hundreds of trials or by averaging over extended exposure (e.g., Clayards et al., 2008; Harmon et al., 2019; Idemaru & Holt, 2011, 2020; Kleinschmidt & Jaeger, 2016; Munson, 2011; Nixon et al., 2016; Theodore & Monto, 2019). These designs are well-suited to investigate cumulative effects of exposure but are less so to identify constraints on rapidly unfolding incremental adaptation. To be able to detect both incremental and cumulative effects of exposure, within and across exposure conditions, we employed the repeated exposure-test design shown in Figure 3.

The use of test blocks that repeat the same stimuli across blocks and exposure conditions deviates from previous work (Clayards et al., 2008; Kleinschmidt & Jaeger, 2016; Theodore & Monto, 2019). This design feature allowed us to assess how increasing exposure affects listeners’




Figure 3. Exposure-test design of the experiment. Test blocks presented identical stimuli within and across conditions

perception without making strong assumptions about the nature of these changes (e.g., linear changes across trials). We kept test blocks short for two reasons. First, previous work has found that repeated testing over uniform test continua can reduce or undo the effects of informative exposure (Cummings & Theodore, 2023; Liu & Jaeger, 2018, 2019). Second, since we held test stimuli constant across exposure conditions, the distribution—and thus the relative unexpectedness—of test stimuli differed to different degrees from the three exposure distributions. By keeping tests short relative exposure (12 vs. 48 trials), we aimed to minimize the influence of test trials on adaptation. The final three test blocks were intended to ameliorate the potential risks of this novel design: in case adaptation remains stable despite repeated testing, those additional test blocks were meant to provide additional statistical power to detect the effects of cumulative exposure.

We also adjusted the standard distributional learning paradigm to increase the ecological validity of the exposure and test stimuli. The pioneering works that inspired the present study employed speech stimuli that did not exhibit the natural correlations between different acoustic-phonetic cues that characterise human speech, and that were clearly identifiable as robotic speech (Clayards et al., 2008; Kleinschmidt & Jaeger, 2016). These studies also followed the majority of research on distributional learning in language (e.g., Maye, Werker, & Gerken, 2002; Pajak & Levy, 2012) and *designed* rather than *sampled* the exposure distributions. As a consequence, exposure distributions in these experiments tend to be symmetrically balanced around the category means—unlike in everyday speech input. Indeed, all of the works we follow

here further used categories with *identical* variances (e.g., identical variance along VOT for /b/ and /p/, Clayards et al., 2008; Kleinschmidt & Jaeger, 2016; or /g/ and /k/, Theodore & Monto, 2019). This, too, is highly atypical for everyday speech input (Chodroff & Wilson, 2018; Lisker & Abramson, 1964). The present study takes several modest steps to ameliorate these issues.

2.1 Methods

2.1.1 Participants

We recruited 126 participants from the Prolific crowdsourcing platform. We used Prolific’s pre-screening to limit the experiment to participants (1) of US nationality, (2) who reported to be English speaking monolinguals, and (3) had not previously participated in any experiment from our lab on Prolific. Prior to the start of the experiment, participants had to confirm that they (4) had spent the first 10 years of their life in the US, (5) were in a quiet place and free from distractions, and (6) wore in-ear or over-the-ears headphones that cost at least \$15. An additional 115 participants loaded the experiment but did not start or complete it.¹

Participants took an average of 31.6 minutes to complete the experiment (SD = 20 minutes) and were remunerated \$8.00/hour. An optional post-experiment survey recorded participant demographics using NIH prescribed categories, including participant sex (59 = female, 60 = male, 3 = NA), age (mean = NA years; 95% quantiles = 20-62.1 years), race (6 = Black, 31 = White, 85 = NA), and ethnicity (6 = Hispanic, 113 = Non-Hispanic, 3 = NA).

Participants’ responses were collected via Javascript developed by the Human Language Processing Lab at the University of Rochester (**JSEXP?**) and stored via Proliferate developed at, and hosted by, the ALPs lab at Stanford University (Schuster, S, 2020).

2.1.2 Materials

We recorded 8 tokens each of four minimal word pairs (*dill/till*, *dim/tim*, *din/tin*, and *dip/tip*) from a 23-year-old, female L1-US English talker from New Hampshire, judged to have a “general

¹ Unlike in lab-based experiments, for which participants’ right to stop the experiment at any point is costly (both in terms of physical effort and perceived social cost), exercising this right in web-based experiments is essentially cost free—in particular, if exercised early in the experiment.

American” accent. In addition to these critical minimal pairs we also recorded three words that did not contain any stop consonant sounds (“flare”, “share”, and “rare”). These word recordings were used for catch trials. Stimulus intensity was normalized to 70 dB sound pressure level for all recordings.

The critical minimal pair recordings were used to create four VOT continua using a script (Winn, 2020) in Praat (Boersma & Weenink, 2022). This approach resulted in continuum steps that sound natural (unlike the highly robotic-sounding stimuli employed in Clayards et al., 2008; Kleinschmidt & Jaeger, 2016). A post-experiment survey asked participants: “*Did you notice anything in particular about how the speaker pronounced the different words (e.g. till, dill, etc.)?*” No participant reported that the stimuli sounded unnatural. The procedure also maintained the natural correlations between the most important cues to word-initial stop-voicing in L1-US English (VOT, F0, and vowel duration). Specifically, the F0 at vowel onset of each stimulus was set to respect the linear relation with VOT observed in the original recordings of the talker. The duration of the vowel was set to follow the natural trade-off relation with VOT (Allen & Miller, 1999). Further details on the recording and resynthesis procedure are provided in the supplementary information (SI, ??).

The VOTs generated for each continuum ranged from -100 to +130 ms in 5 ms steps.² A norming experiment (N = 24 participants) reported in the SI (??) was used to select the three minimal pair continua that elicited the most similar categorization responses (*dill-till*, *din-tin*, and *dip-tip*). These three continua were used to create the exposure conditions shown in Figure 3.

2.1.3 Procedure

At the start of the experiment, participants acknowledged that they met all requirements and provided consent, as per the Research Subjects Review Board of the University of Rochester.

² We follow previous work (Kleinschmidt, 2020; Lisker & Abramson, 1964) and refer to pre-voicing as negative VOTs though we note that pre-voicing is perhaps better conceived of as a separate phonetic feature (for discussion, see REF?). Estimates of the proportion of voiced stops produced with pre-voicing in L1-US English vary substantially between studies (between 20% and 57%) (Dmitrieva, Llanos, Shultz, & Francis, 2015; e.g. Lisker & Abramson, 1967; Smith, 1978; Westbury, 1979). Because pre-voicing is not regarded as a phonemic determinant of English, some studies either discard such data or ignore them altogether (e.g. Zue (1976); Klatt (1975); Chodroff and Wilson (2017)). In some studies that do report pre-voicing, the majority of the tokens were attributed to a minority of talkers (Flege & Brown Jr, 1982; e.g. Lisker & Abramson, 1967). Although speakers tend to prefer one type of production over the other they do not typically use one type exclusively (Docherty, 2011).

Participants also had to pass a headphone test (Woods, Siegel, Traer, & McDermott, 2017), and were instructed to not change the volume throughout the experiment. Following instructions, participants completed 234 two-alternative forced-choice categorization trials (Figure 4). Participants were instructed that they would hear a female talker say a single word on each trial, and were asked to select which word they heard. Participants were asked to listen carefully and answer as quickly and as accurately as possible. They were also alerted to the fact that the recordings were subtly different and therefore may sound repetitive.




Figure 4. Example trial display. When the green button turned bright green, participants had to click on it to play the recording.

Unbeknownst to participants, the 234 trials were split into exposure (54 trials each) and test blocks (12 trials each). Participants were given the opportunity to take breaks after every 60 trials, which was always during an exposure block. Finally, participants completed an exit survey and an optional demographics survey.

Test blocks. The experiment started with a test block. Test blocks were identical within and across conditions, always including 12 minimal pair trials assessing participants' categorization at 12 different VOTs (-5, 5, 15, 25, 30, 35, 40, 45, 50, 55, 65, 70 ms). A uniform distribution over VOTs was chosen to maximize the statistical power to determine participants' categorization function. The assignment of VOTs to minimal pair continua was randomized for each participant, while counter-balancing it within and across test blocks. Each minimal pair appear equally often within each test block (four times), and each minimal pair appear with each VOT equally often (twice) across all six test blocks (and no more than once per test block).

Each trial started with a dark-shaded green fixation dot being displayed. At 500ms from trial onset, two minimal pair words appeared on the screen, as shown in Figure 4. At 1000ms

from trial onset, the fixation dot would turn bright green and participants had to click on the dot to play the recording. This was meant to reduce trial-to-trial correlations by resetting the mouse pointer to the center of the screen at the start of each trial. Participants responded by clicking on the word they heard and the next trial would begin.

Exposure blocks. Each exposure block consisted of 24 /d/ and 24 /t/ trials, as well as 6 catch trials that served as a check on participant attention throughout the experiment (2 instances for each of three combinations of the three catch recordings). With a total of 144 trials, exposure was substantially shorter than in similar previous experiments (cf. 228 trials in Clayards et al., 2008; 222 trials in Kleinschmidt, 2020; 2 x 236 trials, Theodore & Monto, 2019; 456 trials, Nixon et al., 2016).

The distribution of VOTs across the 48 /d/-/t/ trials depended on the exposure condition. Specifically, we first created a *baseline* condition. Although not critical to the purpose of the experiment, we aimed for the VOT distribution in this condition to closely resemble participants' prior expectations for a 'typical' female talker of L1-US English (for details, see SI, ??). The mean and standard deviations for /d/ along VOT were set at 5 ms and 8.9 ms, respectively. The mean and standard deviations for /t/ were set at 50 ms and 16 ms, respectively. To create more realistic VOT distributions, we *sampled* from the intended VOT distribution (top row of Figure 5). This creates distributions that more closely resemble the type of distributional input listeners experience in everyday speech perception, deviating from previous work, which exposed listeners to highly unnatural fully symmetric samples (Clayards et al., 2008; Kleinschmidt, 2020; Kleinschmidt & Jaeger, 2016).

Half of the /d/ and half of the /t/ trials were labeled, the other half was unlabeled (paralleling one of the conditions in Kleinschmidt, Raizada, & Jaeger, 2015). Unlabeled trials were identical to test trials except that the distribution of VOTs across those trials was bimodal (rather than uniform), and determined by the exposure condition.³ Labeled trials instead

³ Previous studies have estimated changes in participants' categorization responses by analyzing responses on unlabeled exposure trials (e.g., Clayards et al., 2008; Kleinschmidt & Jaeger, 2016; Theodore & Monto, 2019). This approach compares responses across different values of acoustic-phonetic cues (since the exposure inputs differed by exposure condition), so that assumptions baked into the analysis approach (e.g., linearity along the acoustic-phonetic continuum) can potentially bias the results. Here we avoid this issue by holding test stimuli constant (see also Kleinschmidt, 2020, Experiment 4).

presented two response options with identical stop onsets (e.g., *din* and *dill*). This effectively labeled the input as belonging to the intended category (e.g., /d/).

Next, we created the two additional exposure conditions by shifting these VOT distributions by +10 or +40 ms (see Figure 5). This approach exposes participants to heterogeneous approximations of normally distributed VOTs for /d/ and /t/ that varied across blocks, while holding all aspects of the input constant across conditions except for the shift in VOT. The order of trials was randomized within each block and participant, with the constraint that no more than two catch trials would occur in a row. Participants were randomly assigned to one of 3 (exposure condition) x 3 (block order) x 2 (placement of response options) lists.




Figure 5. Histogram of voice onset times (VOTs) for each of the three exposure blocks A-C by trial type (/d/ or /t/, labeled or unlabeled) and exposure condition (baseline vs. +10 vs. +40). Each exposure block contained 12 labeled /d/, 12 labeled /t/, 12 unlabeled /d/, and 12 unlabeled /t/ trials, as well as 6 catch trials (not shown). Except for the shift in VOTs (+0, 10 or 40 ms VOT to each trial), the VOT distribution of these trials was identical across exposure conditions. The order of exposure blocks A-C was counter-balanced across participants using a Latin-square design.

2.1.4 Exclusions

Due to data transfer errors 4 participants' data were not stored and therefore excluded from analysis. We further excluded from analysis participants who committed more than 3 errors out of the 18 catch trials ($<83\%$ accuracy, $N = 1$), participants who committed more than 4 errors out of the 72 labelled trials ($<94\%$ accuracy, $N = 0$), participants with an average reaction time more than three standard deviations from the mean of the by-participant means ($N = 0$), participants who had atypical categorization functions at the start of the experiment ($N = 2$, see SI, ?? for details), and participants who reported not to have used headphones ($N = 0$). This left for analysis 17,136 exposure and 8,568 test observations from 119 participants (94% of total), evenly split across the three exposure conditions.

2.2 Results

We analyzed participants' categorization responses during exposure and test blocks in two separate Bayesian mixed-effects psychometric models, using *brms* (Bürkner, 2017) in R (R Core Team, 2022; RStudio Team, 2020, for details, see SI, ??). Psychometric models account for attentional lapses while estimating participants' categorization functions. Failing to account for attentional lapses—while commonplace in research on speech perception (but see Clayards et al., 2008; Kleinschmidt & Jaeger, 2016)—can lead to biased estimates of categorization boundaries (Prins, 2011; Wichmann & Hill, 2001). For the present experiment, however, lapse rates were negligible (0.8%, 95%-CI: 0.4 to 1.5%), and all results replicate in simple mixed-effects logistic regressions (Jaeger, 2008).

Each psychometric model regressed participants' categorization responses against the full factorial interaction of VOT, exposure condition, and block, while including the maximal random effect structure (see SI, ??). Figure 6 summarizes the results that we describe in more detail next. Panels A and B show participants' categorization responses during exposure and test blocks, along with the categorization function estimated from those responses via the mixed-effects psychometric models. These panels facilitate comparison between exposure conditions within each block. Panels C and D show the slope and point of subject equality (PSE)—i.e., the point at

which participants are equally likely to respond “d” and “t”—of the categorization function across blocks and conditions. These panels facilitate comparison across blocks within each exposure condition. Here we focus on the test blocks, which were identical within and across exposure conditions. Analyses of the exposure blocks are reported in the SI (??), and replicate all effects found in the test blocks.

We begin by presenting the overall effects, averaging across all test blocks. This part of our analysis matches previous work, which has focused on the overall effect of exposure across the entire experiment (‘batch tests,’ e.g., Clayards et al., 2008; Kleinschmidt & Jaeger, 2016; Nixon et al., 2016; Theodore & Monto, 2019) and/or during a single post-exposure test block (e.g., Kleinschmidt, 2020). Then we turn to the goals of this study—to characterize the incremental changes in participants’ categorization responses as a function of exposure and, in particular, to test 1) whether we replicate the sublinear effects of exposure observed in previous work under the ecologically more valid stimuli and distributions employed in the present work, and 2) whether we can begin to distinguish between the predictions of the model learning and selection hypotheses.

```
## [1] "VOT test mean: 35.8333333333333"
```

```
## [1] "VOT test mean: 35.8333333333333"
```

```
## [1] "VOT test mean: "
```

```
## [1] "VOT test mean: "
```




Figure 6. Summary of results. **Panel A:** Changes in listeners psychometric categorization functions as a function of exposure, from Test 1 to Test 4 with all intervening exposure blocks (only unlabeled trials were included in the analysis of exposure blocks since labeled trials provide no information about listeners' categorization function). Point ranges indicate the mean proportion of “t”-responses and their 95% bootstrapped CI. Lines and shaded intervals show the MAP predictions and 95% posterior CIs of a Bayesian mixed-effects psychometric model fit to participants' responses. **Panel B:** Same as Panel A but for the final three test blocks without intervening exposure. Test 4 is shown as part of both Panels A and B. **Panels C & D:** Changes across blocks in the slope and boundary (point-of-subjective-equality, PSE) of the categorization functions shown in Panels A-B. Point ranges represent the posterior medians and their 95% CI. Dashed reference lines show the intercepts and PSEs that naive (non-rational) learner would be expected to converge against after sufficient exposure (an ideal observer model that knows the exposure distributions). Percentage labels indicate the amount of shift

2.2.1 Does exposure affect participants' categorizations (averaging across all blocks)?

We first used the psychometric mixed-effects model to assess whether the exposure conditions had the expected effects across all test blocks *relative to each other*. Unsurprisingly, participants were more likely to respond “t” the larger the VOT ($\hat{\beta} = 15.09$, 90%–CI = [12.377, 17.625], $BF = Inf$, $p_{posterior} = 1$). Critically, exposure affects participants' categorization responses in the expected direction. Marginalizing across all blocks, participants in the +40 condition were less likely to respond “t” than participants in the +10 condition ($\hat{\beta} = -2.26$, 90%–CI = [-3.258, -1.228], $BF = 162.3$, $p_{posterior} = 0.994$) or the baseline condition ($\hat{\beta} = -3.08$, 90%–CI = [-4.403, -1.669], $BF = 215.2$, $p_{posterior} = 0.995$). There was also evidence—albeit less decisive—that participants in the +10 condition were less likely to respond “t” than participants in the baseline condition ($\hat{\beta} = -0.82$, 90%–CI = [-1.887, 0.282], $BF = 8.9$, $p_{posterior} = 0.899$). That is, the +10 and +40 conditions resulted in categorization functions that were shifted rightwards compared to the baseline condition, as also visible in Figures 6.

This replicates previous findings that exposure to changed VOT distributions changes listeners' categorization responses (for /b/-/p/: Clayards et al., 2008; Kleinschmidt, 2020; Kleinschmidt & Jaeger, 2016; for /g/-/k/, Theodore & Monto, 2019). Having established that exposure affected categorization, we turn to the questions of primary interest. Incremental changes in participants' categorization responses can be assessed from three mutually complementing perspectives. First, we compare how exposure affects listeners' categorization responses relative to other exposure conditions. This tests how early in the experiment differences between exposure conditions began to emerge. Second, we compare how exposure affects listeners' categorization responses within each condition relative to listeners' responses prior to any exposure. This assesses how the exposure conditions relate to participants' prior expectations. Most importantly, however, it tests the subtly different predictions of the model learning and selection hypotheses—whether changes in listeners' categorization responses are strongly constrained. Third and finally, we compare changes in listeners' responses to those expected from an ideal observer that has fully learned the exposure distributions. This tests whether the

sublinear effects observed in Kleinschmidt and Jaeger (2016) replicate in our repeated exposure-test paradigm with the improvements the present study makes to ecological validity.

2.2.2 Comparing across exposure conditions: How quickly does exposure begin to affect participants' responses?

Figure 6A suggests that differences between exposure conditions emerged early in the experiment: already in Test 2, listener's categorization functions seem to be shifted rightwards (larger PSEs) in the +40 condition compared to the +10 condition, and in the +10 condition compared to the baseline condition. This is confirmed by the Bayesian hypothesis tests summarized in Table 1. Prior to any exposure, during Test 1, participants' responses did not differ across exposure condition (all BFs > XXX). After exposure to only 24 /d/ and 24 /t/ stimuli, during Test 2, participants' responses differed between exposure conditions (BFs > 13.7). The difference between the +40 condition and the +10 or baseline condition kept increasing with exposure up to Test 4. Additional hypothesis tests in Table 2 show that the change from Test 1 to 2 was largest (BF = 57.82), followed by the change from Test 2 to 3 (BF = 10), with only minimal changes from Test 3 to 4 (BF = 1.68). Qualitatively paralleling the changes across blocks for the +40 condition, the change in the difference between the +10 and baseline conditions was largest from Test 1 to 2 (BF = 5.42), and then somewhat decreased from Test 2 to Test 4 (BFs < 1). The comparison across exposure conditions thus suggests that changes in listeners' categorization responses emerged quickly—indeed, they were present already *during* the first exposure block (see SI, ??)—but then leveled off. The comparison across exposure conditions also yields one result that is, at first blush, surprising: while the difference between the +10 and the baseline condition emerged already after the first exposure block, this difference *decreased*, rather than increased, with additional exposure from Test 2 to 3 (see second row of Table 2). We return to this effect below.

Tables 1 and 2 also reveal the consequences of repeated testing. The difference between exposure conditions decreased from Test 4 to 6 (BFs > 4.3; see also Figure 6B & D). On the final test block, the +10 condition did not differ any longer from the baseline condition. Only the differences between the +40 condition relative to the +10 and baseline conditions persisted, albeit substantially reduced compared to Test 4. This pattern of results replicates previous findings that

repeated testing over uniform test continua can undo the effects of exposure (Cummings & Theodore, 2023; Liu & Jaeger, 2018, 2019), and extends them from perceptual recalibration paradigms to distributional learning paradigms (see also Kleinschmidt, 2020). One important methodological consequence of these findings is that longer test phases do not necessarily increase the statistical power to detect effects of adaptation (unless analyses take the effects of repeated testing into account, as in the approach developed in Liu & Jaeger, 2018). Analyses that average across all test tokens—as remains the norm—are bound to systematically underestimate the adaptivity of human speech perception.

Table 1

When did exposure begin to affect participants' categorization responses? When, if ever, were these changes undone with repeated testing? This table summarizes the simple effects of the exposure conditions for each test block.

Hypothesis	Estimate	SE	90%-CI	BF	$p_{\text{posterior}}$
Test block 1 (pre-exposure)					
+10 vs. baseline = 0	-0.34	0.75	[-2.025, 1.437]	3.3	0.77
+40 vs. +10 = 0	0.25	0.73	[-1.338, 1.903]	3.7	0.79
+40 vs. baseline = 0	-0.08	0.91	[-2.124, 2.082]	4.8	0.83
Test block 2					
+10 vs. baseline	-1.45	0.88	[-2.933, 0.181]	13.7	0.93
+40 vs. +10	-2.08	0.99	[-3.824, -0.173]	24.3	0.96
+40 vs. baseline	-3.49	1.24	[-5.635, -1.072]	54.2	0.98
Test block 3					
+10 vs. baseline	-0.78	0.62	[-1.888, 0.364]	7.9	0.89
+40 vs. +10	-2.80	0.82	[-4.188, -1.113]	86.0	0.99
+40 vs. baseline	-3.56	0.97	[-5.202, -1.582]	110.1	0.99
Test block 4					
+10 vs. baseline	-0.88	0.85	[-2.36, 0.847]	4.8	0.83
+40 vs. +10	-3.32	0.89	[-4.883, -1.636]	128.0	0.99
+40 vs. baseline	-4.16	1.21	[-6.275, -1.882]	122.1	0.99
Test block 5 (no additional exposure)					
+10 vs. baseline	-1.33	0.71	[-2.556, -0.003]	19.1	0.95
+40 vs. +10	-2.38	0.86	[-3.893, -0.796]	65.1	0.98
+40 vs. baseline	-3.25	1.24	[-5.307, -0.923]	53.0	0.98
Test block 6 (no additional exposure)					
+10 vs. baseline	-0.22	0.72	[-1.485, 1.114]	1.6	0.62
+40 vs. +10	-1.70	0.79	[-3.078, -0.171]	25.0	0.96
+40 vs. baseline	-2.57	1.22	[-4.58, -0.191]	24.0	0.96

Table 2

Was there incremental change from test block 1 to 4? Did these changes dissipate with repeated testing from block 4 to 6? This table summarizes the interactions between exposure condition and block, whether the differences between exposure conditions changed from test block to test block.

Hypothesis	Estimate	SE	90%-CI	BF	$p_{posterior}$
Difference in +10 vs. baseline					
Block 1 to 2: increased Δ_{PSE}	-0.85	0.78	[-2.166, 0.632]	5.42	0.84
Block 2 to 3: increased Δ_{PSE}	0.34	0.77	[-1.144, 1.761]	0.48	0.32
Block 3 to 4: increased Δ_{PSE}	0.06	0.77	[-1.382, 1.532]	0.89	0.47
<i>Block 1 to 4: increased Δ_{PSE}</i>	-0.42	1.26	[-2.759, 1.963]	1.70	0.63
Block 4 to 5: decreased Δ_{PSE}	-0.33	0.60	[-1.43, 0.785]	0.41	0.29
Block 5 to 6: decreased Δ_{PSE}	1.03	0.65	[-0.234, 2.164]	11.95	0.92
<i>Block 4 to 6: decreased Δ_{PSE}</i>	0.70	0.82	[-0.896, 2.177]	3.83	0.79
Difference in +40 vs. +10					
Block 1 to 2: increased Δ_{PSE}	-2.36	0.89	[-3.811, -0.754]	57.82	0.98
Block 2 to 3: increased Δ_{PSE}	-1.16	0.83	[-2.592, 0.312]	10.00	0.91
Block 3 to 4: increased Δ_{PSE}	-0.27	0.82	[-1.694, 1.162]	1.68	0.63
<i>Block 1 to 4: increased Δ_{PSE}</i>	-3.78	1.22	[-5.865, -1.447]	84.11	0.99
Block 4 to 5: decreased Δ_{PSE}	1.14	0.77	[-0.244, 2.514]	11.38	0.92
Block 5 to 6: decreased Δ_{PSE}	0.45	0.77	[-0.985, 1.787]	2.58	0.72
<i>Block 4 to 6: decreased Δ_{PSE}</i>	1.59	1.00	[-0.3, 3.323]	12.68	0.93
Difference in +40 vs. baseline					
Block 1 to 2: increased Δ_{PSE}	-3.16	1.02	[-4.958, -1.185]	79.00	0.99
Block 2 to 3: increased Δ_{PSE}	-0.82	1.08	[-2.749, 1.145]	3.39	0.77
Block 3 to 4: increased Δ_{PSE}	-0.20	1.08	[-2.146, 1.741]	1.34	0.57
<i>Block 1 to 4: increased Δ_{PSE}</i>	-4.19	1.71	[-7.219, -0.93]	45.78	0.98
Block 4 to 5: decreased Δ_{PSE}	0.80	0.92	[-0.971, 2.493]	4.16	0.81
Block 5 to 6: decreased Δ_{PSE}	1.48	0.94	[-0.36, 3.117]	10.85	0.92
<i>Block 4 to 6: decreased Δ_{PSE}</i>	2.27	1.27	[-0.12, 4.442]	16.47	0.94

2.2.3 Comparing within exposure conditions: How quickly does exposure begin to affect participants' responses?

Next, we compared how exposure affects listeners' categorization responses within each condition relative to listeners' responses prior to any exposure. These changes are summarized for the slope and PSE in Figure 6C & D, respectively. This visualization makes apparent two aspects of participants' behavior that were not readily apparent in the statistical comparisons we have summarized so far. First, while the PSEs for the +40 and +10 conditions were shifted rightwards compared to the baseline condition, both the +10 and the baseline condition actually shift leftwards relative to their pre-exposure starting point in Test 1. This is confirmed by Bayesian

hypothesis tests summarized in Table ??.

2.2.4 Results summary

This study was set up with several objectives in mind. We aimed to replicate previous findings on distributional learning (Kleinschmidt & Jaeger, 2016) while introducing changes to the design to a) increase the ecological validity of results b) illuminate how soon distributional learning effects can be detected and c) allow investigation into the incremental process of belief updating as predicted by the IA framework. [POSSIBLE TO INCLUDE HERE IF THIS IS INTRODUCED AS A SECONDARY OBJECTIVE WHEN DESCRIBED IN THE METHODS: In setting the three exposure conditions we also noted a fourth possible investigation, that is, to test for the presence of “shrinkage” as first discussed in (Kleinschmidt, 2020; Kleinschmidt & Jaeger, 2016). In implementing the study this last objective could not be satisfactorily answered therefore we leave its elaboration to the discussion section.]

In consonance with previous studies we find that listeners changed their categorization behavior in the direction of the shift in the exposure talker’s VOT distributions. This provides new evidence that listeners do respond to talker statistics when the stimuli are more human-like and sampled from distributions that replicate the variability one would encounter in real life. In test block 1 participants in all groups converged on the same prior categorisation function but then their boundaries spread apart after the first exposure block. Regression analysis showed evidence in favour of the differences in boundary estimates between conditions in test blocks 2 to 4, and these differences were consistent with the direction of the distributional shift. The +10ms condition had a boundary to the right of the baseline condition and the +40ms group had a boundary right of the +10ms condition. This order of the boundary placements was maintained throughout all test blocks after the onset of exposure but their differences began to narrow from test block 5 suggesting a dissipation of distributional learning without further informative exposure.

A second finding from this study which remained opaque in previous work was that categorization differences between the groups emerged very early on after exposure. It took as few as 48 exposure trials for a clear difference to emerge between the groups. Although we do not yet

know if learning was already present prior to the 48 trials, that it does not take hundreds of exposures for listeners to exhibit changes in categorizations aligns with other speech adaptation studies employing different paradigms such as perceptual recalibration and L2 accent adaptation (Bradlow and Bent (2008); Clarke and Garrett (2004); (norris2006?)).

We found some evidence for incremental change in categorisation boundaries as listeners received more input of the talker’s cue distributions although this was not always clear from one block to another due to the uncertainty in boundary estimates. Looking at the PSE estimates at each block as a proportion of the ideal boundary implied by their respective distributions (labels Fig. 6), in the +40ms condition listeners increased the shift by roughly 10 percent in the third test block (after 96 exposure trials) from the second block but appeared to regress slightly in test block 4. In the +10ms condition boundaries did shift incrementally after each exposure block but the proportion of while in the baseline condition, listeners showed a slight regression in test block 3 before increasing their shift towards the implied boundary in test block 4. These mixed patterns between the conditions do not clearly tell us

In this experiment we also found that the bulk of the maximum boundary shift that each group would make by the end of all 144 exposures was achieved after the first 48 exposure trials. In the +40ms condition listeners achieved their maximum shift in test block 3

What is common to all three conditions is that none of the groups converged on the category boundary implied by the exposure distributions of their respective conditions.

To understand this pattern, it is helpful to relate our exposure conditions to the distribution of VOT in listeners’ prior experience. Figure 7 shows the mean and covariance of our exposure conditions relative to the distribution of VOT by talkers of L1-US English (based on Chodroff & Wilson, 2018). This comparison offers an explanation as to why the baseline condition (and to some extent the +10 condition) shift leftwards with increasing exposure, whereas the +40 condition shifts rightwards: relative to listeners’ prior experience our baseline condition actually presented lower-than-expected category means; of our three exposure conditions, only the +40 condition presented larger-than-expected category means. That is, once we take into account how our exposure conditions relate to listeners’ prior experience, both the direction of changes from Test 1 to 4 *within* each exposure condition, and the direction of




Figure 7. Placement of exposure stimuli relative to an estimate of typical phonetic distributions for 6914 word-initial /d/ and /t/ productions in L1-US English (based on 92 talkers in Chodroff & Wilson, 2018). The outermost contour of each category shows the 95% density quantile. Points show the category means of the exposure condition.

differences *between* exposure conditions receive an explanation.

Second, the reason for the slight decrease in the difference between the +10 and baseline conditions observed in Tables 1 and 2 (visible in Figure 6D as the decreasing difference between the green and red line) is *not* due to a reversal of the effects in the +10 condition. Rather, both conditions are changing in the same direction but the baseline condition stops changing after Test 2, which reduces the difference between the +10 and baseline conditions (see Table 1). The comparison across blocks thus suggests a rather uniform picture across all exposure conditions: participants' responses initially changed rapidly with exposure; with increasing exposure, these changes did not only slow down but seem to hit a hard constraint. Participants in the leftwards-shifted baseline condition did not exhibit any further changes in their categorization responses beyond Test 2. Similarly, participants in the rightwards-shifted +40 condition did not exhibit any further changes in their categorization responses beyond Test 3. Only participants in the leftward-shifted +10 condition still exhibit changes across blocks even from Test 3 to 4. But, perhaps tellingly, those participants also never reached the degree of shift that was evident in the baseline condition.

2.2.5 Constraints on cumulative changes

Finally, Figures 6C & D also compare participants' responses against those of an ideal observer that has fully learned the exposure distributions.

3 General discussion

- discuss consequences of findings for other accounts (decision-making; normalization)
- discuss fact that test stimuli deviate from exposure stimuli to different extent. on the one hand, it's just 1/4 of all trials. on the other hand, we do see relatively systematic changes in slopes each time we test. so there is evidence that even these 12 trials can affect categorisation slopes (though it is worth keeping in mind that this is a comparison across different sets of stimuli). could this explain shrinkage? unlikely since it wasn't the case in kleinschmidt and jaeger. could it explain the constraint on adaptation? that's less clear. we can, however, compare the relative mean of exposure and test.
- could some form of moving window with historical decay explain the findings? On the one hand if the moving window is very small, that would not explain why we do see some *cumulative* changes across blocks (window must be at least $48 + 12 = 60$ trials). on the other hand, the qualitative changes in the PSEs and slopes suggest that 12 trials can be enough to change some aspects of the categorisation function. it's thus *possible* that something that weighs recent input much more strongly but also considers less recent input beyond 48 trials might explain the overall pattern.
- discuss potential that observed adaptation maximizes accuracy under the choice rule. use psychometric function fit during unlabeled exposure trials to calculate *accuracy* (not likelihood) on labeled trials under criterion and under proportional matching decision rules. compare against accuracy if ideal observers categorization functions are used instead.

4 References

- Allen, J. S., & Miller, J. L. (1999). Effects of syllable-initial voicing and speaking rate on the temporal characteristics of monosyllabic words. *The Journal of the Acoustical Society of America*, 106(4), 2031–2039.
- Apfelbaum, K. S., & McMurray, B. (2015). Relative cue encoding in the context of sophisticated models of categorization: Separating information from categorization. *Psychonomic Bulletin & Review*, 22, 916–943.
- Baayen, R. H., Davidson, D. J., & Bates, D. M. (2008). Mixed-effects modeling with crossed random effects for subjects and items. *Journal of Memory and Language*, 59(4), 390–412.
- Bejjanki, V. R., Beck, J. M., Lu, Z.-L., & Pouget, A. (2011). Perceptual learning as improved probabilistic inference in early sensory areas. *Nature Neuroscience*, 14(5), 642–648.
- Boersma, P., & Weenink, D. (2022). *Praat: Doing phonetics by computer. Version 6.2. 12*.
- Bradlow, A. R., & Bent, T. (2008). Perceptual adaptation to non-native speech. *Cognition*, 106(2), 707–729.
- Bürkner, P.-C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical Software*, 80(1), 1–28. <https://doi.org/10.18637/jss.v080.i01>
- Chodroff, E., & Wilson, C. (2017). Structure in talker-specific phonetic realization: Covariation of stop consonant VOT in american english. *Journal of Phonetics*, 61, 30–47.
- Chodroff, E., & Wilson, C. (2018). Predictability of stop consonant phonetics across talkers: Between-category and within-category dependencies among cues for place and voice. *Linguistics Vanguard*, 4(s2).
- Clarke, C. M., & Garrett, M. F. (2004). Rapid adaptation to foreign-accented english. *The Journal of the Acoustical Society of America*, 116(6), 3647–3658.
- Clayards, M., Tanenhaus, M. K., Aslin, R. N., & Jacobs, R. A. (2008). Perception of speech reflects optimal use of probabilistic speech cues. *Cognition*, 108(3), 804–809.
- Cummings, S. N., & Theodore, R. M. (2023). Hearing is believing: Lexically guided

perceptual learning is graded to reflect the quantity of evidence in speech input.

Cognition, 235, 105404.

Davis, M. H., & Sohoglu, E. (2020). Three functions of prediction error for bayesian inference in speech perception. *The Cognitive Neurosciences*, 177–189.

Dmitrieva, O., Llanos, F., Shultz, A. A., & Francis, A. L. (2015). Phonological status, not voice onset time, determines the acoustic realization of onset f0 as a secondary voicing cue in spanish and english. *Journal of Phonetics*, 49, 77–95.

Docherty, G. J. (2011). The timing of voicing in british english obstruents. In *The timing of voicing in british english obstruents*. De Gruyter Mouton.

Fenn, K. M., & Hambrick, D. Z. (2013). What drives sleep-dependent memory consolidation: Greater gain or less loss? *Psychonomic Bulletin & Review*, 20, 501–506.

Flege, J. E., & Brown Jr, W. S. (1982). The voicing contrast between english/p/and/b/as a function of stress and position-in-utterance. *Journal of Phonetics*, 10(4), 335–345.

Harmon, Z., Idemaru, K., & Kapatsinski, V. (2019). Learning mechanisms in cue reweighting. *Cognition*, 189, 76–88.

Hitczenko, K., & Feldman, N. H. (2016). Modeling adaptation to a novel accent. *Proceedings of the Annual Conference of the Cognitive Science Society*.

Idemaru, K., & Holt, L. L. (2011). Word recognition reflects dimension-based statistical learning. *Journal of Experimental Psychology: Human Perception and Performance*, 37(6), 1939.

Idemaru, K., & Holt, L. L. (2020). Generalization of dimension-based statistical learning. *Attention, Perception, & Psychophysics*, 82, 1744–1762.

Jaeger, T. F. (2008). Categorical data analysis: Away from ANOVAs (transformation or not) and towards logit mixed models. *Journal of Memory and Language*, 59(4), 434–446.

Johnson, K. (1997). Speech perception without speaker normalization. In K. Johnson & J. W. Mullennix (Eds.), *Talker variability in speech processing* (pp. 145–146). San Diego: Academic Press.

Klatt, D. H. (1975). Voice onset time, frication, and aspiration in word-initial consonant clusters. *Journal of Speech and Hearing Research*, 18(4), 686–706.

- Kleinschmidt, D. (2020). *What constrains distributional learning in adults?*
- Kleinschmidt, D., & Jaeger, T. F. (2011). A bayesian belief updating model of phonetic recalibration and selective adaptation. *Proceedings of the 2nd Workshop on Cognitive Modeling and Computational Linguistics*, 10–19.
- Kleinschmidt, D., & Jaeger, T. F. (2012). A continuum of phonetic adaptation: Evaluating an incremental belief-updating model of recalibration and selective adaptation. *Proceedings of the Annual Meeting of the Cognitive Science Society*, 34.
- Kleinschmidt, D., & Jaeger, T. F. (2015). Robust speech perception: Recognize the familiar, generalize to the similar, and adapt to the novel. *Psychological Review*, 122(2), 148.
- Kleinschmidt, D., & Jaeger, T. F. (2016). What do you expect from an unfamiliar talker? *CogSci*.
- Kleinschmidt, D., Raizada, R. D., & Jaeger, T. F. (2015). Supervised and unsupervised learning in phonetic adaptation. *CogSci*.
- Lisker, L., & Abramson, A. S. (1964). A cross-language study of voicing in initial stops: Acoustical measurements. *Word*, 20(3), 384–422.
- Lisker, L., & Abramson, A. S. (1967). Some effects of context on voice onset time in english stops. *Language and Speech*, 10(1), 1–28.
- Liu, L., & Jaeger, T. F. (2018). Inferring causes during speech perception. *Cognition*, 174, 55–70.
- Liu, L., & Jaeger, T. F. (2019). Talker-specific pronunciation or speech error? Discounting (or not) atypical pronunciations during speech perception. *Journal of Experimental Psychology: Human Perception and Performance*, 45(12), 1562.
- Maye, J., Werker, J. F., & Gerken, L. (2002). Infant sensitivity to distributional information can affect phonetic discrimination. *Cognition*, 82(3), B101–B111.
- McMurray, B., & Jongman, A. (2011). What information is necessary for speech categorization? Harnessing variability in the speech signal by integrating cues computed relative to expectations. *Psychological Review*, 118(2), 219.
- Munson, C. M. (2011). *Perceptual learning in speech reveals pathways of processing* ({PhD} dissertation). The University of Iowa.

- Nixon, J. S., Rij, J. van, Mok, P., Baayen, R. H., & Chen, Y. (2016). The temporal dynamics of perceptual uncertainty: Eye movement evidence from cantonese segment and tone perception. *Journal of Memory and Language*, 90, 103–125.
- Pajak, B., & Levy, R. (2012). Distributional learning of L2 phonological categories by listeners with different language backgrounds. *Proceedings of the 36th Boston University Conference on Language Development*, 2, 400–413. Cascadilla Press Somerville, MA.
- Prins, N. (2011). The psychometric function: Why we should not, and need not, estimate the lapse rate. *Journal of Vision*, 11(11), 1175–1175.
- R Core Team. (2022). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- RStudio Team. (2020). *RStudio: Integrated development environment for r*. Boston, MA: RStudio, PBC. Retrieved from <http://www.rstudio.com/>
- Schertz, J., & Clare, E. J. (2020). Phonetic cue weighting in perception and production. *Wiley Interdisciplinary Reviews: Cognitive Science*, 11(2), e1521.
- Schuster, S. (2020). *Praat: Doing phonetics by computer [computer program]*. Stanford, CA: Interactive Language Processing Lab Stanford. Retrieved from <https://docs.proliferate.alps.science/en/latest/contents.html>
- Smith, B. L. (1978). Effects of place of articulation and vowel environment on 'voiced' stop consonant production. *Glossa*, 12, 163–175.
- Sumner, M. (2011). The role of variation in the perception of accented speech. *Cognition*, 119(1), 131–136.
- Tamminen, J., Davis, M. H., Merkx, M., & Rastle, K. (2012). The role of memory consolidation in generalisation of new linguistic information. *Cognition*, 125(1), 107–112.
- Tan, M., Xie, X., & Jaeger, T. F. (2021). Using rational models to interpret the results of experiments on accent adaptation. *Frontiers in Psychology*, 4523.
- Theodore, R. M., & Monto, N. R. (2019). Distributional learning for speech reflects cumulative exposure to a talker's phonetic distributions. *Psychonomic Bulletin &*

680 *Review*, 26, 985–992.

681 Vroomen, J., Linden, S. van, De Gelder, B., & Bertelson, P. (2007). Visual recalibration
682 and selective adaptation in auditory–visual speech perception: Contrasting build-up
683 courses. *Neuropsychologia*, 45(3), 572–577.

684 Westbury, J. R. (1979). Aspects of the temporal control of voicing in consonant clusters in
685 english. *Texas Linguistic Forum Austin, Tex*, 1–304.

686 Wichmann, F. A., & Hill, N. J. (2001). The psychometric function: I. Fitting, sampling,
687 and goodness of fit. *Perception & Psychophysics*, 63(8), 1293–1313.

688 Winn, M. B. (2020). Manipulation of voice onset time in speech stimuli: A tutorial and
689 flexible praat script. *The Journal of the Acoustical Society of America*, 147(2),
690 852–866.

691 Woods, K. J., Siegel, M. H., Traer, J., & McDermott, J. H. (2017). Headphone screening
692 to facilitate web-based auditory experiments. *Attention, Perception, & Psychophysics*,
693 79, 2064–2072.

694 Xie, X., Earle, F. S., & Myers, E. B. (2018). Sleep facilitates generalisation of accent
695 adaptation to a new talker. *Language, Cognition and Neuroscience*, 33(2), 196–210.

696 Xie, X., Jaeger, T. F., & Kurumada, C. (2023). What we do (not) know about the
697 mechanisms underlying adaptive speech perception: A computational framework and
698 review. *Cortex*.

699 Xie, X., Liu, L., & Jaeger, T. F. (2021). Cross-talker generalization in the perception of
700 nonnative speech: A large-scale replication. *Journal of Experimental Psychology:*
701 *General*, 150(11), e22.

702 Xie, X., Weatherholtz, K., Bainton, L., Rowe, E., Burchill, Z., Liu, L., & Jaeger, T. F.
703 (2018). Rapid adaptation to foreign-accented speech and its transfer to an unfamiliar
704 talker. *The Journal of the Acoustical Society of America*, 143(4), 2013–2031.

705 Zue, V. W. (1976). *Acoustic characteristics of stop consonants: A controlled study*.

706 MASSACHUSETTS INST OF TECH LEXINGTON LINCOLN LAB.