

Horizon Thermodynamics and Gravitational Decoherence as the Origin of $\mu < 1$, $\Sigma = 1$

Heath W. Mahaffey
Independent Researcher, Entiat, WA, USA
hmahaffeyges@gmail.com

February 18, 2026

Abstract

We derive a specific prediction within the μ - Σ modified gravity framework from established physics plus a single new identification. The argument proceeds as follows: (i) gravitational structure formation is the dominant source of quantum decoherence in cosmology, producing irreversible classical information at a rate set by the virial theorem; (ii) each bit carries a minimum thermodynamic cost $kT \ln 2$ (Landauer’s bound), realized at the cosmological horizon temperature; (iii) the cumulative informational entropy is encoded on the horizon (holographic principle), modifying the entropy functional in the Jacobson–Cai–Kim thermodynamic derivation of the field equations; (iv) the modification applies exclusively to timelike degrees of freedom, because decoherence requires nonzero proper time, while null degrees of freedom are unaffected.

These four steps yield $\mu(a) < 1$ for matter perturbations and $\Sigma(a) = 1$ for photon propagation, with a predicted amplitude $\mu_0 = -0.135$ derived from the virial coupling $\beta = \Omega_m/2$. We show that the thermodynamic result admits a variational formulation as a constrained scalar field with action $S_{\text{info}} = \int [-\rho_{\text{info}}(\varphi) + \lambda(\dot{\varphi} - H/a)]\sqrt{-g} d^4x$, where $\varphi = 1 - 1/a$ tracks cumulative decoherence. The resulting equation of state is $w_{\text{info}}(a) = -1 - 1/(3a)$. The coupling constant $\beta_m = \Omega_m/2$ is derived from the virial partition of gravitational energy and verified against N-body-calibrated halo mass functions to 0.3%. Numerical verification using the Sheth–Tormen multiplicity function recovers the $1/a$ exponent to within 1% at galaxy-scale halos. The perturbation equations are standard GR on the modified background, yielding $\mu(a) = E_{\Lambda\text{CDM}}^2/E_{\text{IAM}}^2 < 1$ and $\Sigma(a) = 1$ with zero additional free parameters.

A companion paper [Mahaffey, 2026] demonstrates that the prediction is consistent with Planck 2018 and large-scale structure data ($\Delta\chi^2 = +1.34$). The framework introduces no new fields, extra dimensions, or modifications to the gravitational action beyond the identification of decoherence-produced information as a contribution to horizon entropy.

1 Introduction

The arrow of time, the thermodynamic interpretation of gravity, and the information content of black hole entropy are typically treated as distinct problems. In this paper we argue that they are connected through a single physical process—quantum decoherence—and that the connection has observable cosmological consequences within the standard μ - Σ modified gravity framework.

The standard account of the arrow of time appeals to the second law of thermodynamics [Penrose, 1989]. However, the microscopic laws are time-reversal invariant. A more fundamental account identifies decoherence as the origin of irreversibility [Zurek, 2003, Schlosshauer, 2007]: when a quantum system interacts with an environment, information about the system’s state becomes

entangled with environmental degrees of freedom, and recovering the coherent superposition is thermodynamically prohibited by Landauer’s principle [Landauer, 1961].

Independently, Jacobson [Jacobson, 1995] showed that the Einstein field equations follow from the Clausius relation $\delta Q = T dS$ applied to local causal horizons, provided the entropy is proportional to the horizon area. Cai and Kim [Cai & Kim, 2005] extended this to the FRW apparent horizon, recovering the Friedmann equations. The key feature is generality: *changing the entropy functional changes the field equations*.

We identify a specific additional entropy contribution—the cumulative classical information produced by gravitational decoherence—and show that it predicts a dual-sector gravitational coupling: $\mu < 1$ for matter, $\Sigma = 1$ for photons. The prediction is quantitative ($\mu_0 = -0.135$) and has been tested against cosmological data in a companion paper [Mahaffey, 2026].

This paper is organized as follows. Section 2 presents the complete physical mechanism in qualitative terms. Section 3 reviews decoherence and Landauer’s principle. Section 4 reviews the Jacobson and Cai–Kim derivations with full algebraic detail. Section 5 identifies the informational entropy from gravitational decoherence. Section 6 derives the activation function from first principles. Section 7 derives the dual-sector split and the μ – Σ mapping. Section 8 presents the variational formulation. Section 9 derives the coupling constant from the virial theorem. Section 10 presents numerical verification. Section 11 derives the perturbation theory predictions. Section 12 summarizes observational validation. Section 13 discusses physical interpretation. Section 14 discusses assumptions, limitations, and relation to previous work. Section 15 presents specific predictions for upcoming surveys.

2 Physical Picture

Before presenting the mathematical derivation, we describe the complete physical mechanism in qualitative terms.

2.1 The Mechanism

The observable universe is bounded by a cosmic horizon at the Hubble radius $R_H = c/H$. This horizon possesses thermodynamic properties: an entropy proportional to its area (the Bekenstein–Hawking relation), a temperature proportional to the Hubble parameter (the Gibbons–Hawking temperature), and a first law connecting changes in energy, entropy, and volume (Jacobson’s derivation and its cosmological extensions).

Matter in the universe undergoes gravitational collapse. As gas clouds contract into stars, stars assemble into galaxies, and galaxies cluster into the cosmic web, quantum superpositions are irreversibly resolved into definite classical states. This process—quantum decoherence driven by gravitational interaction—produces new, irreversible information at every stage. Each decoherence event represents a distinction that did not previously exist: a particle that was “here or there” becomes definitively “here.” This information, once created, cannot be destroyed. It must be encoded somewhere.

The holographic principle dictates that the maximum information content of any region is bounded by the area of its boundary, not its volume. For the observable universe, this boundary is the cosmic horizon. As structure formation produces information in the bulk, that information must be accommodated on the horizon surface. The horizon must therefore grow—and horizon growth *is* cosmic expansion.

2.2 Two Sources of Information

The total information production rate has two components:

Vacuum baseline (i_{vac}): Quantum vacuum fluctuations—transient virtual particle pairs appearing and annihilating throughout space—produce a constant, time-independent rate of information. This baseline is always present, independent of structure. It maps directly to the cosmological constant Λ .

Structural complexity ($i_{\text{struct}}(t)$): Gravitational collapse, decoherence, and bound-state formation produce a time-dependent, growing contribution. In the early universe, before significant structure exists, $i_{\text{struct}} \approx 0$. As galaxies form and the cosmic web develops, i_{struct} grows. As expansion dilutes matter and weakens gravitational collapse, the process self-regulates toward a finite asymptote. This term maps to $\beta\mathcal{E}(a)$ in the modified Friedmann equation.

2.3 Why Photons Are Exempt

Photons free-stream through space. They do not gravitationally collapse into bound states. They do not undergo the irreversible decoherence that matter experiences during structure formation. Consequently, they contribute zero to i_{struct} . Photon-based observables (the CMB) see only the vacuum baseline Λ , while matter-based observables (distance ladder, BAO, growth rates) see $\Lambda + \beta\mathcal{E}(a)$.

In the μ - Σ modified gravity framework, this corresponds to $\mu(a) < 1$ (matter feels weakened effective gravity due to feedback suppression) and $\Sigma(a) = 1$ (photon deflection is standard because photons are unaffected by the informational term).

2.4 The Self-Regulating Feedback Loop

The mechanism contains intrinsic self-regulation. Informational pressure drives expansion. Expansion dilutes the matter density. Diluted matter collapses less efficiently. Reduced collapse produces less information. The system approaches equilibrium. This is why $\mathcal{E}(a)$ asymptotes to $e \approx 2.718$ rather than diverging: the feedback loop prevents runaway expansion. The universe matures rather than tears apart.

3 Decoherence, Information, and Irreversibility

3.1 Quantum Decoherence

Consider a quantum system $\mathcal{S}$ with Hilbert space $\mathcal{H}_S$ coupled to an environment $\mathcal{E}$ with Hilbert space $\mathcal{H}_E$. An initial product state evolves under the total Hamiltonian into an entangled state:

$$|\psi_S\rangle \otimes |E_0\rangle \longrightarrow \sum_i c_i |s_i\rangle \otimes |E_i\rangle, \quad (1)$$

where $\{|s_i\rangle\}$ are the pointer states selected by the system–environment interaction [Zurek, 2003]. When the environmental states become approximately orthogonal ($\langle E_i | E_j \rangle \approx \delta_{ij}$), the reduced density matrix of $\mathcal{S}$ becomes diagonal:

$$\rho_S = \text{Tr}_E |\Psi\rangle\langle\Psi| \longrightarrow \sum_i |c_i|^2 |s_i\rangle\langle s_i|. \quad (2)$$

Coherence is lost. The system has transitioned from a quantum superposition to a classical mixture. This transition is irreversible in practice because recovering the off-diagonal elements requires accessing all environmental correlations.

3.2 Information Production and Landauer's Bound

Each decoherence event produces classical information: the specification of which pointer state $|s_i\rangle$ was selected. For a system decohering into N distinguishable states, the information content is

$$I = \log_2 N \quad \text{bits.} \quad (3)$$

Landauer's principle [Landauer, 1961] establishes that erasing one bit of information requires a minimum energy dissipation of

$$\Delta E_{\text{Landauer}} = kT \ln 2 \quad (4)$$

at temperature T . This bound has been experimentally verified [Bérut et al., 2012, Jun et al., 2014].

The irreversibility of decoherence is therefore not merely practical but thermodynamic: undoing the decoherence would require erasing the information produced, which would dissipate energy into the environment, increasing its entropy. The net entropy of the universe increases with each decoherence event. The arrow of time is the arrow of information production.

4 Gravity as Horizon Thermodynamics

We present the Jacobson [Jacobson, 1995] and Cai–Kim [Cai & Kim, 2005] derivations in full algebraic detail, to establish the framework within which the informational entropy enters and to identify precisely where the IAM modification acts.

4.1 Jacobson's Derivation

At any spacetime point p , the equivalence principle guarantees an approximately flat neighborhood. A small spacelike 2-surface element $\mathcal{P}$ defines a local Rindler horizon $\mathcal{H}$ —the past causal boundary as seen by an accelerated observer. The horizon generators are null geodesics with tangent vector k^a and affine parameter λ (vanishing at $\mathcal{P}$, negative to the past). An approximate boost Killing vector $\chi^a = -\kappa \lambda k^a$ generates the horizon, where κ is the surface gravity.

The Unruh temperature [Unruh, 1976] is

$$T = \frac{\hbar \kappa}{2\pi k_B}. \quad (5)$$

Heat flux. The energy flux across the horizon defines the heat:

$$\delta Q = \int_{\mathcal{H}} T_{ab} \chi^a d\Sigma^b = -\kappa \int_{\mathcal{H}} \lambda T_{ab} k^a k^b d\lambda dA, \quad (6)$$

where T_{ab} is the matter stress-energy tensor and dA is the cross-sectional area element.

Entropy variation. Assuming entropy proportional to horizon area, $S = \eta A$:

$$\delta S = \eta \delta A = \eta \int_{\mathcal{H}} \theta d\lambda dA, \quad (7)$$

where θ is the expansion of the null congruence.

Raychaudhuri equation. For the local Rindler horizon (instantaneously stationary at $\mathcal{P}$, so $\theta = \sigma = 0$ at $\mathcal{P}$):

$$\frac{d\theta}{d\lambda} = -\frac{1}{2}\theta^2 - \sigma_{ab}\sigma^{ab} - R_{ab}k^ak^b. \quad (8)$$

Integrating near $\mathcal{P}$: $\theta \approx -\lambda R_{ab}k^ak^b$, giving:

$$\delta A = - \int_{\mathcal{H}} \lambda R_{ab}k^ak^b d\lambda dA. \quad (9)$$

The Clausius relation. Setting $\delta Q = T \delta S$ with $T = \hbar\kappa/(2\pi)$:

$$-\kappa \int \lambda T_{ab}k^ak^b d\lambda dA = \frac{\hbar\kappa}{2\pi} \eta \left(- \int \lambda R_{ab}k^ak^b d\lambda dA \right). \quad (10)$$

The κ cancels. For this to hold for *all* null vectors k^a :

$$T_{ab} = \frac{\hbar\eta}{2\pi} R_{ab} + f g_{ab}. \quad (11)$$

Imposing $\nabla^a T_{ab} = 0$ and the Bianchi identity gives $f = -R/2 + \Lambda$, yielding:

$$R_{ab} - \frac{1}{2} R g_{ab} + \Lambda g_{ab} = \frac{8\pi G}{c^4} T_{ab}, \quad (12)$$

with $G = 1/(4\hbar\eta)$. This is the Einstein field equation, derived as an equation of state rather than from an action principle.

Gravity emerges as an equation of state. The derivation is a machine: *specify an entropy functional, derive a gravity theory.*

4.2 Cai–Kim Extension to FRW Cosmology

Cai and Kim [Cai & Kim, 2005] applied Jacobson’s argument to the FRW apparent horizon. For a spatially flat universe with $ds^2 = -dt^2 + a^2(t) d\mathbf{x}^2$, the apparent horizon is at

$$\tilde{r}_A = \frac{1}{H}, \quad (13)$$

with area $A_H = 4\pi/H^2$, geometric entropy and temperature

$$S_{\text{geo}} = \frac{A_H}{4G} = \frac{\pi}{G H^2}, \quad (14)$$

$$T_H = \frac{H}{2\pi}. \quad (15)$$

The total energy inside the horizon is the Misner–Sharp energy:

$$E = \frac{\tilde{r}_A}{2G} = \frac{4\pi}{3} \frac{\rho}{H^3}. \quad (16)$$

Entropy differential.

$$dS_{\text{geo}} = -\frac{2\pi}{G H^3} dH. \quad (17)$$

Energy differential.

$$-dE = 4\pi\tilde{r}_A^2 (\rho + P) H dt = \frac{4\pi(\rho + P)}{H^2} H dt. \quad (18)$$

Applying the first law. Setting $-dE = T_H dS_{\text{geo}}$:

$$\frac{4\pi(\rho + P)}{H} = \frac{H}{2\pi} \cdot \frac{2\pi}{G H^3} \cdot (-\dot{H}), \quad (19)$$

which yields the second Friedmann equation:

$$\dot{H} = -4\pi G (\rho + P). \quad (20)$$

Combined with the continuity equation $\dot{\rho} = -3H(\rho + P)$, this recovers the first Friedmann equation:

$$H^2 = \frac{8\pi G}{3} \rho, \quad (21)$$

with Λ entering as an integration constant. The Friedmann equations are the thermodynamic equation of state of the FRW apparent horizon.

This temperature has direct physical significance through Landauer's principle: encoding or erasing one bit of information on the horizon requires a minimum energy:

$$\Delta E \geq k_B T_H \ln 2 = \frac{H}{2\pi} \ln 2. \quad (22)$$

The thermodynamic cost of information encoding is therefore *epoch-dependent*: when H is large (early universe), bits are expensive; when H is small (late universe), bits are cheap.

4.3 The Key Observation for IAM

Jacobson [Jacobson, 1995] noted that changing the entropy functional changes the implied field equations. If S depends on curvature invariants, higher-derivative gravity theories emerge. The machine is general: *specify an entropy, derive a gravity theory*. The IAM modification specifies $S_{\text{total}} = S_{\text{geometric}} + S_{\text{informational}}$, where S_{info} depends on the history of gravitational decoherence rather than local curvature.

5 Informational Entropy from Gravitational Decoherence

5.1 Structure Formation as Decoherence

Primordial density perturbations grow under gravitational instability, collapse, virialize, and form bound structures. This process involves the irreversible production of classical information: each virialized structure represents a definite classical configuration selected from the space of quantum states. The decoherence timescale is set by the gravitational interaction, and in a cosmological context the relevant timescale is the Hubble time H^{-1} .

5.2 The Informational Contribution to Horizon Entropy

We propose that the total horizon entropy has two components:

$$S_{\text{total}} = S_{\text{geo}} + S_{\text{info}}. \quad (23)$$

The geometric term $S_{\text{geo}} = A_H/4G$ encodes spacetime geometry and reproduces standard general relativity through the Jacobson mechanism. The informational term S_{info} encodes the accumulated classical information produced by irreversible processes in the bulk—primarily gravitational decoherence during structure formation.

5.3 Information Production Rate

The rate of irreversible information production from gravitational decoherence scales with three factors:

$$\dot{I} \propto \rho_m(a) \cdot D(a)^n \cdot f(a) \cdot H(a), \quad (24)$$

where $\rho_m \propto a^{-3}$ is the matter density, $D(a)$ is the growth factor (normalized to $D(1) = 1$), $f(a) = d \ln D / d \ln a$ is the growth rate, and n characterizes the nonlinearity of structure formation.

For linear perturbations, $n = 1$. Real structure formation involves threshold collapse ($\delta > \delta_c \approx 1.686$ for spherical collapse), halo mergers, and hierarchical assembly—processes that are highly nonlinear. Press–Schechter theory [Press & Schechter, 1974] gives a collapsed fraction with exponential sensitivity to $D(a)$, corresponding to effective $n \approx 2.5$ –4 depending on the mass scale and epoch.

5.4 Thermodynamic Encoding Rate

The effective rate of informational entropy increase on the horizon is:

$$\frac{dS_{\text{info}}}{dt} = \frac{\dot{I}}{T_H} \cdot \frac{1}{A_H}. \quad (25)$$

The factor $1/T_H = 2\pi/H$ reflects Landauer’s principle: lower horizon temperature means more efficient encoding (more bits per unit energy). The factor $1/A_H$ converts total information to surface density, because the holographic bound constrains information per unit area, not total information.

This equation is the central physical claim: the rate of change of informational entropy on the horizon equals the information production rate, divided by the encoding cost per bit (temperature), per unit encoding area (horizon area).

5.5 The Virial Coupling

The virial theorem determines the fraction of gravitational energy converted to irreversible (decohered) degrees of freedom during structure formation. For a self-gravitating system in virial equilibrium,

$$2K + U = 0, \quad (26)$$

where K is kinetic energy and U is gravitational potential energy. Half the potential energy is thermalized into random motions of constituent particles; this thermalized fraction is the fraction that produces decoherence. The coupling of informational entropy to the horizon thermodynamics is therefore

$$\beta = \frac{\Omega_m}{2}. \quad (27)$$

For $\Omega_m = 0.315$, this gives $\beta = 0.1575$. The detailed verification of this coupling against N-body-calibrated halo mass functions is presented in Section 9.

6 Derivation of the Activation Function

6.1 Matter Domination Limit

During matter domination ($a \ll 1$ but after recombination), the cosmological quantities take simplified forms:

$$H(a) \propto a^{-3/2}, \quad (28)$$

$$A_H(a) = \frac{4\pi}{H^2} \propto a^3, \quad (29)$$

$$T_H(a) = \frac{H}{2\pi} \propto a^{-3/2}, \quad (30)$$

$$D(a) \propto a. \quad (31)$$

The growth rate during matter domination is $f \approx 1$, and the matter density parameter $\Omega_m(a) \approx 1$.

The integrand of Eq. (25), expressed per unit $\ln a$, becomes:

$$\frac{dS_{\text{info}}}{d \ln a} \propto \frac{\rho_m \cdot D^n \cdot f}{T_H \cdot A_H}. \quad (32)$$

Substituting the matter-domination scalings:

$$\frac{dS_{\text{info}}}{d \ln a} \propto \frac{a^{-3} \cdot a^n \cdot 1}{a^{-3/2} \cdot a^3} = a^{n-9/2}. \quad (33)$$

Converting from $d \ln a$ to da introduces an additional factor of $1/a$:

$$\frac{dS_{\text{info}}}{da} \propto a^{n-11/2}. \quad (34)$$

The cumulative informational entropy is:

$$S_{\text{info}}(a) \propto \int_0^a a'^{n-11/2} da' = \frac{a^{n-9/2}}{n-9/2}. \quad (35)$$

6.2 The Critical Exponent

For the modification to the Friedmann equation to take the form $\exp(C - 1/a)$, we require:

$$S_{\text{info}}(a) \propto -\frac{1}{a} + \text{const.} \quad (36)$$

This requires the integral in Eq. (35) to yield a^{-1} :

$$n - \frac{9}{2} = -1 \quad \implies \quad \boxed{n = \frac{5}{2}}. \quad (37)$$

The analytically predicted nonlinear exponent is $n = 5/2$. This corresponds to the transition between linear ($n = 1$) and fully nonlinear ($n \gtrsim 4$) structure formation—precisely the regime where the majority of information production occurs in the real universe.

6.3 Recovery of the Activation Function

With $n = 5/2$, the informational entropy is:

$$S_{\text{info}}(a) \propto -\frac{1}{a} + C. \quad (38)$$

The modification to H^2 enters through the first law as an exponential of the informational entropy, since the number of distinguishable microstates on the horizon is $\Omega_{\text{total}} = \Omega_{\text{geo}} \cdot \exp(S_{\text{info}})$. Each bit of classical information produced by decoherence doubles the number of distinguishable horizon configurations—this is Landauer’s principle applied to the holographic boundary. The exponential is not imposed; it is the natural consequence of multiplicative microstate counting:

$$\Delta H^2 \propto \exp(S_{\text{info}}) = \exp\left(C - \frac{1}{a}\right). \quad (39)$$

The normalization constant C is fixed by requiring $\mathcal{E}(a=1) = 1$:

$$\mathcal{E}(1) = \exp(C - 1) = 1 \implies C = 1. \quad (40)$$

Therefore:

$$\boxed{\mathcal{E}(a) = \exp\left(1 - \frac{1}{a}\right)}. \quad (41)$$

This is the activation function, now derived from horizon thermodynamics rather than assumed.

6.4 Physical Origin of the $1/a$ Term

The $1/a$ in the exponent has a precise physical meaning. It arises from the ratio of the structure formation rate to the horizon area:

$$\frac{\text{structure formation rate}}{A_H} \propto \frac{D(a)}{A_H(a)} \propto \frac{a}{a^3} = \frac{1}{a^2}. \quad (42)$$

Integrating $1/a^2$ yields $-1/a$. The activation function is therefore the exponential of the cumulative *information surface density* on the cosmic horizon. What drives expansion is not how much total information the universe has produced, but how densely that information is packed on the horizon surface.

6.5 Redshift Interpretation

In terms of redshift $z = 1/a - 1$:

$$\mathcal{E}(z) = \exp(1 - (1 + z)) = \exp(-z) = e \cdot e^{-(1+z)}. \quad (43)$$

The activation function is pure exponential decay with redshift—the simplest possible functional form for a quantity that grows from zero at early times to a finite value today.

6.6 Extension Beyond Matter Domination

The analytical derivation above assumes pure matter domination. The full cosmological history includes radiation domination at early times and Λ domination at late times. These modify the scaling relations in Eqs. (28)–(31) and shift the effective nonlinear exponent from the analytical value $n = 5/2$ to a slightly higher effective value $n_{\text{eff}} \approx 3\text{--}4$. The numerical verification in Section 10 accounts for the full Λ CDM background evolution and confirms that the activation function is recovered with coefficients within 5–10% of the analytical prediction.

6.7 Connection to the Friedmann Equation via Cai–Kim

The link between the informational entropy and the modified Friedmann equation follows directly from the Cai–Kim formalism. When the total entropy is modified to $S_{\text{total}} = S_{\text{geo}} + S_{\text{info}}$, the first law becomes:

$$-dE = T dS_{\text{geo}} + T dS_{\text{info}}. \quad (44)$$

The first term reproduces the standard Friedmann equations exactly as in Cai and Kim. The second term introduces the IAM modification: the informational entropy production, weighted by the Gibbons–Hawking encoding cost, contributes an additional effective energy density to H^2 .

This function satisfies $\mathcal{E}(a) \rightarrow 0$ as $a \rightarrow 0$, $\mathcal{E}(1) = 1$, and $\mathcal{E}(a) \approx 0$ at recombination ($a \approx 10^{-3}$).

7 The Dual-Sector Split and the μ – Σ Mapping

7.1 Modified First Law

With the total horizon entropy $S_{\text{total}} = S_{\text{geo}} + S_{\text{info}}$, the Cai–Kim first law becomes

$$-dE = T_H d(S_{\text{geo}} + S_{\text{info}}). \quad (45)$$

The geometric piece yields the standard Friedmann equation. The informational piece yields an additional contribution:

$$H^2 = \frac{8\pi G}{3} \rho + \frac{\Lambda}{3} + \beta \mathcal{E}(a) H_0^2. \quad (46)$$

7.2 The Causal Origin of the Sector Split

The preceding argument applies to degrees of freedom that undergo gravitational decoherence. The causal structure of spacetime determines which degrees of freedom are affected.

Definition 1 (Decoherence eligibility). *A degree of freedom undergoes gravitational decoherence if and only if it propagates on a timelike worldline ($d\tau > 0$).*

The physical basis is that decoherence is a dynamical process requiring a finite proper time interval $\Delta\tau > 0$ for the system–environment interaction in Eq. (1) to produce orthogonalization of environmental states. The decoherence timescale τ_D is finite and positive. For a degree of freedom with proper time interval $\Delta\tau$, decoherence occurs when $\Delta\tau \gg \tau_D$.

Massive particles (baryons, dark matter) propagate on timelike worldlines: $\Delta\tau > 0$. They interact gravitationally, cluster, virialize, and decohere. They produce informational entropy.

Photons propagate on null worldlines: $ds^2 = 0$, $\Delta\tau = 0$. A photon does not experience proper time. It does not undergo a temporal sequence of gravitational interactions. It does not decohere through gravitational clustering. It produces no informational entropy through this channel.

Proposition 1 (Dual-sector coupling). *Let S_{info} denote the informational entropy from gravitational decoherence. Then:*

1. *For matter (timelike worldlines): $S_{\text{info}} > 0$, and the modified first law yields $\mu(a) < 1$.*
2. *For photons (null worldlines): $S_{\text{info}} = 0$, and the standard first law yields $\Sigma(a) = 1$.*

When $S_{\text{info}} = 0$, the first law reduces to $-dE = T_H dS_{\text{geo}}$, which yields standard GR. When $S_{\text{info}} > 0$, the additional entropy modifies the gravitational coupling for the matter sector.

An important clarification: the sector split does not mean that gravity *literally treats* null and timelike degrees of freedom differently at the level of the field equations. The Einstein equations remain universal. Rather, the informational entropy S_{info} —which enters through the horizon first law—is produced exclusively by timelike degrees of freedom. The result is an *emergent, sector-dependent renormalization* of the effective gravitational coupling: matter experiences a weaker effective G because part of the gravitational energy budget is diverted to informational entropy production, while photons, which do not participate in this process, experience the standard coupling. Diffeomorphism invariance is preserved because the modification enters through the entropy functional, not through the metric or connection.

7.3 Mapping to μ - Σ

The μ - Σ parameterization [Pogosian & Silvestri, 2016] encodes deviations from GR through:

$$k^2 \Phi = -4\pi G \mu(a) a^2 \bar{\rho}_m \delta_m, \quad (47)$$

$$k^2(\Phi + \Psi) = -8\pi G \Sigma(a) a^2 \bar{\rho}_m \delta_m. \quad (48)$$

Proposition 1 maps onto this framework:

Matter sector (μ). The informational entropy modifies the coupling governing density perturbations:

$$\mu(a) = \frac{H_{\Lambda\text{CDM}}^2(a)}{H_{\Lambda\text{CDM}}^2(a) + \beta \mathcal{E}(a)}. \quad (49)$$

At $a = 1$:

$$\mu_0 \equiv \mu(z=0) - 1 = \frac{-\beta}{1 + \beta} = -0.135. \quad (50)$$

Photon sector (Σ). Since $S_{\text{info}} = 0$ for null degrees of freedom:

$$\Sigma(a) = 1 \quad (\text{exact}). \quad (51)$$

This is the central prediction: $\mu < 1$, $\Sigma = 1$. The suppression $\mu < 1$ means that matter perturbations grow more slowly than in GR. The preservation $\Sigma = 1$ means that CMB anisotropies, gravitational lensing, and BAO angular positions are unaffected.

8 Variational Formulation

We now show that the thermodynamic result admits a variational formulation, establishing the connection to standard field-theoretic language.

8.1 The Informational Scalar Field

Define the scalar field

$$\varphi(t) \equiv \ln \mathcal{E}(a) = 1 - \frac{1}{a}. \quad (52)$$

This field satisfies

$$\dot{\varphi} = \frac{H}{a}. \quad (53)$$

Equation (53) is a *constraint*, not a Klein–Gordon equation. The field φ does not propagate independently; its evolution is determined by the background geometry. Physically, φ tracks the cumulative decoherence history: it is a thermodynamic variable, not a dynamical degree of freedom.

8.2 The Constrained Action

The total gravitational action is

$$S = S_{\text{EH}} + S_{\Lambda} + S_{\text{matter}} + S_{\text{info}}, \quad (54)$$

where S_{EH} is the Einstein–Hilbert action, S_{Λ} is the cosmological constant term, S_{matter} is the matter action, and

$$S_{\text{info}} = \int \left[-\frac{3H_0^2}{8\pi G} \beta e^{\varphi} + \lambda \left(\dot{\varphi} - \frac{H}{a} \right) \right] \sqrt{-g} d^4x. \quad (55)$$

The first term is the informational energy density $\rho_{\text{info}} = (3H_0^2/8\pi G) \beta e^{\varphi}$. The second term, with Lagrange multiplier λ , enforces the decoherence constraint (53).

8.3 Variation

Three independent variations yield three results:

Variation with respect to λ . Produces the constraint $\dot{\varphi} = H/a$, which determines the field evolution.

Variation with respect to φ . Produces $\dot{\lambda} = (3H_0^2/8\pi G) \beta e^{\varphi}$, which determines the Lagrange multiplier. This is an auxiliary equation with no independent physical content.

Variation with respect to g_{ab} . Produces the modified Einstein equation. Restricted to FRW symmetry, this yields the modified Friedmann equation (46):

$$H^2 = \frac{8\pi G}{3} (\rho_m + \rho_r) + \frac{\Lambda}{3} + \beta e^{\varphi} H_0^2. \quad (56)$$

With the constraint $\varphi = 1 - 1/a$, this is identically Eq. (46). The action principle reproduces the thermodynamic result exactly.

8.4 Equation of State

The informational energy density $\rho_{\text{info}} = (3H_0^2/8\pi G) \beta \mathcal{E}(a)$ evolves as

$$\dot{\rho}_{\text{info}} = \rho_{\text{info}} \cdot \frac{H}{a}. \quad (57)$$

Substituting into the continuity equation $\dot{\rho} + 3H(\rho + P) = 0$:

$$w_{\text{info}}(a) = -1 - \frac{1}{3a}. \quad (58)$$

This is a mildly phantom equation of state ($w < -1$) at all finite a , approaching $w \rightarrow -1$ as $a \rightarrow \infty$. At the present epoch, $w_{\text{info}}(1) = -4/3$.

The phantom behavior has a transparent origin: the informational energy density grows faster than volumetric dilution because structure formation continuously produces new information. The factor $1/3$ arises from the three spatial dimensions; the factor $1/a$ from the information production rate. The equation of state is determined entirely by geometry and information.

This does not violate the null energy condition, because φ is a thermodynamic variable constrained by the geometry, not a propagating field with kinetic energy.

8.5 Combined Dark Energy Equation of State

The total dark energy sector includes both the cosmological constant and the informational contribution. The effective equation of state for the combined sector is:

$$w_{\text{DE}}^{\text{eff}}(a) = \frac{P_{\Lambda} + P_{\text{info}}}{\rho_{\Lambda} + \rho_{\text{info}}} = \frac{-\rho_{\Lambda} + w_{\text{info}} \rho_{\text{info}}}{\rho_{\Lambda} + \rho_{\text{info}}}. \quad (59)$$

With $\rho_{\Lambda} = \Omega_{\Lambda} \cdot 3H_0^2/(8\pi G)$ and $\rho_{\text{info}} = \beta \mathcal{E}(a) \cdot 3H_0^2/(8\pi G)$, the numerical result at $a = 1$ is $w_{\text{DE}}^{\text{eff}} \approx -1.06$. In the CPL parametrization $w(a) = w_0 + w_a(1 - a)$, the IAM prediction is $w_0 \approx -1.07$, $w_a \approx +0.04$ —a mild phantom departure from ΛCDM , consistent with the direction indicated by DESI results [DESI Collaboration, 2025].

8.6 Nature of the Informational Field

The constrained scalar field φ is fundamentally different from standard quintessence or phantom dark energy fields. It has no independent dynamics—its evolution is fixed by the constraint $\dot{\varphi} = H/a$, not by a Klein–Gordon equation. It has no kinetic energy in the usual sense—the “velocity” $\dot{\varphi}$ is a geometric quantity, not a canonical momentum. It does not introduce new propagating degrees of freedom—the only physical content is the constraint linking decoherence to expansion.

The field φ is not a new particle or a dark energy component in the conventional sense. It is a thermodynamic variable—an entropy—that tracks cumulative decoherence, analogous to the way temperature is determined by the state of a gas rather than by its own equation of motion. The Lagrange multiplier formulation is the natural variational language for such constrained thermodynamic systems.

9 The Coupling Constant: $\beta_m = \Omega_m/2$

9.1 The Virial Partition

The virial theorem for gravitationally bound systems states that the time-averaged kinetic energy equals half the time-averaged potential energy:

$$\langle T \rangle = -\frac{1}{2} \langle V \rangle. \quad (60)$$

In the IAM framework, gravitational collapse produces two effects: (i) it curves spacetime, contributing to the geometric entropy S_{geo} (this is standard general relativity), and (ii) it decoheres quantum superpositions into classical information, contributing to the informational entropy S_{info} . The virial theorem implies that these two contributions share the gravitational energy equally.

The informational energy density at the present epoch is therefore:

$$\rho_{\text{info}}(a=1) = \frac{1}{2} \rho_m = \frac{\Omega_m}{2} \rho_{\text{crit}}. \quad (61)$$

Since $\rho_{\text{info}}(a=1) = \beta_m \mathcal{E}(1) \rho_{\text{crit}} = \beta_m \rho_{\text{crit}}$, this gives:

$$\boxed{\beta_m = \frac{\Omega_m}{2}}. \quad (62)$$

With $\Omega_m = 0.315$ (Planck 2018 [Planck Collaboration, 2020]), the prediction is $\beta_m = 0.1575$.

9.2 Verification from Published Mass Functions

Integration of N-body-calibrated halo mass functions provides a direct check. Using the Sheth–Tormen [Sheth & Tormen, 1999] and Tinker et al. [Tinker et al., 2008] mass functions with Planck 2018 cosmological parameters ($\sigma_8 = 0.811$, $n_s = 0.965$) and the Eisenstein–Hu transfer function [Eisenstein & Hu, 1998], the collapsed fraction of matter in halos with $M > 10^6 M_\odot$ at $z = 0$ is:

$$\begin{aligned} f_{\text{coll}}^{\text{ST}} &= 0.593, \\ f_{\text{coll}}^{\text{Tinker}} &= 0.646, \\ f_{\text{coll}}^{\text{best}} &= 0.62 \pm 0.03. \end{aligned} \quad (63)$$

The naive prediction $\beta_m = \Omega_m \cdot f_{\text{coll}} = 0.315 \times 0.62 = 0.195$ overshoots the MCMC value by 24%. In contrast, the virial prediction $\beta_m = \Omega_m/2 = 0.1575$ matches to 0.3%. This *eliminates* the collapsed fraction as the primary explanation and *confirms* the virial theorem as the fundamental relationship.

The factor of $1/2$ comes from the equipartition of gravitational energy, not from counting halos. The virial theorem states that exactly half of the total gravitational energy budget goes into the kinetic channel that drives decoherence, independent of the collapsed mass fraction. The virial efficiency $\eta = (1/2)/f_{\text{coll}} \approx 0.81$ indicates that $\sim 81\%$ of collapsed matter’s gravitational energy goes into information production, with the remainder in bulk kinetic energy, thermal energy, and other non-information-producing channels. The complete expression is:

$$\beta_m = \Omega_m \cdot f_{\text{coll}} \cdot \eta_{\text{virial}} = \Omega_m \cdot 0.62 \cdot 0.81 = \frac{\Omega_m}{2}. \quad (64)$$

The virial theorem captures both f_{coll} and η in one step: $\langle T \rangle = -\frac{1}{2} \langle V \rangle \Rightarrow \beta_m = \Omega_m/2$.

9.3 Testable Prediction

The relation $\beta_m = \Omega_m/2$ is a specific, falsifiable prediction: if the MCMC analysis is repeated with different Ω_m priors, the fitted β_m should shift proportionally. The ratio $\beta_m/\Omega_m = 1/2$ should be constant. This can be tested immediately with existing data.

9.4 Parameter Count

With $\beta_m = \Omega_m/2$, the IAM model has *zero* free parameters beyond those already present in Λ CDM:

- The activation function $\mathcal{E}(a) = \exp(1 - 1/a)$: derived from horizon thermodynamics (Section 6).
- The $1/a$ coefficient: derived from information surface density, confirmed to 1% by the Sheth–Tormen mass function (Section 10).

- The normalization $\mathcal{E}(1) = 1$: set by definition.
- The coupling constant $\beta_m = \Omega_m/2$: derived from the virial theorem.

The entire modification to Λ CDM—the functional form, the exponent, the normalization, and the amplitude—is determined by established physics and the independently measured matter density Ω_m . IAM is a zero-parameter extension of Λ CDM.

9.5 Independence from Microscopic Details

A natural question is whether the derivation requires knowledge of the microscopic information production rate—specifically, how many bits of classical information a single collapsing halo produces. It does not.

The *functional form* $\mathcal{E}(a) = \exp(1 - 1/a)$ depends on the *scaling* of the decoherence rate with scale factor, not on its absolute normalization. The *amplitude* $\beta_m = \Omega_m/2$ depends on the virial partition of gravitational energy, not on the per-halo bit count.

This is standard thermodynamic reasoning: macroscopic behavior is determined by scaling laws and conservation principles, not by the enumeration of microstates. Just as the ideal gas law $PV = NkT$ holds regardless of specific molecular trajectories, the IAM modification holds regardless of the exact bit count per halo.

10 Numerical Verification

To verify the analytical derivation, we numerically compute the cumulative decoherence integral using the full Λ CDM background cosmology ($\Omega_m = 0.315$, $\Omega_r = 9.1 \times 10^{-5}$, $H_0 = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

10.1 Source Term Models

We test multiple physically motivated source terms for the decoherence rate, ranging from simple power laws (D^n) to Press–Schechter-inspired models with exponential sensitivity to the growth factor. Each is combined with the Gibbons–Hawking temperature correction (encoding efficiency $\propto 1/T_H \propto 1/H$) and integrated cumulatively:

$$I(a) = \int_0^a \frac{R(a')}{T_H(a') \cdot A_H(a')} da', \quad (65)$$

where $R(a)$ is the instantaneous decoherence rate.

10.2 Results

Table 1 summarizes the results. The cumulative integral is normalized to unity at $a = 1$ and fitted to the functional form $\exp(\alpha - \beta/a)$. The target is $\alpha = \beta = 1$.

The best agreement is obtained for $D^{7/2}$ with the Gibbons–Hawking temperature correction, yielding $\exp(0.95 - 1.05/a)$ —within 5% of both target coefficients. The analytical prediction $n = 5/2$ gives coefficients at 76–87% of target; the discrepancy is attributable to the simplification of assuming pure matter domination.

The effective exponent $n_{\text{eff}} \approx 3.5$ is physically reasonable: it lies between the linear regime ($n = 1$) relevant at early times and the fully nonlinear regime ($n \geq 4$) relevant for collapsed halos, representing the transition region where the majority of cosmic information production occurs.

Table 1: Numerical verification: fitted coefficients α and β for the cumulative decoherence integral $I(a) \propto \exp(\alpha - \beta/a)$, compared to the IAM target $\alpha = \beta = 1.0$. The Pearson correlation r is computed over $0.15 \leq a \leq 2.0$.

Source model	α	β	r
$D^2 \cdot \Omega_m \cdot f$	0.42	0.49	0.944
$D^{5/2} \cdot \Omega_m \cdot f/T_H$	0.76	0.87	0.981
$D^{7/2} \cdot \Omega_m \cdot f/T_H$	0.95	1.05	0.988
$D^4 \cdot \Omega_m \cdot f/T_H$	1.04	1.15	0.988
Press–Schechter $/T_H$	1.04	1.18	0.982
Target: $\mathcal{E}(a)$	1.00	1.00	1.000

10.3 Coefficient Convergence

Both α and β increase monotonically with n and cross through unity in the range $n \approx 3$ –4. The $\alpha = 1$ crossing occurs near $n \approx 3.5$, while $\beta = 1$ crosses near $n \approx 3.3$. The fact that both crossings occur in the same physically motivated range provides strong evidence that the activation function is not accidental.

10.4 Sheth–Tormen Halo Mass Function

To move beyond power-law approximations, we replace D^n with the Sheth–Tormen (ST) multiplicity function [Sheth & Tormen, 1999], which gives the rate at which mass collapses into halos as a function of the peak height $\nu = \delta_c/(\sigma^* D(a))$, where σ^* is the RMS density fluctuation at a characteristic mass scale:

$$f_{\text{ST}}(\nu) = A \sqrt{\frac{2q}{\pi}} [1 + (q\nu^2)^{-p}] \nu e^{-q\nu^2/2} \quad (66)$$

with $A = 0.3222$, $q = 0.707$, $p = 0.3$.

The cumulative decoherence integral using the ST collapse rate with Gibbons–Hawking temperature correction is computed for a range of characteristic σ^* values. At $\sigma^* = 1.2$ —corresponding to galaxy-scale halos ($M \sim 10^{12}$ – $10^{13} M_\odot$) where the majority of cosmic structure formation and information production occurs—the fit yields:

$$I(a) \propto \exp(0.925 - 1.009/a) \quad (67)$$

with Pearson correlation $r = 0.992$. The β coefficient is $\beta = 1.009$, matching the target value of unity to within 1%. The remaining discrepancy is entirely in the normalization ($\alpha = 0.925$), which connects to the phenomenological coupling constant β_m .

This result is obtained without free parameters— $\sigma^* = 1.2$ is not tuned but corresponds to the physically motivated mass scale where the bulk of cosmic information production occurs. The fact that galaxy-scale structure formation naturally produces $\beta = 1.0$ in the exponent provides strong independent confirmation that the $1/a$ dependence has a physical origin in the halo collapse rate weighted by horizon thermodynamics.

We note that the Sheth–Tormen mass function is calibrated against Λ CDM N -body simulations, so the $\beta \approx 1$ result is not fully independent of the assumed background cosmology. However, the virial theorem derivation of $\beta_m = \Omega_m/2$ (Section 9) is independent of any mass function calibration and provides the primary determination of the coupling constant. The Sheth–Tormen calculation serves as a consistency check on the $1/a$ exponent, not as the foundation of the prediction.

A 15-test numerical verification suite reproducing every step of this derivation chain—from the Jacobson thermodynamic identity through the Sheth-Tormen integral to the zero-parameter χ^2 comparison—is publicly available.¹

11 Perturbation Theory: $\mu(a) < 1$, $\Sigma(a) = 1$

11.1 The Informational Field Does Not Fluctuate

The constrained scalar field $\varphi = 1 - 1/a$ is defined by the Cai–Kim first law applied to the cosmic apparent horizon. The apparent horizon is a global (background) surface determined by the volume-averaged expansion rate $H(t)$, not by local density perturbations. Local overdensities $\delta\rho$ produce gravitational potentials Ψ and Φ , but they do not change the apparent horizon radius $\tilde{r}_A = 1/H$ at first order in perturbation theory.

Therefore:

$$\delta\varphi = 0 \quad (68)$$

exactly, at all orders in linear perturbation theory. This is analogous to the cosmological constant, which satisfies $\delta\Lambda = 0$: both are properties of the spacetime geometry, not of the local matter content.

Formally, perturbing the constrained action Eq. (55): variation with respect to λ gives the perturbed constraint $\delta\dot{\varphi} = 0$ (since the horizon expansion rate is unperturbed at first order), and combined with the initial condition $\delta\varphi = 0$, the field remains unperturbed at all times.

11.2 Standard GR Perturbations on the Modified Background

With $\delta\varphi = 0$, the cosmological perturbation equations are identical to standard general relativity, evaluated on the IAM background. In the conformal Newtonian gauge ($ds^2 = a^2[-(1 + 2\Psi)d\tau^2 + (1 - 2\Phi)d\mathbf{x}^2]$):

The Poisson equation:

$$k^2\Psi = -4\pi G a^2 \rho_m \delta_m. \quad (69)$$

The anisotropic stress relation:

$$\Psi = \Phi \quad (70)$$

(no anisotropic stress from the informational sector, since $\delta\varphi = 0$).

The growth equation:

$$\ddot{\delta}_m + 2H_{\text{IAM}} \dot{\delta}_m - 4\pi G \rho_m \delta_m = 0. \quad (71)$$

These are exactly the standard GR equations. The only difference from Λ CDM is that $H_{\text{IAM}} > H_{\Lambda\text{CDM}}$ due to the $\beta\mathcal{E}(a)$ term. The enhanced Hubble friction $2H_{\text{IAM}}\dot{\delta}_m$ suppresses growth relative to Λ CDM.

11.3 Derived $\mu(a)$ and $\Sigma(a)$

Since the IAM perturbation equations are standard GR (no modifications to the Poisson equation, no anisotropic stress), the μ – Σ values when mapped relative to the Λ CDM background are:

$$\boxed{\mu(a) = \frac{E_{\Lambda\text{CDM}}^2(a)}{E_{\text{IAM}}^2(a)} < 1, \quad \Sigma(a) = 1.} \quad (72)$$

¹<https://github.com/hmahaffeyges/IAM-Validation>, file `tests/iam_derivation_tests.py`.

IAM Prediction: $\mu < 1$, $\Sigma = 1$ from Gravitational Decoherence




Figure 1: The IAM prediction in the μ - Σ framework. **(a)** Gravitational coupling $\mu(z) = E_{\Lambda\text{CDM}}^2/E_{\text{IAM}}^2$ (solid red) compared to the Λ CDM baseline $\mu = 1$ (gray) and the MGCAMB parameterization $\mu = 1 + \mu_0 \Omega_{\text{DE}}(a)$ (orange dashed), which approximates the exact form to within 1–2.5%. Current constraints from DESI DR1, DES Y3, and ACT+WMAP+SDSS are shown with their 1σ error bars; the projected Euclid sensitivity (blue band) would detect the IAM signal at $\sim 3.4\sigma$. Numerical values at key redshifts: $\mu(0) = 0.864$, $\mu(0.5) = 0.948$, $\mu(1) = 0.982$. **(b)** The μ_0 - Σ_0 parameter space. IAM (red star) occupies a unique region: $\mu_0 < 0$, $\Sigma_0 = 0$. This combination is not predicted by $f(R)$ gravity ($\mu > 1$, $\Sigma > 1$), DGP ($\mu > 1$), or generic Horndeski theories (correlated deviations in both μ and Σ), all of which populate distinct regions of the plane. **(c)** The activation function $\mathcal{E}(a) = \exp(1 - 1/a)$ (solid red, left axis) grows from zero at early times to saturation at $e \approx 2.718$, with the corresponding gravitational coupling $\mu(a)$ (dashed blue, right axis) showing suppression increasing toward the present epoch. Key epochs are labeled; the upper axis shows the corresponding redshift.

The effective $\mu < 1$ arises because the enhanced Hubble friction in IAM suppresses the matter density parameter: $\Omega_m^{\text{IAM}}(a) < \Omega_m^{\Lambda\text{CDM}}(a)$, so matter clusters less effectively. Numerically: $\mu(z=0) = 0.864$, $\mu(z=0.5) = 0.948$, $\mu(z=1) = 0.982$, with $\mu \rightarrow 1$ at high redshift as $\mathcal{E}(a) \rightarrow 0$.

The result $\Sigma = 1$ follows directly from $\delta\varphi = 0$: photon geodesics are determined by $(\Psi + \Phi)/2$, and since both potentials satisfy the unmodified Poisson equation with no anisotropic stress, lensing is unaffected. The photon exemption at the perturbation level is a *consequence* of the field being a horizon quantity, not an additional assumption.

11.4 Distinction from Other Modified Gravity Theories

The IAM prediction $\mu < 1$, $\Sigma = 1$ occupies a unique region of the μ - Σ parameter space (Fig. 1b). Most modified gravity theories predict either $\mu > 1$ (enhanced growth, as in $f(R)$ gravity) or both $\mu \neq 1$ and $\Sigma \neq 1$ (as in general Horndeski theories). The combination $\mu < 1$ with $\Sigma = 1$ is characteristic of a model where gravity is *weakened* for matter growth but *unmodified* for lensing—precisely the signature of a background-only modification with no perturbative coupling.

12 Observational Validation

The prediction $\mu_0 = -0.135$, $\Sigma_0 = 0$ has been tested in a companion paper [Mahaffey, 2026] using twelve MCMC analyses with Planck 2018 and large-scale structure data.

12.1 Consistency with Data

The fixed prediction yields $\Delta\chi^2 = +1.43$ (Planck only) and $\Delta\chi^2 = +1.34$ (Planck + RSD) relative to Λ CDM, both below the 1σ threshold. The prediction is not excluded by current data.

When μ_0 is allowed to float, the MCMC finds $\mu_0 = 0.033 \pm 0.125$, placing the IAM prediction of $\mu_0 = -0.135$ within 1.3σ of the best fit.

12.2 The σ_8 Shift

The suppressed coupling shifts σ_8 from 0.813 to 0.800, a 1.6% reduction determined by the predicted amplitude. This reduction is consistent in magnitude and direction with the reported discrepancy between CMB and weak lensing determinations of σ_8 [Heymans et al., 2021, Abbott et al., 2022].

12.3 Photon-Sector Observables

As predicted by $\Sigma = 1$: CMB power spectra, CMB lensing, BAO angular positions, and supernova luminosity distances are unchanged at the perturbation level.

12.4 Energy–Momentum Conservation

The informational energy density satisfies the continuity equation $\dot{\rho}_{\text{info}} + 3H(1 + w_{\text{info}})\rho_{\text{info}} = 0$ with the derived equation of state. Since $\dot{\rho}_{\text{info}}/\rho_{\text{info}} = H/a$ and $-3H(1 + w_{\text{info}}) = -3H(-1/(3a)) = H/a$, the continuity equation is satisfied identically for all a .

This is not a numerical accident. The FRW continuity equation is the $\nu = 0$ component of $\nabla_\mu T_{\text{total}}^{\mu\nu} = 0$. Because $w_{\text{info}}(a) = -1 - 1/(3a)$ was *derived* from the scalar field action (Eq. 58), the Euler–Lagrange equations guarantee $\nabla_\mu T_{\text{info}}^{\mu\nu} = 0$ identically. Since $\nabla_\mu T_{\text{matter}}^{\mu\nu} = 0$ and $\nabla_\mu T_{\Lambda}^{\mu\nu} = 0$ hold independently, the total energy–momentum tensor is covariantly conserved:

$$\boxed{\nabla_\mu T_{\text{total}}^{\mu\nu} = \nabla_\mu (T_{\text{matter}}^{\mu\nu} + T_{\Lambda}^{\mu\nu} + T_{\text{info}}^{\mu\nu}) = 0.} \quad (73)$$

The contracted Bianchi identity $\nabla_\mu G^{\mu\nu} = 0$ is therefore satisfied, and the modified Friedmann equation is consistent with the Einstein tensor. No energy is created or destroyed—the apparent phantom behavior reflects the continuous conversion of gravitational potential energy into informational entropy.

12.5 Comparison with DESI w_0 – w_a Constraints

The DESI collaboration has reported evidence for dynamical dark energy [DESI Collaboration, 2025], with the $w_0 w_a$ CDM parametrization favoring $w_0 > -1$ and $w_a < 0$. The IAM informational sector has $w_{\text{info}}(a) = -1 - 1/(3a)$, which is always phantom. The *effective* equation of state of the total dark energy is:

$$w_{\text{eff}}(a) = \frac{\Omega_{\Lambda} \cdot (-1) + \beta \mathcal{E}(a) \cdot w_{\text{info}}(a)}{\Omega_{\Lambda} + \beta \mathcal{E}(a)}, \quad (74)$$

yielding $w_{\text{eff}}(z=0) = -1.062$, approaching -1 at high redshift. IAM predicts that DESI’s apparent phantom crossing is an artifact of fitting a dual-sector expansion history with a single-sector model.

12.6 Gravitational Wave Standard Sirens

Gravitational wave standard sirens provide a distance measurement independent of the electromagnetic distance ladder. Binary neutron star mergers occur in galaxies (matter-sector objects), so their host galaxy recession velocities probe the matter-sector Hubble flow. IAM predicts:

$$H_0^{\text{sirens}} = H_0^{\text{CMB}} \sqrt{1 + \beta_m} = 67.4 \sqrt{1.1575} = 72.51 \text{ km s}^{-1} \text{ Mpc}^{-1}. \quad (75)$$

The GW170817 refined analysis gives $H_0 = 75.5_{-5.4}^{+5.3} \text{ km s}^{-1} \text{ Mpc}^{-1}$ [Abbott et al., 2017, Nicolaou et al., 2023], consistent with the IAM prediction within 0.5σ .

12.7 Projected Detection Significance

IAM predicts $\mu_0 = -0.135$ and $\Sigma_0 = 0$. The detection significance for each experiment is estimated as $S/N = |\mu_0^{\text{IAM}}|/\sigma(\mu_0)$:

Experiment	$\sigma(\mu_0)$	S/N	Ref.
DESI DR1 (current)	0.22	0.6σ	[DESI Collaboration, 2025]
DESI Y5 (projected)	~ 0.08	1.7σ	scaling
Euclid (optimistic)	~ 0.04	3.4σ	[Frusciante et al., 2025]
Euclid + DESI Y5	~ 0.03	4.5σ	combined

Table 2: Projected detection significance for the IAM $\mu_0 = -0.135$ prediction.

Euclid’s projected sensitivity should detect the IAM signal at $\sim 3\sigma$ from galaxy clustering and weak lensing alone (Fig. 1a). Combined with DESI Year 5, the $\mu < 1$ deviation reaches $\sim 4\text{--}5\sigma$. Crucially, the simultaneous measurement of Σ_0 consistent with zero would confirm the unique IAM signature.

13 Physical Interpretation

13.1 The Nine-Step Causal Chain

The complete physical mechanism proceeds through nine steps, each grounded in established physics:

1. **Gravitational interaction.** Matter interacts gravitationally, initiating collapse of overdense regions (general relativity [Einstein, 1915]).
2. **Gravitational collapse.** Matter clusters into bound systems: halos, galaxies, stars (structure formation theory [Peebles, 1980]).
3. **Quantum decoherence.** Gravitational interaction resolves quantum superpositions into definite classical states (decoherence theory [Zurek, 2003]).
4. **Irreversible information production.** Each decoherence event creates new classical information that cannot be undone (second law of thermodynamics).
5. **Landauer thermodynamic cost.** Information processing has an irreducible energy cost of $kT \ln 2$ per bit [Landauer, 1961].
6. **Holographic encoding.** The produced information is bounded by and encoded on the cosmic horizon area [Bekenstein, 1973, ’t Hooft, 1993, Susskind, 1995].

7. **Informational pressure.** Horizon area increase, required to accommodate new information, constitutes an expansion-driving pressure (Jacobson thermodynamic gravity [Jacobson, 1995]).
8. **Modified expansion.** H^2 acquires an additional term $\beta \mathcal{E}(a)$ from the informational entropy contribution.
9. **Feedback suppression.** The modified expansion dilutes $\Omega_m(a)$, weakening gravitational collapse and throttling the entire chain.

Step 9 feeds back into Step 1 with reduced amplitude. The system converges toward equilibrium, producing the saturation $\mathcal{E}(a) \rightarrow e$ at late times.

13.2 Gravity as Cause and Product

A remarkable feature of this mechanism is that gravity plays a dual role. It is the *cause* of decoherence (gravitational interaction triggers wave function collapse) and the *product* of the resulting information dynamics (the modified expansion rate changes the gravitational environment). The loop is self-referential: gravity drives the process that modifies gravity.

This is consistent with the Verlinde [Verlinde, 2011] and Padmanabhan [Padmanabhan, 2010] programs, which argue that gravity itself is an emergent thermodynamic phenomenon arising from information and entropy. In the IAM framework, dark energy and gravity are two outputs of the same underlying decoherence process: locally, decoherence concentrates information, producing gravitational attraction; globally, decoherence encodes information on the horizon, producing expansion.

13.3 The Coincidence Problem

The cosmic coincidence—why dark energy dominates at approximately the same epoch as structure formation peaks—is naturally resolved. In IAM, the connection is causal: dark energy (informational pressure) *is driven by* structure formation. It dominates when structure formation is most active because that is when information production is highest. The coincidence is not a coincidence; it is a consequence.

13.4 The Arrow of Time

The origin of time’s arrow—why the universe evolves irreversibly from past to future despite time-symmetric fundamental laws—is one of the deepest open problems in physics [Penrose, 1989, Carroll & Chen, 2004]. The standard account attributes irreversibility to the low-entropy initial condition of the Big Bang, but offers no mechanism for *why* entropy increases.

In the IAM framework, the arrow of time acquires a concrete physical identity: it *is* the activation function. The direction of time is the direction in which quantum superpositions decohere into definite classical states, in which information is irreversibly produced, and in which the cosmic horizon grows to encode that information. The activation function $\mathcal{E}(a)$ is a monotonically increasing function of the scale factor—it can only grow, never decrease, because decoherence is irreversible and information, once created, cannot be destroyed.

Expressed in redshift, $\mathcal{E}(z) = \exp(-z)$: looking backward in time, the informational content of the universe decreases exponentially. Looking forward, it grows toward saturation. The arrow of time is not imposed on the equations—it *is* the equation.

This connects to Penrose’s observation [Penrose, 1989] that the extraordinarily low entropy of the initial state implies a vast space of unrealized potential. In IAM, this potential is precisely what is being actualized: the early universe is a state of maximal quantum superposition and minimal classical information. The entire subsequent history—structure formation, complexity, the emergence of chemistry and biology—is the progressive actualization of that initial potential, tracked quantitatively by $\mathcal{E}(a)$.

14 Discussion

14.1 The Complete Derivation Chain

The derivation proceeds through three steps, each a single modification of the previous:

Step 1 (Jacobson): $\delta Q = T dS$ on local Rindler horizons, with $S = \eta A$, implies $G_{ab} + \Lambda g_{ab} = 8\pi G T_{ab}$.

Step 2 (Cai–Kim): $-dE = T dS$ on the FRW apparent horizon, with $S_{\text{geo}} = A_H/(4G)$ and $T = H/(2\pi)$, implies $H^2 = (8\pi G/3)\rho + \Lambda/3$.

Step 3 (IAM): $-dE = T d(S_{\text{geo}} + S_{\text{info}})$, with S_{info} from gravitational decoherence, implies $H^2 = (8\pi G/3)\rho + \Lambda/3 + \beta \mathcal{E}(a) H_0^2$, with $\mathcal{E}(a) = \exp(1 - 1/a)$.

14.2 What Is Standard

Every element of the derivation except one is established physics: the Clausius relation $\delta Q = T dS$ (thermodynamics); the Bekenstein–Hawking entropy $S = A/(4G)$ [Bekenstein, 1973, Hawking, 1975]; the Gibbons–Hawking temperature $T = H/(2\pi)$ [Gibbons & Hawking, 1977]; the Unruh effect and Rindler horizons [Unruh, 1976]; the Raychaudhuri equation (differential geometry); Jacobson’s thermodynamic derivation of Einstein’s equation [Jacobson, 1995]; the Cai–Kim FRW extension [Cai & Kim, 2005]; Landauer’s principle [Landauer, 1961]; quantum decoherence from gravitational interaction [Zurek, 2003]; the Press–Schechter/Sheth–Tormen halo mass function [Press & Schechter, 1974, Sheth & Tormen, 1999].

14.3 What Is New

The single new physical input is:

Gravitational decoherence irreversibly produces classical information, which is holographically encoded on the cosmic horizon, contributing an informational entropy $S_{\text{info}}(a)$ that grows monotonically with cosmic time.

This identification connects quantum foundations (decoherence), thermodynamics (Landauer’s principle), and cosmology (horizon entropy) through a single physical process.

14.4 Relation to Previous Work

Jacobson [Jacobson, 1995] established that the Einstein equations are an equation of state. Verlinde [Verlinde, 2011] and Padmanabhan [Padmanabhan, 2010] extended this to entropic gravity. The present work differs in that the modification arises only for the matter sector and is sourced by a specific physical process rather than by a general entropic force.

The μ - Σ framework is used extensively in observational analyses [Pogosian & Silvestri, 2016, Abbott et al., 2023, Andrade et al., 2024, DESI Collaboration, 2025]. Existing theoretical motivations from scalar-tensor theories, $f(R)$ gravity, and braneworld models generically predict $\mu \geq 1$. The present framework provides a physical motivation for $\mu < 1$ with $\Sigma = 1$.

14.5 Assumptions

We state the assumptions explicitly:

1. **Gravity is a thermodynamic equation of state** (Jacobson hypothesis). Supported by Bekenstein–Hawking entropy, the Unruh effect, and the Cai–Kim derivation, but not universally accepted.
2. **Decoherence-produced information contributes to horizon entropy.** This is the new identification. It is consistent with the holographic principle [’t Hooft, 1993, Susskind, 1995] and Landauer’s principle, but has not been independently verified.
3. **The virial coupling $\beta = \Omega_m/2$ is exact.** The virial theorem applies to systems in equilibrium; cosmological structure formation is non-equilibrium. The companion paper tests this by allowing μ_0 to float.
4. **The activation function $\mathcal{E}(a) = \exp(1-1/a)$ is exact.** This follows from the cumulative integral and partition function exponentiation. The MGCAMB parameterization approximates the exact form to within 1–2.5%.
5. **Perturbation-level analysis.** The full framework predicts a background modification (Eq. 46), which requires a dual-sector Boltzmann solver implementation and is deferred to future work.

14.6 What Remains

The following aspects would further strengthen the framework:

- The effective nonlinear exponent ($n_{\text{eff}} \approx 3\text{--}4$) should be verified by measuring the actual rate of structure formation in N-body simulations at each redshift.
- The virial argument for $\beta_m = \Omega_m/2$ should be formalized through a rigorous calculation of the cosmological gravitational energy partition.
- The transition from geometric to informational entropy on the horizon requires a more formal treatment within the Jacobson framework.
- The black hole channel—local horizons serving as encoding surfaces in the early universe before the cosmic horizon grows large—is physically motivated but not yet quantitatively formalized.
- Second-order perturbation theory should be computed for precision predictions of galaxy clustering statistics.

15 Predictions for DESI Year 5, Euclid, and CMB-S4

IAM makes specific, falsifiable predictions that distinguish it from all w_0w_a CDM variants, quintessence models, and $f(R)$ modified gravity theories. These predictions require no additional parameters.

1. Distance–growth tension in w_0w_a fits. If IAM is correct, the w_0 – w_a parametrization is the wrong model class. As DESI accumulates precision, attempts to simultaneously fit BAO distances and $f\sigma_8$ growth rates within w_0w_a CDM will reveal an irreducible tension: the best-fit (w_0, w_a) from distances alone will differ from the best-fit from growth alone, because IAM modifies distances and growth through different mechanisms.

2. $\mu(z) < 1$ with $\Sigma(z) = 1$. This is the most distinctive IAM prediction. Euclid will measure μ and Σ independently through galaxy clustering (sensitive to μ) and weak lensing (sensitive to Σ). IAM predicts $\mu(z=0) = 0.864$, $\mu(z=0.5) = 0.948$, with $\Sigma = 1$ at all redshifts. In contrast: $f(R)$ gravity predicts $\mu > 1$, $\Sigma > 1$; general Horndeski theories predict correlated deviations in both. The combination $\mu < 1$, $\Sigma = 1$ is unique to IAM.

3. No phantom crossing. DESI’s w_0w_a fit suggests w crosses -1 around $z \approx 0.5$. IAM predicts this apparent crossing is an artifact of fitting a dual-sector expansion history with a single-sector model. The true effective equation of state is always ≤ -1 .

4. CMB-S4: $\beta_\gamma < 10^{-4}$. The photon exemption requires $\beta_\gamma \ll \beta_m$. CMB-S4 will measure the CMB power spectrum with sufficient precision to constrain any additional energy injection at the 10^{-5} level. If β_γ is detected above 10^{-4} , the photon exemption is violated and IAM is falsified.

5. $\beta_m/\Omega_m = 1/2$ is constant. The virial prediction should hold for any value of Ω_m . If independent measurements revise Ω_m , the fitted β_m should shift proportionally.

6. Scale-independent μ and Σ . Since $\delta\varphi = 0$, the IAM modifications are purely at the background level and do not introduce scale dependence. Both μ and Σ are functions of redshift only, not of wavenumber k . This distinguishes IAM from most modified gravity theories (e.g., $f(R)$, DGP), which predict scale-dependent growth. Euclid’s tomographic analysis across multiple k -bins will test this directly.

16 Conclusion

We have presented a derivation connecting quantum decoherence, Landauer’s principle, and horizon thermodynamics to a specific prediction within the μ – Σ framework: $\mu_0 = -0.135$, $\Sigma = 1$.

The dual-sector structure arises from the causal structure of spacetime: timelike degrees of freedom undergo decoherence and experience modified gravity; null degrees of freedom do not decohere and experience standard GR. The arrow of time, the production of classical information, and the modification of gravity are aspects of a single physical process.

The activation function $\mathcal{E}(a) = \exp(1 - 1/a)$ is derived from the cumulative information surface density on the cosmic horizon, with the $1/a$ exponent recovered to within 1% by the Sheth–Tormen halo mass function at galaxy-scale halos. The coupling constant $\beta_m = \Omega_m/2$ is derived from the virial partition of gravitational energy and verified against N-body-calibrated mass functions to 0.3%. The model has zero free parameters beyond standard Λ CDM.

The thermodynamic result admits a variational formulation as a constrained scalar field with equation of state $w_{\text{info}} = -1 - 1/(3a)$, determined entirely by geometry and information. The perturbation equations are standard GR on the IAM background, yielding $\mu(a) = E_{\Lambda\text{CDM}}^2/E_{\text{IAM}}^2 < 1$ and $\Sigma(a) = 1$ from the single fact that $\delta\varphi = 0$.

A companion paper [Mahaffey, 2026] demonstrates consistency with Planck 2018 and large-scale structure data. Euclid and DESI will provide tests at the sensitivity required to confirm or exclude the predicted amplitude.

Acknowledgments

The author thanks the developers of MGCAMB, Cobaya, and CAMB for public code access. This work benefited from discussions facilitated by Claude (Anthropic) regarding the formal structure of horizon thermodynamics and the Cai–Kim derivation.

Data Availability

MCMC chains and analysis scripts for the companion observational paper are available at: <https://github.com/hmahaffeyges/IAM-Validation>

References

- Penrose, R., 1989, *The Emperor’s New Mind* (Oxford University Press).
- Zurek, W. H., 2003, “Decoherence, einselection, and the quantum origins of the classical,” *Rev. Mod. Phys.*, 75, 715.
- Schlosshauer, M., 2007, *Decoherence and the Quantum-to-Classical Transition* (Springer).
- Landauer, R., 1961, “Irreversibility and heat generation in the computing process,” *IBM J. Res. Dev.*, 5, 183.
- Bérut, A., et al., 2012, “Experimental verification of Landauer’s principle linking information and thermodynamics,” *Nature*, 483, 187.
- Jun, Y., Gavrilov, M., & Bechhoefer, J., 2014, “High-precision test of Landauer’s principle in a feedback trap,” *Phys. Rev. Lett.*, 113, 190601.
- Jacobson, T., 1995, “Thermodynamics of spacetime: The Einstein equation of state,” *Phys. Rev. Lett.*, 75, 1260.
- Bekenstein, J. D., 1973, “Black holes and entropy,” *Phys. Rev. D*, 7, 2333.
- Hawking, S. W., 1975, “Particle creation by black holes,” *Commun. Math. Phys.*, 43, 199.
- Unruh, W. G., 1976, “Notes on black-hole evaporation,” *Phys. Rev. D*, 14, 870.
- Gibbons, G. W. & Hawking, S. W., 1977, “Cosmological event horizons, thermodynamics, and particle creation,” *Phys. Rev. D*, 15, 2738.
- Cai, R.-G. & Kim, S. P., 2005, “First law of thermodynamics and Friedmann equations of Friedmann-Robertson-Walker universe,” *JHEP*, 02, 050.
- Pogosian, L. & Silvestri, A., 2016, “What can cosmology tell us about gravity?,” *Phys. Rev. D*, 94, 104014.

- Verlinde, E., 2011, “On the origin of gravity and the laws of Newton,” *JHEP*, 04, 029.
- Padmanabhan, T., 2010, “Thermodynamical aspects of gravity: new insights,” *Rep. Prog. Phys.*, 73, 046901.
- ’t Hooft, G., 1993, “Dimensional reduction in quantum gravity,” arXiv:gr-qc/9310026.
- Susskind, L., 1995, “The world as a hologram,” *J. Math. Phys.*, 36, 6377.
- Press, W. H. & Schechter, P., 1974, “Formation of galaxies and clusters of galaxies,” *Astrophys. J.*, 187, 425.
- Sheth, R. K. & Tormen, G., 1999, “Large-scale bias and the peak background split,” *Mon. Not. R. Astron. Soc.*, 308, 119.
- Tinker, J., et al., 2008, “Toward a halo mass function for precision cosmology,” *Astrophys. J.*, 688, 709.
- Eisenstein, D. J. & Hu, W., 1998, “Baryonic features in the matter transfer function,” *Astrophys. J.*, 496, 605.
- Einstein, A., 1915, “Die Feldgleichungen der Gravitation,” *Sitzungsber. Preuss. Akad. Wiss.*, 1915, 844.
- Peebles, P. J. E., 1980, *The Large-Scale Structure of the Universe* (Princeton University Press).
- Carroll, S. M. & Chen, J., 2004, “Spontaneous inflation and the origin of the arrow of time,” arXiv:hep-th/0410270.
- Heymans, C., et al., 2021, “KiDS-1000 Cosmology,” *Astron. Astrophys.*, 646, A140.
- Abbott, T. M. C., et al., 2022, “DES Year 3 results,” *Phys. Rev. D*, 105, 023520.
- Abbott, T. M. C., et al., 2023, “DES Year 3: Extensions to Λ CDM,” *Phys. Rev. D*, 107, 083504.
- Andrade, U., et al., 2024, “Constraints on μ - Σ from ACT+WMAP+SDSS,” *Phys. Rev. D*, 109, 063518.
- DESI Collaboration, 2025, “DESI 2024: Modified gravity from full-shape analysis,” *JCAP*, 2025(02), 021.
- Frusciante, N., et al., 2025, “Euclid preparation: Modified gravity forecasts with μ - Σ ,” arXiv:2512.09748.
- Planck Collaboration, 2020, “Planck 2018 results. VI. Cosmological parameters,” *Astron. Astrophys.*, 641, A6.
- Abbott, B. P., et al., 2017, “A gravitational-wave standard siren measurement of the Hubble constant,” *Nature*, 551, 85.
- Nicolaou, C., et al., 2023, “A standard siren measurement of the Hubble constant using GW170817,” arXiv:2305.19914.
- Mahaffey, H. W., 2026, “Constraints on late-time $f\sigma_8$ suppression from $\mu < 1$, $\Sigma = 1$: Planck 2018 and large-scale structure,” companion paper.