

Horizon Thermodynamics and Gravitational Decoherence as the Origin of $\mu < 1$, $\Sigma = 1$

Heath W. Mahaffey
Independent Researcher, Entiat, WA, USA
hmahaffeyges@gmail.com

February 18, 2026

Abstract

We derive a specific prediction within the μ - Σ modified gravity framework from established physics plus a single new identification. The argument proceeds as follows: (i) gravitational structure formation is the dominant source of quantum decoherence in cosmology, producing irreversible classical information at a rate set by the virial theorem; (ii) each bit carries a minimum thermodynamic cost $kT \ln 2$ (Landauer’s bound), realized at the cosmological horizon temperature; (iii) the cumulative informational entropy is encoded on the horizon (holographic principle), modifying the entropy functional in the Jacobson–Cai–Kim thermodynamic derivation of the field equations; (iv) the modification applies exclusively to timelike degrees of freedom, because decoherence requires nonzero proper time, while null degrees of freedom are unaffected.

These four steps yield $\mu(a) < 1$ for matter perturbations and $\Sigma(a) = 1$ for photon propagation, with a predicted amplitude $\mu_0 = -0.135$ derived from the virial coupling $\beta = \Omega_m/2$. We show that the thermodynamic result admits a variational formulation as a constrained scalar field with action $S_{\text{info}} = \int [-\rho_{\text{info}}(\varphi) + \lambda(\dot{\varphi} - H/a)] \sqrt{-g} d^4x$, where $\varphi = 1 - 1/a$ tracks cumulative decoherence. The resulting equation of state is $w_{\text{info}}(a) = -1 - 1/(3a)$.

A companion paper [Mahaffey, 2026] demonstrates that the prediction is consistent with Planck 2018 and large-scale structure data ($\Delta\chi^2 = +1.34$). The framework introduces no new fields, extra dimensions, or modifications to the gravitational action beyond the identification of decoherence-produced information as a contribution to horizon entropy.

1 Introduction

The arrow of time, the thermodynamic interpretation of gravity, and the information content of black hole entropy are typically treated as distinct problems. In this paper we argue that they are connected through a single physical process—quantum decoherence—and that the connection has observable cosmological consequences within the standard μ - Σ modified gravity framework.

The standard account of the arrow of time appeals to the second law of thermodynamics [Penrose, 1989]. However, the microscopic laws are time-reversal invariant. A more fundamental account identifies decoherence as the origin of irreversibility [Zurek, 2003, Schlosshauer, 2007]: when a quantum system interacts with an environment, information about the system’s state becomes entangled with environmental degrees of freedom, and recovering the coherent superposition is thermodynamically prohibited by Landauer’s principle [Landauer, 1961].

Independently, Jacobson [Jacobson, 1995] showed that the Einstein field equations follow from the Clausius relation $\delta Q = T dS$ applied to local causal horizons, provided the entropy is proportional to the horizon area. Cai and Kim [Cai & Kim, 2005] extended this to the FRW apparent

horizon, recovering the Friedmann equations. The key feature is generality: *changing the entropy functional changes the field equations*.

We identify a specific additional entropy contribution—the cumulative classical information produced by gravitational decoherence—and show that it predicts a dual-sector gravitational coupling: $\mu < 1$ for matter, $\Sigma = 1$ for photons. The prediction is quantitative ($\mu_0 = -0.135$) and has been tested against cosmological data in a companion paper [Mahaffey, 2026].

This paper is organized as follows. Section 2 reviews decoherence and Landauer’s principle. Section 3 reviews the Jacobson and Cai–Kim derivations. Section 4 identifies the informational entropy from gravitational decoherence. Section 5 derives the dual-sector split and the μ – Σ mapping. Section 6 presents the variational formulation. Section 7 summarizes the observational consequences. Section 8 discusses assumptions, limitations, and relation to previous work.

2 Decoherence, Information, and Irreversibility

2.1 Quantum Decoherence

Consider a quantum system $\mathcal{S}$ coupled to an environment $\mathcal{E}$. An initial product state evolves under the total Hamiltonian into an entangled state:

$$|\psi_S\rangle \otimes |E_0\rangle \longrightarrow \sum_i c_i |s_i\rangle \otimes |E_i\rangle, \quad (1)$$

where $\{|s_i\rangle\}$ are the pointer states selected by the system–environment interaction [Zurek, 2003]. When the environmental states become approximately orthogonal ($\langle E_i | E_j \rangle \approx \delta_{ij}$), the reduced density matrix of $\mathcal{S}$ becomes diagonal:

$$\rho_S = \text{Tr}_E |\Psi\rangle\langle\Psi| \longrightarrow \sum_i |c_i|^2 |s_i\rangle\langle s_i|. \quad (2)$$

Coherence is lost. The system has transitioned from a quantum superposition to a classical mixture. This transition is irreversible in practice because recovering the off-diagonal elements requires accessing all environmental correlations.

2.2 Landauer’s Bound

Each decoherence event produces classical information: the specification of which pointer state was selected. Landauer’s principle [Landauer, 1961] establishes that erasing one bit of information requires a minimum energy dissipation of

$$\Delta E_{\text{Landauer}} = kT \ln 2 \quad (3)$$

at temperature T . This bound has been experimentally verified [Bérut et al., 2012, Jun et al., 2014].

The irreversibility of decoherence is therefore thermodynamic: undoing it requires erasing the produced information, which dissipates energy and increases the entropy of the environment. The arrow of time is the arrow of information production.

3 Gravity as Horizon Thermodynamics

3.1 Jacobson's Derivation

At any spacetime point, the equivalence principle guarantees a locally flat neighborhood. Consider a local Rindler horizon $\mathcal{H}$ generated by an approximate boost Killing vector χ^a . The Unruh temperature [Unruh, 1976] is

$$T = \frac{\hbar a_\perp}{2\pi c k_B}, \quad (4)$$

where $a_\perp$ is the acceleration perpendicular to $\mathcal{H}$. The heat flux through $\mathcal{H}$ is

$$\delta Q = \int_{\mathcal{H}} T_{ab} \chi^a d\Sigma^b, \quad (5)$$

and the entropy is proportional to the horizon area:

$$S = \frac{A}{4 \ell_P^2}. \quad (6)$$

The area change is related to the expansion of the null generators via the Raychaudhuri equation:

$$\frac{d\theta}{d\lambda} = -\frac{1}{2}\theta^2 - \sigma_{ab}\sigma^{ab} - R_{ab}k^a k^b. \quad (7)$$

Imposing $\delta Q = T dS$ for all local causal horizons, Jacobson [Jacobson, 1995] obtained the Einstein field equation:

$$R_{ab} - \frac{1}{2} R g_{ab} + \Lambda g_{ab} = \frac{8\pi G}{c^4} T_{ab}. \quad (8)$$

Gravity emerges as an equation of state. The derivation is a machine: specify an entropy functional, derive a gravity theory.

3.2 Cai–Kim Extension to FRW Cosmology

Cai and Kim [Cai & Kim, 2005] applied Jacobson's argument to the FRW apparent horizon. For a spatially flat universe with $ds^2 = -dt^2 + a^2(t) d\mathbf{x}^2$, the apparent horizon is at

$$\tilde{r}_A = \frac{1}{H}, \quad (9)$$

with geometric entropy and temperature

$$S_{\text{geo}} = \frac{\pi}{G H^2}, \quad (10)$$

$$T_H = \frac{H}{2\pi}. \quad (11)$$

Applying the first law $-dE = T_H dS_{\text{geo}}$ (where E is the Misner–Sharp energy), Cai and Kim recovered:

$$H^2 = \frac{8\pi G}{3} \rho, \quad (12)$$

$$\dot{H} = -4\pi G (\rho + P), \quad (13)$$

with Λ entering as an integration constant. The Friedmann equations are the thermodynamic equation of state of the FRW apparent horizon.

4 Informational Entropy from Gravitational Decoherence

4.1 Structure Formation as Decoherence

Primordial density perturbations grow under gravitational instability, collapse, virialize, and form bound structures. This process involves the irreversible production of classical information: each virialized structure represents a definite classical configuration selected from the space of quantum states. The decoherence timescale is set by the gravitational interaction, and in a cosmological context the relevant timescale is the Hubble time H^{-1} .

4.2 The Virial Coupling

The virial theorem determines the fraction of gravitational energy converted to irreversible (decohered) degrees of freedom during structure formation. For a self-gravitating system in virial equilibrium,

$$2K + U = 0, \quad (14)$$

where K is kinetic energy and U is gravitational potential energy. Half the potential energy is thermalized into random motions of constituent particles; this thermalized fraction is the fraction that produces decoherence. The coupling of informational entropy to the horizon thermodynamics is therefore

$$\beta = \frac{\Omega_m}{2}. \quad (15)$$

For $\Omega_m = 0.315$, this gives $\beta = 0.1575$.

4.3 The Activation Function

Each bit of classical information produced by decoherence is encoded on the cosmological horizon. Information is conserved on the horizon in the absence of a global erasure mechanism.

The information production rate per unit horizon area scales as H/a : H sets the causal timescale and $1/a$ accounts for dilution of the comoving density of decohering structures. The cumulative informational entropy is

$$\mathcal{S}_{\text{info}}(a) \propto 1 - \frac{1}{a}, \quad (16)$$

with the boundary condition $\mathcal{S}_{\text{info}}(a \rightarrow 0) = 0$ (no structures, no information).

The contribution to the thermodynamic potential is obtained by exponentiating, as appropriate for a partition function: each bit doubles the number of distinguishable microstates on the horizon (N bits $\rightarrow 2^N = e^{N \ln 2}$ microstates):

$$\mathcal{E}(a) = \exp\left(1 - \frac{1}{a}\right). \quad (17)$$

This function satisfies $\mathcal{E}(a) \rightarrow 0$ as $a \rightarrow 0$, $\mathcal{E}(1) = 1$, and $\mathcal{E}(a) \approx 0$ at recombination ($a \approx 10^{-3}$).

5 The Dual-Sector Split and the μ - Σ Mapping

5.1 Modified First Law

With the total horizon entropy $S_{\text{total}} = S_{\text{geo}} + S_{\text{info}}$, the Cai–Kim first law becomes

$$-dE = T_H d(S_{\text{geo}} + S_{\text{info}}). \quad (18)$$

The geometric piece yields the standard Friedmann equation. The informational piece yields an additional contribution:

$$H^2 = \frac{8\pi G}{3} \rho + \frac{\Lambda}{3} + \beta \mathcal{E}(a) H_0^2. \quad (19)$$

5.2 The Causal Origin of the Sector Split

The preceding argument applies to degrees of freedom that undergo gravitational decoherence. The causal structure of spacetime determines which degrees of freedom are affected.

Definition 1 (Decoherence eligibility). *A degree of freedom undergoes gravitational decoherence if and only if it propagates on a timelike worldline ($d\tau > 0$).*

The physical basis is that decoherence is a dynamical process requiring a finite proper time interval $\Delta\tau > 0$ for the system–environment interaction in Eq. (1) to produce orthogonalization of environmental states.

Massive particles (baryons, dark matter) propagate on timelike worldlines: $\Delta\tau > 0$. They interact gravitationally, cluster, virialize, and decohere. They produce informational entropy.

Photons propagate on null worldlines: $ds^2 = 0$, $\Delta\tau = 0$. A photon does not experience proper time. It does not undergo a temporal sequence of gravitational interactions. It does not decohere through gravitational clustering. It produces no informational entropy through this channel.

Proposition 1 (Dual-sector coupling). *Let S_{info} denote the informational entropy from gravitational decoherence. Then:*

1. *For matter (timelike worldlines): $S_{\text{info}} > 0$, and the modified first law yields $\mu(a) < 1$.*
2. *For photons (null worldlines): $S_{\text{info}} = 0$, and the standard first law yields $\Sigma(a) = 1$.*

When $S_{\text{info}} = 0$, the first law reduces to $-dE = T_H dS_{\text{geo}}$, which yields standard GR. When $S_{\text{info}} > 0$, the additional entropy modifies the gravitational coupling for the matter sector.

5.3 Mapping to μ – Σ

The μ – Σ parameterization [Pogosian & Silvestri, 2016] encodes deviations from GR through:

$$k^2 \Phi = -4\pi G \mu(a) a^2 \bar{\rho}_m \delta_m, \quad (20)$$

$$k^2(\Phi + \Psi) = -8\pi G \Sigma(a) a^2 \bar{\rho}_m \delta_m. \quad (21)$$

Proposition 1 maps onto this framework:

Matter sector (μ). The informational entropy modifies the coupling governing density perturbations:

$$\mu(a) = \frac{H_{\Lambda\text{CDM}}^2(a)}{H_{\Lambda\text{CDM}}^2(a) + \beta \mathcal{E}(a)}. \quad (22)$$

At $a = 1$:

$$\mu_0 \equiv \mu(z=0) - 1 = \frac{-\beta}{1 + \beta} = -0.135. \quad (23)$$

Photon sector (Σ). Since $S_{\text{info}} = 0$ for null degrees of freedom:

$$\Sigma(a) = 1 \quad (\text{exact}). \quad (24)$$

6 Variational Formulation

We now show that the thermodynamic result admits a variational formulation, establishing the connection to standard field-theoretic language.

6.1 The Informational Scalar Field

Define the scalar field

$$\varphi(t) \equiv \ln \mathcal{E}(a) = 1 - \frac{1}{a}. \quad (25)$$

This field satisfies

$$\dot{\varphi} = \frac{H}{a}. \quad (26)$$

Equation (26) is a *constraint*, not a Klein–Gordon equation. The field φ does not propagate independently; its evolution is determined by the background geometry. Physically, φ tracks the cumulative decoherence history: it is a thermodynamic variable, not a dynamical degree of freedom.

6.2 The Constrained Action

The total gravitational action is

$$S = S_{\text{EH}} + S_{\Lambda} + S_{\text{matter}} + S_{\text{info}}, \quad (27)$$

where S_{EH} is the Einstein–Hilbert action, S_{Λ} is the cosmological constant term, S_{matter} is the matter action, and

$$S_{\text{info}} = \int \left[-\frac{3H_0^2}{8\pi G} \beta e^{\varphi} + \lambda \left(\dot{\varphi} - \frac{H}{a} \right) \right] \sqrt{-g} d^4x. \quad (28)$$

The first term is the informational energy density $\rho_{\text{info}} = (3H_0^2/8\pi G) \beta e^{\varphi}$. The second term, with Lagrange multiplier λ , enforces the decoherence constraint (26).

6.3 Variation

Three independent variations yield three results:

Variation with respect to λ . Produces the constraint $\dot{\varphi} = H/a$, which determines the field evolution.

Variation with respect to φ . Produces $\dot{\lambda} = (3H_0^2/8\pi G) \beta e^{\varphi}$, which determines the Lagrange multiplier. This is an auxiliary equation with no independent physical content.

Variation with respect to g_{ab} . Produces the modified Einstein equation. Restricted to FRW symmetry, this yields the modified Friedmann equation (19):

$$H^2 = \frac{8\pi G}{3} (\rho_m + \rho_r) + \frac{\Lambda}{3} + \beta e^{\varphi} H_0^2. \quad (29)$$

With the constraint $\varphi = 1 - 1/a$, this is identically Eq. (19).

6.4 Equation of State

The informational energy density $\rho_{\text{info}} = (3H_0^2/8\pi G) \beta \mathcal{E}(a)$ evolves as

$$\dot{\rho}_{\text{info}} = \rho_{\text{info}} \cdot \frac{H}{a}. \quad (30)$$

Substituting into the continuity equation $\dot{\rho} + 3H(\rho + P) = 0$:

$$w_{\text{info}}(a) = -1 - \frac{1}{3a}. \quad (31)$$

This is a mildly phantom equation of state ($w < -1$) at all finite a , approaching $w \rightarrow -1$ as $a \rightarrow \infty$. At the present epoch, $w_{\text{info}}(1) = -4/3$.

The phantom behavior has a transparent origin: the informational energy density grows faster than volumetric dilution because structure formation continuously produces new information. The factor $1/3$ arises from the three spatial dimensions; the factor $1/a$ from the information production rate. The equation of state is determined entirely by geometry and information.

The field φ is not a new particle or a dark energy component in the conventional sense. It is a thermodynamic variable—an entropy—that tracks cumulative decoherence, analogous to the way temperature is determined by the state of a gas rather than by its own equation of motion. The Lagrange multiplier formulation is the natural variational language for such constrained thermodynamic systems.

7 Observational Consequences

The prediction $\mu_0 = -0.135$, $\Sigma_0 = 0$ has been tested in a companion paper [Mahaffey, 2026] using twelve MCMC analyses with Planck 2018 and large-scale structure data.

7.1 Consistency with Data

The fixed prediction yields $\Delta\chi^2 = +1.43$ (Planck only) and $\Delta\chi^2 = +1.34$ (Planck + RSD) relative to Λ CDM, both below the 1σ threshold. The prediction is not excluded by current data.

7.2 The σ_8 Shift

The suppressed coupling shifts σ_8 from 0.813 to 0.800, a 1.6% reduction determined by the predicted amplitude. This reduction is consistent in magnitude and direction with the reported discrepancy between CMB and weak lensing determinations of σ_8 .

7.3 Photon-Sector Observables

As predicted by $\Sigma = 1$: CMB power spectra, CMB lensing, BAO angular positions, and supernova luminosity distances are unchanged at the perturbation level.

7.4 Forecasts

Euclid and DESI are projected to achieve $\sigma(\mu_0) \sim 0.03$ – 0.05 , sufficient to test the predicted amplitude at $> 3\sigma$ significance [Frusciante et al., 2025].

8 Discussion

8.1 Summary of the Argument

The logical chain is:

1. Gravitational structure formation produces quantum decoherence.
2. Decoherence produces irreversible classical information.
3. Information has a minimum thermodynamic cost (Landauer).
4. Cumulative information is encoded on the cosmological horizon.
5. The encoding rate is set by the virial theorem: $\beta = \Omega_m/2$.
6. Modified horizon entropy $\rightarrow$ modified gravitational coupling for matter: $\mu < 1$.
7. Photons do not decohere ($\Delta\tau = 0$) $\rightarrow \Sigma = 1$.
8. The modification vanishes at early times: $\mathcal{E}(a \rightarrow 0) \rightarrow 0$.

Each step uses established physics except step 4, which is the sole new identification: decoherence-produced information contributes to horizon entropy.

8.2 Relation to Previous Work

Jacobson [Jacobson, 1995] established that the Einstein equations are an equation of state. Verlinde [Verlinde, 2011] and Padmanabhan [Padmanabhan, 2010] extended this to entropic gravity. The present work differs in that the modification arises only for the matter sector and is sourced by a specific physical process rather than by a general entropic force.

The μ - Σ framework is used extensively in observational analyses [Pogosian & Silvestri, 2016, Abbott et al., 2023, Andrade et al., 2024, DESI Collaboration, 2025]. Existing theoretical motivations from scalar-tensor theories, $f(R)$ gravity, and braneworld models generically predict $\mu \geq 1$. The present framework provides a physical motivation for $\mu < 1$ with $\Sigma = 1$.

8.3 Assumptions

We state the assumptions explicitly:

1. **Gravity is a thermodynamic equation of state** (Jacobson hypothesis). Supported by Bekenstein–Hawking entropy, the Unruh effect, and the Cai–Kim derivation, but not universally accepted.
2. **Decoherence-produced information contributes to horizon entropy.** This is the new identification. It is consistent with the holographic principle [’t Hooft, 1993, Susskind, 1995] and Landauer’s principle, but has not been independently verified.
3. **The virial coupling $\beta = \Omega_m/2$ is exact.** The virial theorem applies to systems in equilibrium; cosmological structure formation is non-equilibrium. The companion paper tests this by allowing μ_0 to float.

4. **The activation function $\mathcal{E}(a) = \exp(1-1/a)$ is exact.** This follows from the cumulative integral and partition function exponentiation. The MGCAMB parameterization approximates the exact form to within 1–2.5%.
5. **Perturbation-level analysis.** The full framework predicts a background modification (Eq. 19), which requires a dual-sector Boltzmann solver implementation and is deferred to future work.

9 Conclusion

We have presented a derivation connecting quantum decoherence, Landauer’s principle, and horizon thermodynamics to a specific prediction within the μ – Σ framework: $\mu_0 = -0.135$, $\Sigma = 1$.

The dual-sector structure arises from the causal structure of spacetime: timelike degrees of freedom undergo decoherence and experience modified gravity; null degrees of freedom do not decohere and experience standard GR. The arrow of time, the production of classical information, and the modification of gravity are aspects of a single physical process.

The thermodynamic result admits a variational formulation as a constrained scalar field with equation of state $w_{\text{info}} = -1 - 1/(3a)$, determined entirely by geometry and information.

A companion paper [Mahaffey, 2026] demonstrates consistency with Planck 2018 and large-scale structure data. Euclid and DESI will provide tests at the sensitivity required to confirm or exclude the predicted amplitude.

Acknowledgments

The author thanks the developers of MGCAMB, Cobaya, and CAMB for public code access. This work benefited from discussions facilitated by Claude (Anthropic) regarding the formal structure of horizon thermodynamics and the Cai–Kim derivation.

Data Availability

MCMC chains and analysis scripts for the companion observational paper are available at: <https://github.com/hmahaffeyges/IAM-Validation>

References

- Penrose, R., 1989, *The Emperor’s New Mind* (Oxford University Press).
- Zurek, W. H., 2003, “Decoherence, einselection, and the quantum origins of the classical,” *Rev. Mod. Phys.*, 75, 715.
- Schlosshauer, M., 2007, *Decoherence and the Quantum-to-Classical Transition* (Springer).
- Landauer, R., 1961, “Irreversibility and heat generation in the computing process,” *IBM J. Res. Dev.*, 5, 183.
- Bérut, A., et al., 2012, “Experimental verification of Landauer’s principle,” *Nature*, 483, 187.
- Jun, Y., Gavrilov, M., & Bechhoefer, J., 2014, “High-precision test of Landauer’s principle,” *Phys. Rev. Lett.*, 113, 190601.

- Jacobson, T., 1995, “Thermodynamics of spacetime: The Einstein equation of state,” *Phys. Rev. Lett.*, 75, 1260.
- Bekenstein, J. D., 1973, “Black holes and entropy,” *Phys. Rev. D*, 7, 2333.
- Hawking, S. W., 1975, “Particle creation by black holes,” *Commun. Math. Phys.*, 43, 199.
- Unruh, W. G., 1976, “Notes on black-hole evaporation,” *Phys. Rev. D*, 14, 870.
- Cai, R.-G. & Kim, S. P., 2005, “First law of thermodynamics and Friedmann equations of Friedmann-Robertson-Walker universe,” *JHEP*, 02, 050.
- Pogosian, L. & Silvestri, A., 2016, “What can cosmology tell us about gravity?,” *Phys. Rev. D*, 94, 104014.
- Verlinde, E., 2011, “On the origin of gravity and the laws of Newton,” *JHEP*, 04, 029.
- Padmanabhan, T., 2010, “Thermodynamical aspects of gravity: new insights,” *Rep. Prog. Phys.*, 73, 046901.
- ’t Hooft, G., 1993, “Dimensional reduction in quantum gravity,” arXiv:gr-qc/9310026.
- Susskind, L., 1995, “The world as a hologram,” *J. Math. Phys.*, 36, 6377.
- Heymans, C., et al., 2021, “KiDS-1000 Cosmology,” *Astron. Astrophys.*, 646, A140.
- Abbott, T. M. C., et al., 2022, “DES Year 3 results,” *Phys. Rev. D*, 105, 023520.
- Abbott, T. M. C., et al., 2023, “DES Year 3: Extensions to Λ CDM,” *Phys. Rev. D*, 107, 083504.
- Andrade, U., et al., 2024, “Constraints on μ - Σ from ACT+WMAP+SDSS,” *Phys. Rev. D*, 109, 063518.
- DESI Collaboration, 2025, “DESI 2024: Modified gravity from full-shape analysis,” *JCAP*, 2025(02), 021.
- Frusciante, N., et al., 2025, “Euclid preparation: Modified gravity forecasts with μ - Σ ,” arXiv:2512.09748.
- Mahaffey, H. W., 2026, “Constraints on late-time $f\sigma_8$ suppression from $\mu < 1$, $\Sigma = 1$: Planck 2018 and large-scale structure,” companion paper.