

# Corolario sobre la regularidad de la convolución

**Corolario 1.** Sea  $f \in \mathcal{L}^1(\mathbb{R}^n)$  y  $g \in \mathcal{C}^k(\mathbb{R}^n)$  tal que  $\forall \alpha \in (\mathbb{N} \cup \{0\})^n$  con  $|\alpha| \leq k$  se tiene que  $\|D_x^\alpha g\|_\infty < \infty$ . Entonces,

$$f * g \in \mathcal{C}^k(\mathbb{R}^n) \quad \wedge \quad \forall \alpha \in (\mathbb{N} \cup \{0\})^n : |\alpha| \leq k : D_x^\alpha (f * g) = f * (D_x^\alpha g).$$

**Demostración:** Basta con probar el caso  $\alpha = (1, 0, \dots, 0)$  y el resto se tiene por inducción.

Sea  $x \in \mathbb{R}^n$  y definimos  $F(x_1) = (g * f)(x) = \int_{\mathbb{R}^n} g(x - y) f(y) dy$  donde  $(x_2, \dots, x_n)$  son fijos. Queremos demostrar que

$$F'(x_1) = \frac{\partial}{\partial x_1} \int_{\mathbb{R}^n} g(x - y) f(y) dy = \int_{\mathbb{R}^n} \frac{\partial g}{\partial x_1}(x - y) f(y) dy.$$

Aplicamos el ~~teo-fundamental-calculo-generalizado/??~~ con  $X = \mathbb{R}^n$ ,  $\mu$  es la medida de Lebesgue,  $t = x_1 \in [a, b]$  arbitrario, y  $f(y, x_1) = g(x_1, x_2, \dots, x_n - y) f(y) = g(x - y) f(y)$ .

Debemos verificar que  $\left| \frac{\partial}{\partial x_1} [g(x - y) f(y)] \right| \leq g(y) \in \mathcal{L}^1(\mathbb{R}^n)$  donde  $g$  es la mayorante:

$$\left| \frac{\partial g(x - y) f(y)}{\partial x_1} \right| = \left| \frac{\partial g}{\partial x_1}(x - y) f(y) \right| \leq \left\| \frac{\partial g}{\partial x_1} \right\|_\infty \cdot |f(y)| \leq M \cdot |f(y)| \in \mathcal{L}^1(\mathbb{R}^n),$$

donde  $M = \left\| \frac{\partial g}{\partial x_1} \right\|_\infty < \infty$  por hipótesis. Por tanto, obtenemos

$$\frac{\partial}{\partial x_1} \int_{\mathbb{R}^n} g(x - y) f(y) dy = \int_{\mathbb{R}^n} \frac{\partial g}{\partial x_1}(x - y) f(y) dy = \frac{\partial g}{\partial x_1} * f(x) = f * \frac{\partial g}{\partial x_1}(x),$$

donde la última igualdad se sigue de la conmutatividad de la convolución. ■

### **Temporary page!**

L<sup>A</sup>T<sub>E</sub>X was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L<sup>A</sup>T<sub>E</sub>X now knows how many pages to expect for this document.