

# Prop Aut Disco Unidad Traslada Z1 0 Z2 01

**Proposición 1.** Sean  $z_1, z_2 \in \mathbb{D}$  tales que  $z_1 \neq z_2$ , entonces

$$\exists f \in \text{Aut}(\mathbb{D}) : f(z_1) = 0 \wedge f(z_2) \in (0, 1) \subset \mathbb{R}.$$

**Demostración:** Observamos que  $\tau_{z_1}(z_1) = 0$  y, como  $z_2 \neq z_1$ ,  $\tau_{z_1}(z_2) \neq 0$ . Definimos

$$\gamma := \frac{|\tau_{z_1}(z_2)|}{\tau_{z_1}(z_2)} \in \mathbb{T} \quad \text{y} \quad f := \rho_\gamma \circ \tau_{z_1}.$$

Entonces  $f \in \text{Aut}(\mathbb{D})$ ,  $f(z_1) = 0$  y  $f(z_2) = \gamma \tau_{z_1}(z_2) = |\tau_{z_1}(z_2)| \in (0, 1)$ . ■

## Referenciado en

- Teo Camino Minimo Poincare

### **Temporary page!**

L<sup>A</sup>T<sub>E</sub>X was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L<sup>A</sup>T<sub>E</sub>X now knows how many pages to expect for this document.