

Obs Teo Bloch Imp Teo Bloch

Invariante

Observación 1. La versión invariante del teorema de Bloch se puede demostrar a partir del teorema de Bloch. Sea $f \in \mathcal{H}(\mathbb{D})$ tal que $\exists w \in f(\mathbb{D}) : f(\mathbb{D}) \supset D(w, \frac{1}{4} |f'(0)|)$. Entonces, $R(f(\mathbb{D})) \geq \frac{1}{4} |f'(0)|$, lo que implica que $f^\sharp(0) = \frac{1}{2} |f'(0)| \leq 2R(f(\mathbb{D}))$.

Para $a \in \mathbb{D}$, tomamos $f \circ \tau_a$ que cumple $(f \circ \tau_a)(\mathbb{D}) = f(\mathbb{D}) \subset \Omega$ y $(f \circ \tau_a)^\sharp(0) = f^\sharp(a)$. Por tanto, $f^\sharp(a) \leq 2R(f(\mathbb{D}))$, para cada $a \in \mathbb{D}$, y se concluye que $\sup_{z \in \mathbb{D}} f^\sharp(z) \leq 2R(f(\mathbb{D}))$.

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.