

Desigualdad de Hölder

Teorema 1 (Desigualdad de Hölder). Sea $p \in (1, \infty)$ y $f \in \mathcal{L}^p(\mu)$, $g \in \mathcal{L}^q(\mu)$

$$\implies \|f \cdot g\|_1 \leq \|f\|_p \cdot \|g\|_q$$

donde q es el exponente conjugado de p .

Demostración: Si $\|f\|_p = 0$ o $\|g\|_q = 0$, la desigualdad es trivial, por tanto, suponemos que $\|f\|_p \neq 0$ y $\|g\|_q \neq 0$ y definimos $\widehat{f} := f/\|f\|_p$ y $\widehat{g} := g/\|g\|_q$.

$$\implies \|f \cdot g\|_1 = \int_{\Omega} |f(x) \cdot g(x)| \, d\mu(x) = \|f\|_p \cdot \|g\|_q \int_{\Omega} |\widehat{f}(x) \cdot \widehat{g}(x)| \, d\mu(x).$$

Por la desigualdad de Young, $\forall x \in \Omega : \left| \widehat{f}(x) \cdot \widehat{g}(x) \right| \leq \frac{|\widehat{f}(x)|^p}{p} + \frac{|\widehat{g}(x)|^q}{q}$.

$$\begin{aligned} \implies \|f \cdot g\|_1 &\leq \|f\|_p \cdot \|g\|_q \left(\frac{1}{p} \int_{\Omega} |\widehat{f}(x)|^p \, d\mu(x) + \frac{1}{q} \int_{\Omega} |\widehat{g}(x)|^q \, d\mu(x) \right) = \\ &\leq \|f\|_p \cdot \|g\|_q \left(\frac{1}{p} + \frac{1}{q} \right) = \|f\|_p \cdot \|g\|_q. \end{aligned}$$

■

Referenciado en

- prop-inclusion-lp-general
- prop-convolucion-exp-conjugados
- prop-inclusion-lp-esp-finito
- teo-aproximacion-identidad-convolucion
- desigualdad-minkowski

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.