

Ejer localizacion ideal primo anillo local

Ejercicio 1. Sea R un anillo y sea $\mathfrak{p} \subset R$ un ideal primo. Demuestra que $R_{\mathfrak{p}}$ es un anillo local con ideal maximal $\mathfrak{p}R_{\mathfrak{p}}$.

Solución: Queremos probar que $R_{\mathfrak{p}}$ tiene un único ideal maximal que es $\mathfrak{p}R_{\mathfrak{p}}$. Sea $\mathfrak{m} \subset R_{\mathfrak{p}}$ un ideal maximal. Entonces, en particular $\mathfrak{m}$ es un ideal primo, luego por el cor-ideal-primo-localizacion-extendido/??, $\exists \mathfrak{q} \subset R$ ideal primo tal que $\mathfrak{m} = \mathfrak{q}^e$ y $\mathfrak{q} \cap S = \emptyset$. Pero como $S = R \setminus \mathfrak{p}$, tenemos que $\mathfrak{q} \subset \mathfrak{p}$.

Como $\mathfrak{q} \subset \mathfrak{p}$, tenemos que $\mathfrak{q}^e \subset \mathfrak{p}^e$, es decir, $\mathfrak{m} \subset \mathfrak{p}^e$. Además, como $\mathfrak{p} \cap S = \emptyset$, tenemos que $\mathfrak{p}^e \neq R_{\mathfrak{p}}$. Por tanto, $\mathfrak{m} \subset \mathfrak{p}^e \subsetneq R_{\mathfrak{p}}$. Como $\mathfrak{m}$ es maximal, debe ser $\mathfrak{m} = \mathfrak{p}^e = \mathfrak{p}R_{\mathfrak{p}}$. ■

Referenciado en

- teo-extension-entera-ideal-primo-imp-exists-ideal-primo-contrae

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.