

Un ideal y su radical definen la misma variedad algebraica

Ejercicio 1. Sea $I \subset K[x_1, \dots, x_n]$ un ideal. Demuestra que $\mathbb{V}(I) = \mathbb{V}(\sqrt{I})$.

Solución: Veamos ambas inclusiones por separado:

- $\subseteq$ Sea $a = (a_1, \dots, a_n) \in \mathbb{V}(I)$, entonces $\forall p \in I : p(a) = 0$. Sea $q \in \sqrt{I}$, entonces $\exists m \in \mathbb{N} : q^m \in I$ y, por tanto, $0 = (q^m)(a) = (q(a))^m$. Como K es un dominio de integridad, no contiene nilpotentes no nulos y, por tanto, $q(a) = 0$. Luego, $a \in \mathbb{V}(\sqrt{I})$.
- $\supseteq$ Sea $a = (a_1, \dots, a_n) \in \mathbb{V}(\sqrt{I})$, entonces $\forall p \in \sqrt{I} : p(a) = 0$. Como $I \subset \sqrt{I}$, tenemos que $\forall p \in I : p(a) = 0$ y, por tanto, $a \in \mathbb{V}(I)$.

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.